

DoD COVID-19 PRACTICE MANAGEMENT GUIDE

Clinical Management of COVID-19

This Practice Management Guide does not supersede DoD Policy.

It is based upon the best information available at the time of publication. It is designed to provide information and assist decision making. It is not intended to define a standard of care and should not be construed as one. Neither should it be interpreted as prescribing an exclusive course of management. It was developed by experts in this field. Variations in practice will inevitably and appropriately occur when clinicians take into account the needs of individual patients, available resources, and limitations unique to an institution or type of practice. Every healthcare professional making use of this guideline is responsible for evaluating the appropriateness of applying it in the setting of any particular clinical situation. The Practice Management Guide is not intended to represent TRICARE policy. Further, inclusion of recommendations for specific testing and/or therapeutic interventions within this guide does not guarantee coverage of civilian sector care. Additional information on current TRICARE benefits may be found at www.tricare.mil or by contacting your regional TRICARE Managed Care Support Contractor.

Leads: Lt Col Renée I. Matos and COL Kevin K. Chung
6-18-2020

DoD COVID-19 Practice Management Guide v4.0 Summary of Changes

Updated throughout with new literature, studies, and society guidelines across disciplines
Major Updates and Changes listed by Section (sections not listed did not have major changes):

Background & Clinical Presentation: Updated information on clinical presentation, including symptoms, dermatologic findings, MIS-C, hypercoagulable state, and data on cases/CFRs.

Screening & Triage: Recent literature was reviewed for any prospective evaluations of NEWS, and although alternative scales were noted, there was no new data or validation to recommend a change. A recently published individual patient triage algorithm was included as an example.

Infection Prevention & Control: Deleted standard precaution table and added two documents: (1) Mitigating risks of Legionella and other waterborne pathogens, and (2) Recovery plan to consider when reopening services pertaining to infection prevention and control. Reemphasized that N95 used for AGR are not reused per current CDC guidance.

Laboratory Diagnosis: Included information about newer testing options, including salivary testing. Included guidance from IDSA on laboratory diagnosis and algorithms that can be referenced. Expanded section on serologic testing. Changed section title to Laboratory Diagnosis (not just specimen collection).

Prevention of Complications: Updated Acute kidney injury and Hemodialysis epidemiology information; Added Cardiopulmonary Return To Exercise and Physical Activity Recommendations. Updated Anticoagulation recommendations and venous thromboembolism recommendation.

Septic Shock and Cardiac Arrest: Updated Code Blue and RRT algorithms. Added a section on COVID-19 Autopsies in the DoD.

Imaging: Modified recommendations regarding rescheduling non-urgent imaging exams to address the current issues related to safely resuming non-urgent exams. Updated references and added recommendations related to cardiac, neuro and abdominal imaging findings in COVID-19 patients.

Adjunctive Therapies: Updated literature related to the potential benefit of remdesivir and the lack of benefit of hydrochloroquine. Information is included related to how to access remdesivir and indications for and how to access convalescent plasma in the DoD. Updated energy requirements and removed glycerol.

Special Populations: Updates related to inpatient obstetric care to reflect guidance from SMFM and the section was reorganized to improve flow. Updates to the newborn care section to reflect new AAP recommendations. Clinical features of COVID-19 in children has been edited for clarity and organization with additional input from critical care on pediatric shock management. The treatment section deemphasizes hydroxychloroquine. MIS-C is discussed. New recommendations in support of child wellness visits. Geriatrics section updated references.

Surgical Implications: Changed name to reflect inclusion of other invasive procedures. Expanded guidance on preoperative testing and patient education with reference to facility readiness checklist. Revised recommendations regarding intraoperative risk management and the use of engineering controls. Revised subsection on Trauma and Emergency Care. Added subsection on Endoscopy.

Behavioral Health: Added content on (1) assessment and management of delirium and (2) psychopharmacology considerations in COVID-19 treatment.

Rehab Medicine: Added a statement regarding telehealth post-discharge.

Telehealth: Changed the plan for COVID-19 tele-consultation support for Critical Care and ID so the MHS only needs to call 1-833-ADVSRLN (238-7756) or DSN 312-429-9089 to be connected via a live care coordinator to the geographically associated facility as per the DHA memo “COVID-19 Pulmonary/Critical Care and Infectious Disease Consultation”, which is simpler for the end user. Added QR Code that links to the ADVISOR program and added links to the VMC site for references.

EMS: Updated protocols in tables to protect first responders and ensure judicious use of EMS resources during the COVID-19 pandemic. Modified terminology, rationale, and provided specific examples for clarity.

Public Health: Updated guidance for healthcare personnel (HCP).

Ethics: This section was removed. However, the Department of Defense Medical Ethics Center (DMEC) has posted a Bioethics Guidance on the COVID-19 Pandemic (Version 4, 20 April 2020), which can be found at: https://www.usuhs.edu/sites/default/files/media/research/pdf/covid-19_dmec_bioethics_guidance-april_2020.pdf.

Other/Facilities: Information provided regarding installation of Plexiglas sneeze guards/barriers, recommendations for dental area isolation rooms, and water management plans for reopening closed facilities.



Background	4
Clinical Presentation	4
Planning and Preparation	6
Screening and Triage (Early Recognition of Patients with COVID-19).....	11
Immediate Implementation of Infection Prevention & Control (IPC) Measures and Personal Protective Equipment (PPE).....	13
Laboratory Diagnosis of COVID-19	14
Management of COVID-19 Based on Illness Category	16
Outpatient Management COVID-19: Symptomatic Treatment & Monitoring.....	16
Management of Severe COVID-19: Treatment of Co-Infections.....	17
Management of Critical COVID-19: Oxygen and Acute Respiratory Distress Syndrome (ARDS).....	18
Management of Critical Illness and COVID-19: Prevention of Complications.....	23
Management of Critical Illness and COVID-19: Septic Shock and Cardiac Arrest	28
Imaging of COVID-19: Radiology Department Guidance and Imaging Findings.....	32
Adjunctive therapies for COVID-19: Investigational Approaches	36
Caring for Special Populations with COVID-19: Pregnancy and Lactation, Infants, Children and the Elderly	39
Palliative Medicine During the COVID-19 Pandemic	54
Implications of Surgical and Invasive Procedures	57
Operational Considerations for COVID-19: Planning and Preparation	62
Behavioral Health and Wellness in COVID-19 Clinical Management	65
Rehabilitation Considerations for Persons with COVID-19	67
Telemedicine Support During the COVID-19 Pandemic.....	68
Emergency Medical Services (EMS) and Ground Transport of Persons with COVID-19	73
En Route Critical Care Considerations of Persons with COVID-19	77
Public Health Considerations and Response.....	79
Whole of Government Response in Coordination of Resources.....	82
Other Considerations Related to COVID-19.....	82
References	83
Appendix A: Crisis Level Surge Critical Care Triage Tool Example.....	96
Appendix B: Infection Prevention and Control (IPC) Relevant Resources and Documents	97
Appendix C: Example Triage Protocols during COVID-19 Pandemic.....	116
Appendix D: Adult Prone Positioning Protocol Example.....	118
Appendix E: COVID-19 Intubation Checklists, Protocols, and Cognitive Aids	121
Appendix F: Sample Protocol for Extubation of COVID-19 Patients	126
Appendix G: Transport Ventilator Set Up Guide.....	127
Appendix H: Post-COVID-19 Cardiopulmonary Return to Exercise Recommendations.....	129
Appendix I: Weight-Based Heparin Dosing Algorithm for Venous Thromboembolism	131
Appendix J: Enteral Nutrition Care Pathway for Patients with COVID-19.....	132
Appendix K: Sample Protocols for Various Intensive Care Unit (ICU) Management.....	133
Appendix L: American Heart Association (AHA) ACLS and PALS Algorithms for COVID-19 Patients	135
Appendix M: Preparation and Cleaning of Ultrasound Rooms in the Context of COVID-19	137
Appendix N: DHA Summary on Multisystem Inflammatory Syndrome in Children (MIS-C)	140
Appendix O: DHA Quick Reference Guide to Virtual Health and Telephone Encounters	141
Appendix P: Example Triage Protocol for Resource Allocations in Times of Crisis	142
Appendix Q: List of Contributors	146

BACKGROUND

Coronavirus disease 2019 (COVID-19) is a respiratory illness caused by a novel coronavirus (SARS-CoV-2). COVID-19 was first described in Wuhan, China in December 2019 and is now a global pandemic with more than five million confirmed cases worldwide. Most (80%) of those affected have milder illness, 15% will be severely ill (most often some degree of hypoxemic respiratory failure) and 5% will require critical care interventions.(1, 2) Of those who are critically ill, most require early intubation and mechanical ventilation. Other complications include septic shock and multi-organ failure, including acute kidney injury and cardiac injury, in the setting of an inflammatory and prothrombotic state.(3) Older age and comorbid conditions, including hypertension, diabetes, and coronary artery disease increase risk of death.(4-6) The virus is highly contagious and spread via respiratory droplets, direct contact, and if aerosolized, airborne routes.

The intent of this publication is to provide clinicians and military medical treatment facilities (MTFs) with best practices based on latest evidence to optimize DoD response to the current COVID-19 pandemic.

CLINICAL PRESENTATION & CLINICAL COURSE

1. Incubation period: ~4-5 days (interquartile range: 2 to 7 days).(7, 8) Some studies have estimated a wider range for the incubation period. Data for human infection with other coronaviruses (e.g. MERS-CoV, SARS-CoV) suggest that the incubation period may range from 2-14 days; a study of 181 COVID-19 patients supported these initial estimates and found that 97.5% of symptomatic patients develop symptoms within 11.5 days of infection.(9)
2. Frequently reported symptoms of patients admitted to the hospital: (4, 10-16)
 - Fever (77–99%)
 - Cough (46%–82%)
 - Myalgia or fatigue (11–70%)
 - Shortness of breath (SOB) or dyspnea (3-31%)
 - GI symptoms, e.g., anorexia, diarrhea, nausea (pooled prevalence 17.6% in meta-analysis of 60 studies, may precede respiratory symptoms)
 - Anosmia/hyposmia or ageusia/dysgeusia (30-87%)
3. Though fever is ultimately reported in the vast majority of patients during the course of illness, a lower proportion (20-44%) of patients are febrile on presentation. Among 1,099 hospitalized COVID-19 patients in China, fever was present in 44% at hospital admission, though developed in 89% during hospitalization.(8) In a study of 5,700 hospitalized COVID-19 patients in New York City, approximately 31% were febrile on presentation.(17)
4. Less commonly reported symptoms: sore throat, rhinorrhea, conjunctivitis, headache, cough with sputum production and/or hemoptysis, and lower respiratory tract signs and symptoms.(4, 5, 8)
5. Dermatologic findings including maculopapular, urticarial and vesicular lesions (“COVID toes”) and livedo reticularis have been reported in association with illness, although a clear association has not yet been established.(18, 19)
6. Severe disease can occur in all persons, although older patients and those with chronic medical conditions appear to be at higher risk for severe illness.(4, 20) Males appear to be at higher risk for hospitalization and severe illness. (21)
7. Pregnant women: Based on limited data, pregnant women do not appear to be at higher risk for severe disease. Emerging reports from the United States suggest that pregnant women may be at higher risk of atypical presentation with severe disease and caesarean delivery. Additionally, women who develop pneumonia appear to have increased risk of preterm labor.(22-26)
8. Children: Limited information is available about the clinical presentation, clinical course, and risk factors for severe COVID-19 in children with approximately 5-6% presenting with severe illness. In China, COVID-19 made up between 1.5-2% of acute respiratory admissions, with a median age of 7 years. In the US, the median age reported thus far was 11 years. Along with the typical symptoms described, emesis and diarrhea appear to be

Clinical Management of COVID-19, v4

prominent with the virus found in stool samples suggesting fecal-oral transmission. Critically ill children have presented with ARDS, septic shock, encephalopathy and myocarditis. Co-infections with other respiratory viruses or bacteria are common. The MMWR study reported that hospitalized children were more commonly <1 year and had underlying conditions, e.g., asthma.(7, 27-33)

9. Most children experience mild disease, though increasing reports of a syndrome similar to Kawasaki Disease or toxic shock, termed multisystem inflammatory syndrome in children (MIS-C), are evolving. These children may present with persistent fevers, GI symptoms, dermatologic manifestations or lesions or edematous extremities and rapidly progress to shock and multi-organ failure in the setting of known SARS-CoV-2 infection or exposure.
10. Prolonged detection of SARS-CoV-2 RNA has been reported and appears to be related to severity of illness; in respiratory specimens (up to 6 weeks) and stool specimens (>30 days).(27, 28, 34)
11. Clinical presentation among cases of COVID-19 varies in severity from asymptomatic to fatal illness. Several reports suggest clinical deterioration can occur during the 2nd week of illness (range: 5 – 13 days).(4, 12)
12. Acute hypoxemic respiratory failure developed in 17–29% of hospitalized patients. Mortality is high and up to almost 90% in those requiring mechanical ventilation. Secondary infection developed in 10%, with a median time from symptom onset to of respiratory failure of 8 days.(4, 10, 11, 16)
13. Approximately 20-30% of hospitalized patients with COVID-19 and pneumonia have required critical care. Compared to patients not admitted to an intensive care unit (ICU), critically ill patients were older (median age 66 years vs. 51 years), and were more likely to have underlying co-morbid conditions (72% vs 37%).(4, 11)
14. Among critically ill patients admitted to an ICU, 11–64% received high-flow oxygen therapy and 47-71% received mechanical ventilation. A small proportion (3-12% of ICU patients) have also been supported with extracorporeal membrane oxygenation (ECMO).(10, 11, 20)
15. Other reported complications include cardiac injury, sudden cardiac death, arrhythmia, septic shock, liver dysfunction, acute kidney injury, venous and arterial thrombosis despite chemoprophylaxis, and multi-organ failure.(35)
16. COVID-19 is associated with a hypercoagulable state. A Dutch review of COVID-19 positive patients admitted to an ICU with pneumonia revealed 31% experienced a thrombotic complication with the majority of these being pulmonary emboli.(36) Viral inclusion bodies have been seen in endothelium of kidneys, small bowel, and heart suggesting that endotheliopathy could be contributing to thrombotic complications.(37) The prevalence of arterial thrombosis such as stroke is not as well described as the significantly increased risk of venous thromboembolism.
17. Case fatality rates (CFR) appear to vary by location and be related to demographics, e.g., median age, of the population. A CFR of 2.3% has been reported among confirmed cases of COVID-19 in China.(20) However, the majority of these cases were hospitalized patients, so this mortality estimate is likely biased upward. Among hospitalized patients with pneumonia, the case fatality proportion has been reported as 4–15%.(4, 10, 11) In a report from one Chinese hospital, 61.5% of critically ill patients with COVID-19 had died by day 28 of ICU admission. Among all critically ill COVID-19 patients in China, the reported case fatality proportion was 49%.(3)
18. As of 29 May 20, the Italian government COVID-19 surveillance group reported 31,936 deaths associated with COVID-19, of which 85.1% ≥70 yr, 10.3% 60-69 yr, 3.5% 50-59 yr, 0.9% 40-49 yr, 0.2% 30-39 yr, 0.04% <30 yr. Of the 2,621 patients for whom data on pre-existing co-morbidities are available, approximately 60% had >3 pre-existing co-morbidities (e.g., hypertension, type 2 diabetes, ischemic heart disease, atrial fibrillation). The CFR for Italy was estimated at 13.7%. <https://www.epicentro.iss.it/en/coronavirus/>
19. In the US, as of 1 Jun 20, the Centers for Disease Control and Prevention (CDC) reports approximately 1.8 million cases and 104,396 COVID-19-related deaths. (<https://www.cdc.gov/coronavirus/2019-ncov/cases-updates/cases-in-us.html>) CFR increases with age (highest in ≥85 yr). As of 23 May, the overall cumulative hospitalization rate is 73.3 per 100,000, with highest rates in people ≥65 yr (229.7 per 100,000) and 50-64 yr (113.4 per 100,000). (<https://www.cdc.gov/coronavirus/2019-ncov/covid-data/covidview/index.html>)




Figure 1. Clinical Courses of Major Symptoms and Outcomes and Duration of Viral Shedding [from Zhou, et al.; Lancet (2020)]. (5) Figure shows median duration of symptoms and onset of complications and outcomes. ICU, intensive care unit; ARDS, acute respiratory distress syndrome.

PLANNING AND PREPARATION

Facility Incident Command and Systems.

1. A local emergency response command structure with clearly defined roles and lines of communication should be defined. (38, 39) These structures should have the ability to coordinate expansion or restriction of resources in conjunction with unit medical directors, help coordinate “just in time” training as well as regional expert consultation (i.e. tele-consultation with critical care, infectious disease, or other specialists), facilitate the flow of staff, critical equipment and patients, and coordinate with Contingency the Crisis Standard of Care (CSC) changes on both a local and regional level. Additionally, the local Incident Command Center (ICC) should liaise and coordinate with the community as transition through crisis care levels and resource triage if needed, depends on regional, not just local, healthcare utilization.
2. Establish and Manage Crisis/Contingency Standards of Care
 - a. Crisis Standards of Care are “a substantial change in usual healthcare operations and the level of care it is possible to deliver, which is made necessary by a pervasive (e.g., pandemic influenza) or catastrophic (e.g., earthquake, hurricane) disaster.” (40)
 - b. The establishment of CSC should enable specific legal and regulatory protections for health care providers. For reference, DODI 6200.03 allows for establishment of a CSC within the DoD.
 - c. Design and implementation of these standards for each agency should remain flexible based on each situation and should be tiered (i.e. normal operations, contingency, crisis) and have specific triggers to engage. In general Contingency when >120% typical capacity and Crisis when >200% capacity.
 - d. Contingency Care is more similar to typical care standards with most staff working in their usual environments but with expanded clinical responsibilities.
 - e. Crisis Standards of Care, if invoked, triggers significantly altered staffing models as described below with incumbent assumed risks of increased morbidity and mortality. CSC should be developed by multi-disciplinary groups and collated by the Incident Command Center (ICC) and should be individualized to a facility. A list of topics that should be included:
 - Authority and triggers for enacting escalating from usual to Contingency then Crisis
 - Just in time training & scope of practice changes as CSC escalate (nursing, physician, etc)
 - Alterations in practice allowed (limiting documentation, changes in work hours and locations, changes in location of patient care and monitoring requirements).

Clinical Management of COVID-19, v4

- Alternations from normal should be limited as much as possible to mitigate patient safety risks.
3. Establish clear lines of communication (LOC) to ensure:
 - a. The ability to communicate updated processes and protocols.
 - b. The ability to transfer clinical information with patients through the system.
 - c. That communication be consistent, from designated sources, and information be trusted by staff.(41-43)
 4. Establish Patient Tracking and Re-unification systems: Plan and coordinate a system for patient tracking, identification, and the ability to communicate with next of kin who may be restricted from visitation.(43)
 5. Establish security, access points, and “clean” areas with access restricted:
 - a. Security should be included in the planning process given increased community stress and security risks during the COVID-19 pandemic.
 - b. Establish “satellite” units in alternative locations to care for patients unaffected by the pandemic to protect non-infected patients and high-risk staff (e.g., underlying medical conditions, age >60).(44)
 - c. Consider access to specialty or routine care that may be needed in these areas with screening as patients enter.
 - d. Establish single or controlled points of entry for every facility and initiate screening procedures for possibly infected patients at entrances.
 6. Coordination of re-prioritization of clinical duties:
 - a. Focus on urgent care, but ensure a process for providing necessary routine care when unsafe to defer.
 - b. Care should be primarily virtual unless a face-to-face visit is necessary as determined by the care team.
 - c. Closely track access and demand and consider expanding or contracting services based on local epidemiology and need.
 - d. Coordinate re-allocation of assets off loaded by limitations to areas of need (Critical Care, Inpatient care, Initial triage, and Urgent/Emergency Care).(45)
 - e. Limit administrative, educational and academic duties to those necessary to directly support patient care.
 - f. Frequently message patients and staff any changes in services, clinic hours, entry procedures, etc. to manage their expectations.




Figure 2. A framework outlining the conventional, contingency, and crisis surge responses. PACU: post-anesthesia care unit. [from Christian, et al.; Chest (2014)].(46)

7. Develop Recall Roster for all assets (nursing, physician, housekeeping, dietary, security, admin, etc) and triggers for re-calling those who may be needed from remote work.
8. Consider logistic/ancillary support needs when determining “Essential Personnel” for tasks including:
 - a. Disposal of personal protective equipment (PPE) and cleaning both “dirty” rooms and shared spaces. These tasks should be prioritized and will be in very high demand.(47)
 - b. Allocation of adequate space for safe, respectful care of the deceased.(48)
 - c. Designating locations and facilities to shelter and feed families of ill patients, staff members, and even families of staff members to augment and limit the up to 40-50% absenteeism anticipated with illness, school/childcare closure, and fear.(44, 45)

Preparing Critical Care Resources & Teams.

1. Understand the following steps provide a framework and are not the “correct” way to manage bed or staffing expansion. Exact staffing models, ratios, logistic and system support models should reflect the needs of the community and resources available at local centers. Transitioning to Crisis Care models carries with it significant increases in both morbidity and mortality above that seen in standard care models. It should be undertaken only when absolutely necessary, with careful consideration, and in an iterative way assessing for increased volume paradoxically leading to excessively increased morbidity and mortality.
2. **Staffing.** In a global pandemic causing a surge of emergency room and admitted patients, additional staffing models should be considered. Although telehealth resources should be optimized, there may still be significant deficits in critical care trained healthcare workers.
 - a. Staff Shortages:
 - i. Illness, fatigue, fear, and care giver duties, particularly with school/daycare closure, limit staff availability with some estimates as high as 40-60% absenteeism.(44, 49)
 - ii. Augmenting staffing initially with increased “mandated overtime” should be avoided as long as possible to avoid early staff burn out.
 - iii. Facility based alteration of staffing ratios (i.e. less provider staff in the inpatient setting overnight) may help reduce staff burden while maintaining reasonable coverage in keeping with typical hospital processes.
 - iv. Strategies listed above may mitigate (facility based child care, cohort care teams, etc.) but planning should consider at least a 25-40% reduction in staff availability. Additional recommendations to augment staff availability include:(50)
 - A PPE officer (can be trained non-clinical staff) to train and monitor PPE and staff exposure on each ward
 - Mental health support or “resiliency teams” with focus on staff wellness and support
 - Team “Safety officers” to monitor/ensure breaks, hydration, toileting and nutrition
 - v. **Critical Care.** The Society of Critical Care Medicine (SCCM) recommends staffing models to support expanded critical care bed capacity in the event of a global pandemic, which includes use of multiple non-ICU trained healthcare workers. Another model with a single ICU physician leading a larger team caring for more critically ill patients is advocated by some but would likely lead to even higher increases in excess mortality. At a minimum, the first four staff positions noted below should be ICU trained and experienced: (51)
 - Critical Care Physician
 - Respiratory Therapist
 - Advanced Practice Providers (APP)
 - Critical Care Nurse (CCRN or experienced active RN working in critical care)
 - In facilities without intensivists, critical care teams may be directed by anesthesiologists, pulmonologists, hospitalists, or others with experience caring for critically ill patients.(51)
 - Staffing for the other roles could include but are not limited to those with some previous critical care training or experience who currently work as:
 - Non-ICU physician: anesthesiologists, hospitalists, general surgeons or others with experience caring for critically ill patients

Clinical Management of COVID-19, v4

- CRNA, CAA, MD/DO: Residents from medical or surgical specialties (with appropriate supervision and graduated responsibility) or other medical or surgical staff preferably with experience in inpatient medicine
- Non-ICU nurse tiered from best to least suited:(50)
 1. RN currently working in progressive care units (telemetry or step down units)
 2. Ambulatory care setting with previous ICU experience (preferably within 3 years)
 3. Paramedics, EMTs or RNs and medical assistants/LPN that work in urgent care




Figure 3. SCCM Tiered Critical Care Staffing Strategy for Pandemic. APP: advanced practice provider; RT: respiratory therapist; CRNA: certified registered nurse anesthetist; MD/DO: physician [modified from SCCM link above].(51)

- vii. **Step-down Care/Intermediate Care Ward (ICW).** Figure 4 provides a framework staffing model for patients requiring more intensive support but not mechanical ventilation/vasopressor support, or those at imminent risk of requiring mechanical ventilation/vasopressor support, such as could be managed in a step-down unit. Ideally, this team would be led by an experienced hospitalist or intensivist who oversees the care of physician-led teams. These staffing models would be supported by a minimum of two teams working no longer than 12-hour shifts. (44) In the setting of COVID-19, these are likely patients that would be hospitalized in fixed facilities not in ICUs.




Figure 4. Tier 2 Staffing Strategy for Step-down Level Care during a Pandemic

- viii. **Routine Inpatient/Ward Care.** Figure 5 provides a framework staffing model for inpatient routine medicine care, with the team led by an experienced hospitalist or physician with hospital experience. In the setting of COVID-19, these would likely be patients housed in “off-site” facilities with limited resources (e.g., tents, gyms, convention centers, etc).




Figure 5. Tier 3 Staffing Strategy for Routine Ward Level Care during a Pandemic

vii. **Pediatric Care.** For MTFs that have a large footprint of pediatric providers (pediatric residencies, pediatric intensivists, pediatricians, pediatric nursing), there should be consideration to flex pediatric age range up to 30 years old in the Contingency Stage and higher as needed in Crisis Standards of Care. This will leverage appropriate expertise to care for young adults, which is common both for these providers especially in the military, and offload patient numbers from the adult care teams. For smaller MTFs that have minimal pediatric beds, minimal pediatricians (i.e., Family Practice caring for children), there should be consideration of diverting inpatient pediatric patients to dedicated children's hospitals. This decision should be made based on available community capacity and there should be communication with local facilities to strategically plan for patient distributions. MTFs must still maintain dedicated non-COVID-19 medical missions, and should not sacrifice care in other areas (e.g., use NICU beds/ventilators for adult patients if needed in the NICU).

b. **Privileging Options.** In accordance with national standards for accreditation, local leadership may cross-level providers to provide patient care, treatment and services necessary as a life-saving or harm reducing measures, provided the care, treatment, and services are within the scope of the individual's license without modification of existing privileges. Disaster privileges can only be granted to volunteer licensed independent practitioners when the organization's Emergency Operations Plan has been activated. During emergencies, providers undergoing "just in time" training for work outside their normal areas may work within the scope of their individual licensure and do not require privilege modification, addition or supervision. Privileging authorities may award disaster privileges on activation of their emergency management plans consistent with provisions established in DHA PM 6025.13, Volume 4.

2. **Staff Training.**

- a. ICU "Just in time training" for augmentees are available at <https://www.sccm.org/covid19>, <https://www.chestnet.org/Guidelines-and-Resources/COVID-19/On-Demand-e-Learning>, <https://www.chestnet.org/Guidelines-and-Resources/COVID-19/Updates-and-Resources>, <https://www.sccm.org/disaster>, and <https://covid19toolbox.com>.
- b. **Training and augmentation platforms.**
 - If local expertise is not available, utilization of existing DHA teleconsultation platforms (PATH, ADVISOR) may augment capabilities.
 - Places with ICU care should develop brief local ICU orientation models focusing on safety practices, unit hierarchy, protocols, and consultative relationships (brief, max 4-8 hours).
 - Training platforms for provider and nursing augmentees should focus on remote learning resources to provide baseline didactic training such as those above or those locally developed.
- c. Critical care considerations for pregnant women online training is available at: <https://www.smfm.org/critical-care/cases/new-2019>.
- d. DHE Clinical RN Refresher Training Packet was released with the intent of helping to refresh inpatient nursing experience. (<https://info.health.mil/edu/Pages/COVID.aspx>)

Clinical Management of COVID-19, v4

- e. PPE; Donning and doffing officers, which can be personnel pulled from non-clinical roles (administrators, support staff, etc.) should be assigned to train and monitor compliance with PPE protocols. Training video: <https://www.youtube.com/watch?v=bG6zISnenPg> (52)
3. **Equipment and Consumables.** Daily assessment of ventilators, ventilator circuits, PPE, fluids, sedating and other critical medication and supplies should be tracked with equipment burn rates estimated and updated as information is available.
 - a. Consider creating intubation/procedure packs with all necessary equipment and supplies to avoid going in and out of the room repeatedly.
 - b. Consider alternative options to reduce and re-use critical items such as PPE and ventilator circuits. Encourage sharing local policies and solutions as they become available.
 - c. Consider utilization of anesthesia ventilators during expansion, but ensure some remain in reserve based on facility needs for acute, non-COVID-19 emergencies.
 - d. Inventory management.
 - Develop a list of key inventory to include PPE, ventilators and supporting equipment, fluids, key medications, fluids, nutrition, IV and other vascular access supplies, etc.
 - Assign personnel to develop a tracking system and publish regularly to ICC. Utilize models to predict inventory burn rate. A sample PPE utilization calculator available at: <https://www.chestnet.org/Guidelines-and-Resources/Resources/Essential-Institutional-Supply-Chain-Management-in-the-Setting-of-COVID-19-Pandemic> (53)
4. **Space:**
 - a. **ICU Contingency Units.** Many modern ICUs have rooms capable of expanding to hold two patients. These spaces need to be assessed to house appropriate ventilators, suction, and monitoring, but if so equipped, should be utilized first. Co-locating COVID-19 patients as much as possible will increase the efficiency of staff and supply use. If these spaces are exhausted, other monitored, ventilator capable areas may be available to use as alternative ICU rooms (OR, PACU, etc).
 - b. **Ward Cohorting:** Consideration should be given to establishing COVID-19 wards. Clean barriers on open units similar to chemical “hot lines” can be used. This includes cohorting staff to “COVID-positive” or “COVID-negative” teams based on which cohort they are caring for to reduce transmission. If possible, COVID-19 inpatient care should be limited to specific areas of the hospital with designated travel routes reserved for flow of COVID-19 positive patients.

Establishment of a DoD Case Registry for Clinical Performance Improvement.

1. Systematic collection and iterative analysis of key clinical data is essential to optimize delivery of care.
2. The registry currently being implemented will support performance improvement in the setting of a learning health system.
3. Standardized electronic health record (EHR) templates have been developed to increase harmonization and completeness of important data elements needed for the registry.

Returning to the “New Normal”

The decision to de-escalate from contingency and crisis care should be governed by similar principles with ICC coordination, triggers for phased de-escalation, and clear communication. The risk of prolonged delay in routine care or altered practice models creating urgent or emergent care needs and increasing morbidity and mortality should be considered as the decision of when/how to transition back to more normal care models. Additionally, institutions should recognize and plan for a prolonged period (months or longer) with low level COVID-19 care needs requiring cohorted outpatient, emergency, and inpatient services as much as possible to avoid Healthcare associated spread. Plans should be in place with clear triggers to re-escalate to contingency or crisis care with the relaxing of social distancing.

SCREENING AND TRIAGE: EARLY RECOGNITION OF PATIENTS WITH COVID-19

1. **Screening:** Screen and isolate all patients with suspected COVID-19 at the first point of contact with the health care system (ER/clinic/drive-through screening/labor and delivery). Establish processes for how to handle people screening positive at entrances. Processes should be clear and easy to follow and be standardized

Clinical Management of COVID-19, v4

across facilities within the Local Command. It is also recommended to direct low-risk patients to drive-through screening facilities as available to reduce exposure and conserve PPE in MTFs.

2. **Triage:** Triage patients using standardized triage tools and initiate the appropriate disposition decision depending on the clinical setting. Ensure standard protocols established in cooperation with Infectious Disease and Public Health that are clear and easy for staff to follow. Try to keep protocols aligned with national (CDC) and local (state or municipal) guidance and update regularly as new guidance emerges. Triage should be conducted telephonically or in a designated outdoor or dirty area when possible. Staff evaluating patients face-to-face should be pre-identified and outfitted and trained on appropriate PPE. Patients can pre-screen themselves using available self-checkers from the CDC and other organizations.
 - a. **A potentially useful tool for initial categorization of clinical severity and aiding in triage is the National Early Warning Score (NEWS),** Figure 6. This clinically derived score is easily measured in a triage area, clinic, emergency department or other initial assessment environment and consists of parameters listed below.
 - b. The score ranges from 0-21 and higher scores have been demonstrated to correlate with worsened mortality. A score of above 5 increases the likelihood of eventual ICU level of care.(54)
 - c. NEWS in COVID-19 has distinct advantages over qSOFA which can underestimate the severity of presentation if confusion, and hypotension are absent as they often are in COVID-19 patients.(55)
 - d. An alternate version of the NEWS has been developed in China incorporating age >65 as a risk factor in the scoring. The development of this is based on different scoring system for retrospectively created on patient data from the 2013 Avian Influenza epidemic,(56) and neither this score, nor the alternate NEWS score have been prospectively validated/evaluated.(57)
3. **Initial treatment of hospitalized inpatients** consists of optimized supportive and symptomatic care in the ward or intensive care unit. Patients with increased risk of severe disease and mortality include:
 - Age >60
 - Diabetes mellitus
 - Hypertension
 - Immunosuppression
 - Cardiopulmonary disease
4. Patients may present with mild symptoms but have high risk of deterioration and should be admitted to a designated unit for close monitoring.
 - a. Additional consideration should be given to a patient’s resource level in their residence (e.g., barrack dwellers), and ability to quarantine and self-monitor when deciding to admit or discharge a mildly symptomatic patient.

PHYSIOLOGICAL PARAMETERS	3	2	1	0	1	2	3
Respiration Rate	≤8		9 - 11	12 - 20		21 - 24	≥25
Oxygen Saturations	≤91	92 - 93	94 - 95	≥96			
Any Supplemental Oxygen		Yes		No			
Temperature	≤35.0		35.1 - 36.0	36.1 - 38.0	38.1 - 39.0	≥39.1	
Systolic BP	≤90	91 - 100	101 - 110	111 - 219			≥220
Heart Rate	≤40		41 - 50	51 - 90	91 - 110	111 - 130	≥131
Level of Consciousness				A			V, P, or U

Figure 6. National Early Warning Score (NEWS).(54)

Clinical Management of COVID-19, v4

- Mild Illness.** For mild illness, hospitalization may not be required unless concern about rapid deterioration. Isolation to contain/mitigate virus transmission should be prioritized. Safe home care can be performed according to CDC guidance (<https://www.cdc.gov/coronavirus/2019-ncov/hcp/guidance-home-care.html>).
- ICU Admission Criteria.** ICU admission and exclusion criteria may be a fluid decision based on the facility. Given that allocation of dedicated ICU beds and surge capabilities amongst individual hospitals are variable, each hospital should provide a specific plan regarding ICU admission/exclusion criteria. This could be based on the percentage of resources utilized (e.g., beds, ventilators). Figure 7 provides an example plan. Individual triage decisions could be made on the basis of a composite of factors including likelihood of recovery, pre-existing functional status, and severity of illness. An example triage schema is shown in **Appendix A.**(58)




Figure 7. Example of an ICU Surge Plan (from the San Antonio Veteran’s Affairs Hospital)

IMMEDIATE IMPLEMENTATION OF APPROPRIATE Infection Prevention Control (IPC) MEASURES/PPE

- All employees working in inpatient units, ambulatory clinic spaces, and procedural areas should wear a surgical face mask, at all times, while in their respective clinical care settings. Such requirements align with current policies from leading healthcare systems, including the University of Nebraska. Additional information can be found at the following link: <https://www.nebraskamed.com/sites/default/files/documents/covid-19/surgical-mask-policy-and-faq-nebraska-med.pdf>. See **Appendix B** for PPE donning/doffing and mask use.
- Appropriate use of PPE plays an important role in the prevention of disease transmission, however ensuring appropriate work practice and environmental controls are in place is critical. In addition to implementing the PPE guidelines provided in Figure 8, MTFs should adhere to the following essential practices:
 - Screen all visitors and healthcare workers before entry into the MTF (i.e., inside as they enter).
 - Implement restricted visitation policies for the facility (refer to example provided by Emory Healthcare: <http://www.emoryhealthcare.org/covid/index.html>, no federal endorsement is intended or implied)
 - Practice social distancing
 - Adhere to frequent hand hygiene and wear a surgical or cloth mask at all times (includes visitors).
 - Surgical masks are preferred over cloth masks for healthcare personnel. Consider continuing to wear

Clinical Management of COVID-19, v4

respirator or facemask (extended use) while in the facility instead of intermittently switching back to cloth face covering which could cause self-contamination.

- f. Limited re-use of N95 Respirators refers to practice of using same respirator by one HCP for multiple encounters with different patients but removing after each encounter. If no manufacturer guidance is available data suggest limiting the number of reuses to no more than five uses per device. (23 April) <https://www.cdc.gov/coronavirus/2019-ncov/hcp/respirators-strategy/index.html#crisis>

CATEGORY	DEFINITION	REQUIRED ISOLATION/PPE
0	Patient not suspected of having COVID-19	STAFF: <ul style="list-style-type: none"> • Surgical mask • PPE according to task. <i>See Standard Precautions.</i>
		PATIENTS: Masking (e.g., cloth, surgical mask)
1	Asymptomatic patient with known exposure to COVID-19 OR Traveled from high-risk areas within last 14 days	STAFF: <ul style="list-style-type: none"> • Surgical mask • PPE according to task. <i>See Standard Precautions.</i>
		PATIENTS: MUST wear surgical mask if traveling outside room for medically essential purposes
2	Patient under investigation (PUI) or positive COVID-19	STAFF: <ul style="list-style-type: none"> • Contact Precautions (gown and gloves) • Droplet Precautions (surgical mask appropriate per current CDC guidelines if no aerosol-generating procedures performed in room) • Eye protection (face shield or goggles)
		PATIENTS: MUST wear surgical mask if traveling outside room for medically essential purposes
3	Positive COVID-19 requiring aerosol-generating procedures (i.e., BiPAP, CPAP, endotracheal intubation, high-flow nasal cannula, nebulizers, tracheal suctioning)	STAFF: <ul style="list-style-type: none"> • Contact Precautions (gown and gloves) • Consider head and foot covers • Airborne Precautions (N95 Respirator or PAPR) • Eye protection (face shield or goggles) • Negative pressure room
		PATIENTS: MUST wear surgical mask if traveling outside room for medically essential purposes

Figure 8. PPE Recommendations for the MHS (Adapted for the MHS using CDC guidelines accessed 24 April 2020:

<https://www.cdc.gov/coronavirus/2019-CoV/hcp/index.html>); PAPR, powered air-purifying respiratory; PPE, personal protective equipment; PUI, patient under investigation; High-risk Area - Areas with a Level 3 Travel Health Notice identified by the CDC Visit www.cdc.gov/coronavirus/2019-ncov/travelers/after-travel-precautions.html for current list.

Special Situations: ED staff and outpatient healthcare workers with any patient encounter with a PUI: Follow Category 2.

- 3. **PPE Visual for Use During Supply Shortages:** The following visuals from Emory University are available as printable PDFs: <https://med.emory.edu/departments/medicine/divisions/infectious-diseases/serious-communicable-diseases-program/covid-19-resources/conserving-ppe.html>
- 4. **Questions related to IPC** can be sent to: dha.ncr.clinic-support.list.ipc-group@mail.mil

LABORATORY DIAGNOSIS OF COVID-19

- 1. **Triage:** Patients should be triaged and initial testing optimally performed in a manner separated from the general patient population such as in a tent or designated area within a facility. Initial laboratory collection will include nucleic acid testing for SARS-CoV2 RNA and additional tests as indicated.
- 2. **Molecular Testing (PCR and other nucleic acid amplification tests [NAAT]):** Molecular testing is the current gold standard for making the diagnosis of COVID-19. Diagnosis of COVID-19 is primarily confirmed by nucleic acid amplification techniques (NAAT) designed to detect SARS-CoV-2. The most common of these are real-time

Clinical Management of COVID-19, v4

reverse transcriptase polymerase chain reaction (RT-PCR) tests, although other NAATs (in-situ hybridization and others) are also used. The initial CDC PCR assay received emergency use authorization (EUA) on March 3, 2020. Since then numerous commercial laboratories and universities have developed assays with subsequent FDA authorization.(59) These assays are highly specific, but sensitivity may depend on the disease process (mild upper respiratory infection vs severe pulmonary disease), the specimen site, and quality of specimen collection. Consider retesting if clinical suspicion for COVID-19 remains high. Sensitivity of SARS-CoV-2 NAATs may also vary depending on the nucleic acid target being used. The WHO assay uses RNA-dependent RNA polymerase, whereas the CDC assay targets genes which target viral nucleocapsid genes (N genes). One non-peer reviewed publication suggested that the nucleocapsid assays have higher sensitivity than the WHO assay. As a result, any data on sensitivity from a NAAT performed under the WHO assay (to include reports from China) may underestimate sensitivity of assays in the United States.(60)

3. **Specimen Collection for NAAT:** Specimen collection has largely been via nasopharyngeal (NP) samples which should be collected via synthetic fiber swabs. Other specimen types, to include oropharyngeal swabs, sputum, endotracheal tube aspirates, bronchoalveolar lavage (BAL), blood and stool may be utilized, depending on disease process. According to the CDC, nasopharyngeal, oropharyngeal, or a combination of both are commonly used. Other options include mid-turbinate swabs, anterior nares swabs, and nasal washing. Nasal washing may lead to increased risk of aerosolization and could increase the risk of infection to HCPs. If unable to collect from the URT, specimens from the lower respiratory tract (LRT) using expectorated sputum or endotracheal aspirate may be indicated in hospitalized patients with lower respiratory tract infection. Bronchoalveolar lavage (BAL) may be required for sampling, but increases the risk to healthcare providers due to aerosolization of virus.(61) Testing for other viral infections such as influenza should be obtained if indicated, depending on local epidemiology of respiratory viruses. One study describing 1070 specimens from 205 patients with COVID-19 suggested that LRT samples were most likely to be positive for viral RNA.(62) In three studies including patients with specimens collected at multiple sites URT samples in patients with SARS-CoV2 detection at one or more sample sites have a have an estimated sensitivity of 76%.(63) Several recent studies compared NP to salivary samples.(64-69) High concordance between NP and saliva samples were reported, with some studies reporting more frequent detection of SARS CoV-2 RNA in salivary vs NP swabs; however these findings were not consistent between studies, and collection method (spitting vs “coughing up” saliva) also varied. The Infectious Diseases Society of America (IDSA) issued guidelines for recommended specimen types in symptomatic and asymptomatic patients, to include recommendations for self-collection of nasal and mid-turbinate swabs by symptomatic patients, and does not recommend use of oropharyngeal or salivary samples alone.(63)
4. **Antigen Testing:** Antigen testing is designed to detect SARS-CoV-2 proteins in a rapid format, without the complexity of a molecular test. Although less sensitive than NAAT, antigen testing is rapid and specific – producing high positive predictive values (PPV), but low negative predictive values (NPV). This allows for the rapid detection of SARS-CoV-2 antigen in upper respiratory samples, but at the cost of more false negatives. The first antigen test received FDA authorization May 8, 2020.(70) Patients with suspected COVID-19 who test positive with these assays, should be considered to have confirmed infection. Those who test negative should be retested with a molecular test.
5. **Serologic testing:** The majority of immunocompetent hosts will develop antibodies (IgG and/or IgM) 2-3 weeks after the onset of symptoms associated with COVID-19. Serologic assays have been developed to assess antibody response to COVID-19 infection. The two major antigenic targets of SARS-CoV-2 virus against which antibodies are detected are spike glycoprotein (S) and nucleocapsid phosphoprotein (N), although other assays detect antibodies to different antigenic sites.(71) These assays include both point-of-care tests, (which use lateral-flow technology), or laboratory tests, which use ELISA (Enzyme-Linked Immunosorbent Assay) or CIA (chemiluminescent immunoassay). Neutralizing assays have not yet been approved by the FDA. FDA provided EUA for the first antibody tests in April 2020. Unfortunately, the market was rapidly flooded by over 70 assays, which were not evaluated or tested by the FDA. Many of these produced high rates of false-positive (likely secondary to cross-reactivity to other coronaviruses) and/or false-negative results (as the development of antibody response is inconsistent). This led to the Infectious Diseases Society of America (IDSA) advising against use of serologic testing on April 22, 2020.(72) The durability of antibody response in Covid-19 is not currently known. To warn of this knowledge gap, the World Health Organization (WHO) issued a statement

Clinical Management of COVID-19, v4

April 24, 2020 warning that prior infection with SARS-CoV-2 has not been proven to confer immunity to reinfection.(73) The Positive Predictive Values (PPV) of serologic assays vary depending on the prevalence of disease in a given population and number of tests obtained, ranging from as low as 27% PPV for a single test in a patient with 2% community prevalence, to 89% in a community with 30% prevalence.(74)

6. **Retesting Persons with Covid-19:** In persons with confirmed COVID-19, repeated URT and LRT samples can be collected to demonstrate viral clearance. Initially, the CDC recommended using 2 negative tests (greater than 24 hours apart, obtained after clinical recovery) to document virus clearance. Emerging data has documented persons testing positive by molecular methods (using upper respiratory specimens) for up to 8 weeks. Other limited viral culture data supports a much shorter course of shedding viable virus (7-9 days). Thus, repeat testing to “clear” patients may lead to prolonged enhanced isolation in the hospital, difficulty in placing clinically improved patients, and delay in return to work in otherwise healthy individuals. CDC currently has recommendations for both symptom-based and test-based strategies for discontinuation of transmission-based precautions.(75)
7. **Personal Protective Equipment (PPE):** Use appropriate PPE for specimen collection (droplet and contact precautions for URT specimens; airborne precautions for LRT specimens).
8. **Pre-operative/pre-procedural testing:** assessing active COVID-19 infection to determine subclinical infection in patients who require invasive or non-invasive ventilation for surgical or other procedures should generally be limited to NAAT-based assays. Use of serologic assays to determine risk of infectivity to HCPs or to consider reduction of PPE is not recommended. The IDSA guidelines on diagnosis of COVID-19 provide additional guidance on preoperative and pre-procedural testing.(63)
9. **For pregnant and recently postpartum patients:** COVID-19 testing of symptomatic women may need to be prioritized due to need for inpatient care with delivery and ongoing outpatient visits, to enable access to specialized care, to allow appropriate maternal PPE, and appropriate care for the newborn.
10. **Co-infection:** Dual infections with other respiratory viral and bacterial infections have been found in SARS, MERS and COVID-19 patients. As a result, a positive test for a non-COVID-19 pathogen does not rule out COVID-19. At this stage, detailed viral and microbiologic studies are needed in all suspected cases.

MANAGEMENT OF COVID-19 BASED ON ILLNESS CATEGORY

Per National Institutes of Health (NIH) COVID-19 Treatment Guidelines, in general, patients with COVID-19 can be grouped into the following illness categories:(76)

- **Asymptomatic or Pre-symptomatic Infection:** Individuals who test positive for SARS-CoV-2 but have no symptoms
- **Mild Illness:** Individuals who have any of various signs and symptoms (e.g., fever, cough, sore throat, malaise, headache, muscle pain) without shortness of breath, dyspnea, or abnormal imaging
- **Moderate Illness:** Individuals who have evidence of lower respiratory disease by clinical assessment or imaging and a saturation of oxygen (SaO₂) >93% on room air at sea level.
- **Severe Illness:** Individuals who have respiratory frequency >30 breaths per minute, SaO₂ ≤93% on room air at sea level, ratio of arterial partial pressure of oxygen to fraction of inspired oxygen (PaO₂/FiO₂) <300, or lung infiltrates >50%.
- **Critical Illness:** Individuals who have respiratory failure, septic shock, and/or multiple organ dysfunction.

OUTPATIENT MANAGEMENT OF COVID-19: SYMPTOMATIC TREATMENT AND MONITORING

1. **Overall management:** The mainstay of treatment for mild cases of COVID-19 is supportive care.
2. **Disposition:** Those with mild or moderate disease may be managed as an outpatient. Moderate cases should be considered for admission for close observation due to the risk of rapid pulmonary disease progression. The determination of outpatient vs inpatient care should be individualized based on consideration of symptom severity, risks for adverse outcomes (e.g., underlying illness and age), and the patient’s social context:
 - a. Their access to resources such as food and other necessities for daily living

Clinical Management of COVID-19, v4

- b. Their access to appropriate caregivers or ability to engage in self-care
 - c. Their ability to engage in symptom and public-health monitoring
 - d. The transmission risk within the home (e.g., the availability of a separate bedroom to minimize sharing of immediate living spaces; their access to PPE such as gloves and a facemask; their ability to adhere to home isolation, respiratory and hand hygiene, and environmental cleaning; and household members at increased risk for COVID-19 complications).(20, 77, 78)
3. **Monitoring for symptomatic progression:** Monitoring for the evolution of symptoms may be conducted by clinical staff or public-health personnel, depending on local policy.
 - a. Although 81% of patients in a Chinese case series had mild symptoms, those who progressed to more severe disease were hospitalized a median of 7-11 days after the onset of illness.(1, 5, 10) Therefore, close monitoring for symptomatic progression through the second week of illness is important for non-hospitalized patients.
 - b. Close monitoring should be emphasized in any patient who is identified as being at higher risk for severe illness per CDC guidelines at <https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/people-at-higher-risk.html>. Monitoring via telehealth may be an option for these patients.
4. **Home care guidance:** Healthcare providers may provide patients and caregivers with available CDC guidance on home care:
 - a. Preventing the Spread of Coronavirus Disease in Homes and Residential Communities (<https://www.cdc.gov/coronavirus/2019-ncov/hcp/guidance-prevent-spread.html>)
 - b. What to Do If You Are Sick (<https://www.cdc.gov/coronavirus/2019-ncov/if-you-are-sick/steps-when-sick.html>)
 - c. Caring for Someone at Home (<https://www.cdc.gov/coronavirus/2019-ncov/if-you-are-sick/care-for-someone.html>)
 - d. Caring for Yourself at Home (<https://www.cdc.gov/coronavirus/2019-ncov/if-you-are-sick/caring-for-yourself-at-home.html>)
5. **Targeted therapy:** There are currently no approved or proven targeted therapies for the treatment of COVID-19. While the majority of clinical trials are focused on hospitalized patients, there are a growing number of clinical trials targeting mild, outpatient cases and investigating the use of repurposed antiviral, antimalarial, and immunomodulatory agents, among others. Further information and updates can be found at <https://www.clinicaltrials.gov>. The NIH COVID-19 Treatment Guidelines (<https://covid19treatmentguidelines.nih.gov/introduction/>) provides updated recommendations and supporting evidence for antiviral therapy and immunomodulatory agents.
6. **Concomitant medications:** The NIH Guidelines provide recommendations and supporting evidence regarding the role of concomitant medications such as angiotensin-converting enzyme (ACE) inhibitors, angiotensin receptor blockers (ARBs), corticosteroids, HMG-CoA reductase inhibitors (statins), and non-steroidal anti-inflammatory drugs (NSAIDs).(76)
7. **Discontinuation of home isolation:** Clinicians should contact local military public health and/or local/state health departments regarding criteria for discontinuation of home isolation and establish clear and easy-to-follow protocols to guide staff, patients, and commands on return to work/duty criteria.(78) Local implementation of discontinuation strategies may be based on availability of testing supplies, laboratory capacity, and community access to testing. Local authorities may also consider differential application of discontinuation strategies to unique populations based on their risk for transmission to susceptible contacts (e.g., following a test-based strategy for healthcare workers or those living in congregate settings, such as nursing-home residents or basic trainees, vs a non-test-based strategy for the general population). Military bases or units may have administrative requirements for service members to be able to return to work/duty independent of clinical standards. Examples of such protocols can be found in **Appendix C**. The CDC guidelines for discontinuing isolation can be found at <https://www.cdc.gov/coronavirus/2019-ncov/hcp/disposition-in-home-patients.html>.

MANAGEMENT OF SEVERE COVID-19: TREATMENT OF CO-INFECTIONS

1. Clinical judgment and patient severity will dictate provider decision on early antibiotic therapy.
2. Procalcitonin levels have been low in COVID-19 mono-infection, with minimal bacterial co-infections reported except in pediatric patients where >80% are reported to be elevated.(79)

Clinical Management of COVID-19, v4

3. Post-mortem results reported from China suggest concern for *Aspergillus* pulmonary superinfections in critically-ill patients. This is well-described in severe influenza as well. The optimum diagnostic strategy remains to be determined, but a syndrome of worsening fever, hypoxemia, and airspace opacification in a previously-improving patient may suggest secondary aspergillosis. Diagnostic options include serum 1,3-beta-D-glucan and galactomannan assays and (potentially) galactomannan measurement in bronchoalveolar lavage (BAL) fluid, although bronchoscopy should be performed only if no less-invasive option is available and only in airborne infection isolation rooms (AIIRs) with appropriate personal protective equipment (PPE). The culture of *Aspergillus* from tracheal aspirates or BAL is suggestive but not diagnostic.
4. Recommend empiric antimicrobials for intubated patients with COVID-19. The recommended empiric antibiotic therapy is as per the 2019 ATS/IDSA Community Acquired Pneumonia (CAP) guidelines or as per critical care or infectious disease consultation.(80) As a starting point upon intubation, Table 1 can be used until consultation is available:
5. Recommend obtaining blood cultures and tracheal aspirate prior to initiation of antibiotics if feasible.
6. As noted in section on diagnostic testing, co-detection of other respiratory pathogens has been observed with SARS-CoV-2. For example, Stanford researchers recently provided rapid communication of experience with 562 SARS-CoV-2 tests; of 49 positive SARS-COV-2 results, 11 (22.4%) also had a co-infection, and of 127 positive for other viruses, 11 (8.66%) had a SARS-COV-2 co-infection.
(<https://medium.com/@nigam/higher-co-infection-rates-in-COVID-19-b24965088333>)

Table 1. Empiric Antimicrobial Considerations for Intubated COVID-19 Patients (or PUI)

	Starting Antibiotic Regimen
No comorbidities or immunosuppression or risk factors for MRSA or <i>Pseudomonas aeruginosa</i> *	<ul style="list-style-type: none">• Ceftriaxone† 2 g once daily, <u>and</u> Azithromycin† 500 mg once daily
With comorbidities‡	<ul style="list-style-type: none">• Cefepime 2 g every 8 hours, <u>and</u> Azithromycin† 500 mg once daily <i>OR</i>• Piperacillin-Tazobactam 4.5 g every 6 hours (or every 8 hours by extended infusion), <u>and</u> Azithromycin† 500 mg once daily

Definition of abbreviations: MRSA = methicillin-resistant *Staphylococcus aureus*

*Risk factors include prior respiratory isolation of MRSA or *P. aeruginosa* or recent hospitalization AND receipt of parenteral antibiotics (in the last 90 d). If concern for MRSA, add **vancomycin** 15-20 mg/kg q 8-12 hours

†If Ceftriaxone is not available, replace with **ampicillin/sulbactam** 3 g q6h; If Azithromycin is not available or contraindicated, replace with **doxycycline** 100 mg q12h


‡Comorbidities include chronic heart, lung, liver, or renal disease; diabetes mellitus; alcoholism; malignancy; immunodeficiency/asplenia.

These are general recommendations: Please refer to local antibiogram for alternative empiric choices.

MANAGEMENT OF CRITICAL COVID-19: OXYGEN & ACUTE RESPIRATORY DISTRESS SYNDROME (ARDS)

1. Give supplemental oxygen therapy immediately to patients with respiratory distress, hypoxemia, or shock and target SpO₂ 92-96%.(81, 82) Hyperoxia (PaO₂ >225mmHg) should be avoided and is associated with worse outcomes.(83)
2. Begin with low flow nasal cannula (1-6 L/min) followed by high flow nasal cannula (Figure 9).
3. **High-flow nasal cannula (HFNC).** Although an area of controversy, early expert opinion favors HFNC over other NIV modalities (https://emcrit.org/ibcc/COVID-19/#high_flow_nasal_cannula) because it appears to be well tolerated and less aerosolizing. There is presently no definitive evidence that HFNC augments transmission of virus, however HFNC will disperse air farther the higher the flow is set, (but not as far as CPAP). (84) A surgical mask should be placed over the HFNC in an effort to minimize aerosolization risk. Consider intubation for higher flow rates (>40 L/min), especially if the patient is not in a negative pressure room.
4. **Non-invasive ventilation (NIV).** It is recommended to avoid NIV because of increased aerosolization generated by the facemask and lack of an exhalation filter. If there is an exception to this such as patients that chronically use NIV or DNI patients, these patients will require airborne isolation regardless of ICU/acute care status.

General schema for respiratory support in patients with COVID-19



The optimal strategy for respiratory support in COVID-19 remains unknown. The above strategy seems reasonable, adapted largely from experience with other types of viral pneumonia. Patients with more complex respiratory disease (e.g. COPD plus COVID-19) might benefit from BiPAP. -The Internet Book of Critical Care, by @PulmCrit

Figure 9. General Schema for Respiratory Support in Patients with COVID-19 (<https://i1.wp.com/emcrit.org/wp-content/uploads/2020/03/respsup.jpg?resize=713%2C600&ssl=1>).

5. **Helmet ventilation.** The helmet can be connected to either a BiPAP circuit or a HFNC circuit (up to 60 L of flow) to increase the PEEP that the patient receives.(85, 86) The helmet has a tight seal around the neck and should decrease the amount of leak usually seen with mask interface NIV (such as BiPAP and CPAP). In one study comparing helmet to mask NIV with ARDS, there was a decreased rate of intubations.(86) There has been concern that CO₂ washout was inefficient using the helmet,(87) but a follow up study did not support that concern.(85) Helmet ventilation can prevent aerosolizing the virus.
6. **Awake proning** of non-intubated patients is currently being performed at some hospitals across the world.(88) A retrospective study of 15 non-intubated, hypoxemic patients placed in the prone position showed improvement of oxygenation. The effects were not sustained upon supine positioning.(89) See **Appendix D** for full protocol for prone positioning of non-intubated patients.
7. **Aggressive fluid resuscitation may worsen oxygenation** and outcomes in both children and adults, so in the absence of shock, fluid boluses should be minimized. Consider no more than 30 ml/kg ideal body weight (IBW) of isotonic crystalloid for adult patients, assuming no ongoing active fluid losses (e.g., from diarrhea).
8. **Avoid nebulizers**, as metered dose inhalers are recommended for staff protection/avoidance of aerosols.(61)
9. **Admission studies and labs:** Consider the following diagnostic studies in Table 2 for diagnosis, prognosis and risk stratification (and/or safety of agents) for all hospitalized patients with confirmed COVID-19 and for PUIs.
10. Due to infection prevention needs, do not allow ICU visitors during a pandemic except under exigent circumstances.
11. Facilities should assess daily operational status via huddle of equipment including ventilators, medications (e.g. analgesics, sedatives, and paralytics), and staffing (including respiratory therapists, physicians and nursing) and initiate contingency or crisis standards of care as appropriate.

Table 2. Laboratory and Study Considerations for Hospitalized Patients with COVID-19 (or PUI)

Recommended Daily Labs:

- Complete Blood Count (CBC) with differential (trend neutrophil-lymphocyte ratio, NLR)*
- Complete metabolic panel (CMP)
- C-reactive protein
- D-dimer

Recommend on Admission (may repeat q2-3 days if abnormal or with clinical deterioration)

- PT/PTT, Fibrinogen
- Ferritin
- LDH
- IL-6
- SARS-CoV-2 RT-PCR testing (e.g., CDC EUA assay, Biofire COVID-19 panel, Hologic, etc.)
- Electrocardiogram (ECG) (consider utilization of telemetry with severe infection; ECG if changes on telemetry)
- Portable CXR

If Clinically Indicated

- Blood cultures
- Troponin and BNP (if suspect acute coronary syndrome or heart failure)
- Tracheal aspirates for intubated patients
- Viral serologies if LFTs are elevated if clinically indicated (HBV sAb/cAb/sAg, HCV Ab, HIV q/2 Ab/Ag)
- For acute kidney injury (i.e. serum creatinine >0.3 above baseline), send urinalysis and spot urine protein:creatinine)
- Procalcitonin

* <https://emcrit.org/pulmcrit/nlr/>

Endotracheal Intubation

1. **Decision to intubate:** Recent non-peer reviewed references regarding COVID-19 respiratory support should be considered with caution. Neither a practice of ‘early intubation’ (reflexive decisions to intubate once a patient requires more than 5-6 L/m of oxygen), nor ‘permissive hypoxemia/happy-hypoxemic’ (allowing patients to persist with an oxygen saturation of lower than 80% for prolonged duration in order to avoid harms of intubation and mechanical ventilation) are evidence based. However, in the absence of significant clinical experience or high quality evidence with COVID-19 ventilation, it may be reasonable to not intubate select patients with stable mild hypoxemia on supplemental oxygen, but low PaCO₂ and without signs or symptoms of end-organ damage who demonstrate pulmonary shunt physiology for which intubation would not be expected to help.(90) Clinical decisions to intubate should be based on existing evidence-based guidelines balanced with evolving knowledge regarding COVID-19 and preserving healthcare staff safety.
2. **Clear indications to intubate include progressive hypoxia and worsening chest infiltrates, hypercarbia or decreasing mental status, and progressive dyspnea.**
3. Intubation (along with subsequent extubation) has the highest risk of aerosolization and exposure to COVID-19 of all procedures, and the person performing intubation is most at risk.(61) For this reason, the most experienced person should perform endotracheal intubation to reduce exposure to the healthcare team and all team members should be in appropriate PPE with PAPR during intubation. If PAPR is unavailable, an appropriate alternative may be the M50 CBRN gas mask. If these options are not available, N95, hair cover, gown, double gloves, face shields, goggles, and shoe covers should be used, along with a protective clear plastic cover over the patient to optimize protection for the providers. Consider intubation teams and limit the number of staff members during airway manipulation to reduce unnecessary exposure.
(<https://www.apsf.org/news-updates/perioperative-considerations-for-the-2019-novel-coronavirus-covid-19/>)
4. A pre-intubation checklist is encouraged, which should include supplies to be brought inside the room by specific team members and others that should remain outside the room. **Appendix E** provides an example intubation checklist (adapted from University of Washington). *Note: a disposable stethoscope should be used to avoid viral transfer and staff should touch as little as possible in the room to avoid fomites.*
5. For patients with a normal airway assessment, awake intubation should be avoided and modified RSI with sufficient muscle relaxation is strongly encouraged. For patients with difficult airways, good preparation of airway devices and detailed intubation plans should be made in advance.(91)
6. Some centers have advocated for further reducing exposure during pre-oxygenation and ventilation through preparing an additional COVID-19 Intubation Pack, in addition to intubation meds, a video laryngoscope (if used, or direct laryngoscopy), and a non-vented BiPAP mask. The following video demonstrates the set-up:
(<https://youtu.be/C78VTEAhhWU>).

7. **Appendix E** also provides a framework for intubation with medications and doses, although this is not a substitute for clinical judgement. Example cognitive aids are also located in this Appendix.
8. **Extubation:** While the risks of aerosolization of COVID-19 during intubation have been well described there has been less attention paid to extubation. During intubation, particularly with RSI, paralytics limit coughing and patient movement. During extubation coughing can be pronounced and difficult to control. A protective algorithm similar to intubation should be used for extubation. **Appendix F** provides an example protocol, which was adapted from University Medical Center in Las Vegas, NV.

Management of ARDS after Intubation

1. **Mechanical Ventilation.** It has been reported that many patients with COVID-19 pneumonia are initially characterized by a low elastance and high compliance despite severe hypoxemia, which is generally not observed in typical ARDS.(92) They presented the concept of a “Light” or “L” type with near normal lung compliance and severe hypoxemia due to a low V/Q ratio as a result of the loss of hypoxic vasoconstriction. These patients have low lung recruitability. The so called “heavy” or “H” type is more typical of ARDS with high elastance and low lung compliance and hypoxemia due to shunting through nonventilated areas of lung that are presumably recruitable.(93) It has been postulated that a high PEEP strategy in the “L” type may be ineffective and cause cardiovascular impairment while in the “H” type may lead to increased recruitment of nonventilated lung and improved oxygenation. It is likely that there is a spectrum of lung pathologies with the “L” type on one end and the “H” type on the other with most patients falling somewhere in-between or progressing towards the “H” type throughout the course of their illness. Despite these differences the best available data demonstrates that a low tidal volume approach with appropriate PEEP as described below is the most effective treatments strategy for ARDS.(82, 94)
 - a. Target an *ARDSnet lung-protective strategy (4-8 mL/kg ideal body weight)*, and lower inspiratory pressures (**plateau pressure <30 cm H₂O**).(82, 94)
 - i. Start with 6 mL/kg ideal body weight tidal volume and titrate to as high as 8 mL/kg as long as the lungs are compliant.
 - ii. In patients with moderate to severe ARDS, suggest titrating to a higher PEEP as tolerated. PEEP tables are available to guide titration: <http://www.ardsnet.org/tools.shtml>
 - b. *Permissive hypercapnia* ensuring adequate hemodynamics and a pH >7.15 may be tolerated
 - c. Humidification will likely be needed to manage thick secretions. However, keep in mind the risk of aerosolization associated with breaking the circuit to change heat and moisture exchangers (HME) if this is all that is available. Ventilators with heated humidifiers do not require breaking the circuit to humidify the inspiratory limb and are preferred. Consider clamping the ETT during any circuit breaks.
2. **Proning.** Evidence has shown that patients who are unable to adequately ventilate in the supine position may benefit from being placed in the prone position to improve oxygen saturation (PaO₂), pulmonary mechanics, and arterial blood gases (ABGs).(95-99) Anecdotal reports from Italy and Singapore have found that patients with COVID-19 usually respond well to early pronation.(88)
 - a. Prone positioning requires proper sedation/pain medications and paralytic agents if necessary.
 - b. Length of pronation cycle should be a minimum of 16 hours in the prone position with a return to supine positioning at least once a day.
 - c. Prone positioning should be performed as clinically indicated within the first 24 hours of the diagnosis of severe hypoxemia.
 - d. Recommend use of a manual proning protocol with coordination if mechanical beds are not available. **Appendix D** provides an example protocol, which was adapted from University Medical Center in Las Vegas, NV. Additional protocols (including videos) are available.(100)
 - e. Pregnancy is *not* a contraindication for proning or neuromuscular blockade.(101)
 - f. Consider the facility and staff safety and capacity for proning. Proning requires significant PPE and personnel resources (occupy staff in patient room for prolonged period of time). Circuit disconnection and loss of vascular access are among potential risks. The loss of manpower during proning and repositioning may be a contraindication in resource limited environments.

Clinical Management of COVID-19, v4

3. **Neuromuscular Blockade.** In patients with moderate-severe ARDS ($\text{PaO}_2/\text{FiO}_2 < 150$), neuromuscular blockade by continuous infusion should **not** be routinely used, but may be considered in the setting of worsening hypoxia or hypercapnia and in situations where the patient's respiratory drive cannot be managed with sedation alone resulting in ventilator dyssynchrony and lung decruitment.
4. **Airway suctioning.** Use in-line catheters for airway suctioning and clamp endotracheal tube when disconnection is required (for example, transfer to a transport ventilator). Avoid disconnecting the patient from the ventilator, which results in loss of PEEP and atelectasis.
5. **Bronchoscopy.** Routine diagnostic bronchoscopy (including nasal endoscopy or any instrumentation of this area) is **not** recommended. It is not necessary for the diagnosis of viral pneumonia and should be avoided to minimize aerosolization. Tracheal aspirate samples for diagnosis of COVID-19 are usually sufficient. If bronchoscopy is required for another reason, it should be performed with the same level of PPE as recommended for intubation.
6. **Inhaled nitric oxide and prostacyclin.** There is no evidence for routine use of inhaled nitric oxide, prostacyclin or other selective pulmonary vasodilators in acute respiratory failure. However, during emerging infectious disease outbreaks when resources are exhausted, inhaled nitric oxide and prostacyclin may be considered as a temporizing measure when patients develop refractory hypoxemia despite prone ventilation, or in the presence of contraindications to proning or ECMO.
7. **Extracorporeal Membrane Oxygenation (ECMO).** In settings with access to expertise in ECMO, consider referral of patients who have refractory hypoxemia despite lung protective ventilation who are otherwise appropriate candidates. For more information: <https://www.else.org/COVID-19>.

Oxygen Delivery and Mechanical Ventilation in Settings with Resource Limitations

1. As the COVID-19 pandemic places additional strain on available resources, the supplies of available ventilators may not meet clinical demand of patients in respiratory failure in need of invasive positive pressure ventilation (IPPV). Facilities should assess respiratory support operational status daily to account for equipment including ventilators, medications (induction agents, anxiolytics, sedatives, analgesics and paralytics), and staffing (respiratory therapists, providers and nurses).
2. Facilities must be prepared with alternate methods to support patients requiring IPPV in the event the number of patients with respiratory failure exceeds the number of ventilators. Alternate strategies in a crisis resource-limited clinical environment include the following:(102-105)
 - a. Deliver manual breaths via a manual resuscitator (“self-inflating bag”).
 - b. Transport mechanical ventilators may be used for prolonged ventilation of stable patients in the MTF (e.g. Impact 754 and 731 transport ventilators, see **Appendix G**), but need to be used with a viral filter.
 - c. Ventilators in storage (Home Station Medical Response materiel, War Reserve Material, and national stockpiles)
 - d. Anesthesia gas machines capable of providing controlled ventilation or assisted ventilation outside of the traditional use for anesthetic indication.
 - e. Some non-invasive ventilators (e.g., for CPAP or BiPAP) can be used for invasive mechanical ventilation, but should only be used if the standard ventilator supply is exhausted and it is confirmed with the manufacturer (e.g V60) that they are invasive capable and can deliver prescribed breaths. In this case, a HEPA filter should be inserted into the expiratory limb to prevent aerosolization.
3. Conserve accessories used with ventilators, but use viral filters if available. Consider extending the duration of use of breathing circuit supplies and in-line heat and moisture exchangers for treating individual patients.(102)
4. In accordance with professional society consensus statements, U.S. Public Health Service, and FDA guidance:(102, 103, 105)
 - a. Use FDA-cleared conventional/standard full-featured ventilators to support patients with respiratory failure.
 - b. Use one ventilator per patient, matching ventilator settings with the patient’s individual respiratory requirements.

- c. While ventilators may have mechanical capacity to split circuits to support multiple patients, it is excessively difficult to safely implement. There is insufficient body of evidence to support consistent application of this practice. Neither research using animals and test lungs nor case reports of crisis or contingency application of this technique establish clinical safety.

MANAGEMENT OF CRITICAL ILLNESS AND COVID-19: PREVENTION OF COMPLICATIONS

Cardiovascular Disease (CVD)

Cardiovascular comorbidities and the presence of CVD are common in patients with COVID-19 infections. The presence of CVD and risk factors correlate with increasing age, and are associated with increased mortality.(16, 106, 107)

1. **Troponins and Basic Natriuretic Peptide (BNP) Evaluation.** Elevated troponin is common (especially high sensitivity troponin), which is a strong predictor of mortality. Mild troponin elevation often does not represent a type-I (plaque rupture) myocardial infarction. The concentrations of BNP/NT-proBNP reflect the presence or extent of pre-existing cardiac disease or the acute hemodynamic stress. Troponin value, velocity of change in troponin level, elevated BNP/NT-proBNP and echocardiographic imaging should guide the management of the elevated biomarkers, although current opinion advises that troponin and BNP should only be measured if clinical evaluation suggests acute coronary syndrome or heart failure.(108)
2. **Electrocardiogram (ECG).** Recommend ECG in suspected or acute coronary syndrome. May consider of obtaining from cardiac tele-monitoring screen.(108)
3. **Echocardiogram.** An echocardiogram should only be ordered if it is likely to provide clinical benefit. Consider repeat echocardiograms only for clear change in clinical status. Point of Care Ultrasound (POCUS) exams may be used to screen/triage patients. Transesophageal echocardiogram (TEE) requests should only be considered when no other alternative imaging modalities are available as the procedure may be aerosol producing.(109)
4. **Acute Myocardial Injury.**
 - a. Definition: An algorithm for the interpretation of myocardial injury is provided for reference and is based on the 4th Universal Definition of Myocardial Infarction.(110)
 - b. Incidence and Prognosis: Recent reports found that up to 19% of hospitalized patients with COVID-19, have a combination of elevated cardiac biomarkers, in addition to electrocardiographic and echocardiographic abnormalities.(4, 5, 10, 111) There are two patterns of myocardial injury, one pattern of a continued rise with inflammatory markers, and a second pattern similar to the pattern seen in patients with predominantly cardiac symptoms.(112) Myocardial injury appears to be a late manifestation (up to 14 days from illness onset) and has been found to be independently associated with an increased risk of mortality.(5, 108, 111)
 - c. Evaluation: Cardiac Computed Tomography (CCTA): There may be a role for the use of CCTA as a non-invasive means to rule out significant coronary pathology as a cause of myocardial injury. Assessment for the appropriateness of testing and imaging protocols should be made in conjunction with a consulting Cardiologist and Radiologist as capabilities are site specific.(113)
5. **Myocarditis.**
 - a. Incidence: In a case series of 150 patients with COVID-19 patients, nearly 10% of deaths were attributed to myocarditis with circulatory failure, and in 33% of cases it was believed to have contributed as a mechanism for multisystem organ failure.(114)
 - b. Diagnosis: There is currently no role for endocardial biopsy. POCUS at initial evaluation to help protocol TTE. Serial TTE/POCUS only if it will impact management.
 - c. Management: Supportive care depending on hemodynamic status. There are case reports on different treatment strategies, but none are validated by clinical trials.(108)
6. **Acute Coronary Syndrome.**
 - a. Incidence: Based on available published data, there is a potential symptom overlap between acute coronary syndrome and COVID-19 infection.(3)

Clinical Management of COVID-19, v4

- b. Evaluation: Goal is to differentiate acute plaque rupture, demand related ischemia or myocarditis. Recommendation is for cardiology consultation when unable to determine etiology.
 - i. ST segment elevation on the 12 lead EKG has been reported in the absence of coronary thrombosis or spasms in COVID-19 patients.(106) The mechanism for these EKG changes is uncertain but is felt to be attributable to myocarditis vs possible endothelial dysfunction with micro thrombus formation.(106, 115) Confirmation of a wall motion abnormality, indicating regional myocardial ischemia, can be made with POCUS prior to invasive angiography to aid selecting a revascularization strategy. Each MTF should consider individualizing its approach to the STEMI patients based on local expertise and patient characteristics.
 - c. Management: Once the diagnosis of acute coronary syndrome is made, medical management should be coordinated with cardiology.
 - i. Cardiac Catheterization Laboratory Considerations: As most cardiac catheterization laboratories are either normal or positive pressure rooms, the benefits of invasive therapeutics must be weighed against the transmission risk to staff and patients. Deferral of invasive management can be considered based on these factors in favor of medical stabilization if necessary. Patients with borderline or deteriorating respiratory status should be considered for intubation prior to transport to the laboratory. Right heart catheterization, pericardiocentesis, and intra-aortic balloon pump placement can be done at bedside when appropriate. Fibrinolytic protocols should be reviewed at each institution with cardiology to discuss care plans if strained resources.(116)
- 7. Cardiac Dysrhythmias.**
- a. Incidence: Common CV manifestation in COVID-19 patients. Current cases series report an occurrence of unspecified arrhythmias in 17% of hospitalized patients with COVID-19 (44% of ICU patients vs 7% non ICU patients).(5) The new onset of malignant tachydysrhythmias in combination with acute myocardial injury should raise suspicion for potential underlying myocarditis.(3)
 - b. Management: Follow recently published COVID-19 specific ACLS/PALS protocols.(117) In patients with atrial fibrillation requiring cardioversion, CCTA may be preferred over TEE to rule out left atrial appendage or intra-cardiac thrombus.(113)
- 8. Heart Failure and Cardiomyopathy.**
- a. Incidence: In a recent report it was observed that 23% of patients with COVID-19 had presentations consistent with heart failure. More frequently observed in patients who did not survive the hospitalization (51.9% vs 11.7%).(5) Fulminant cardiomyopathy can occur and is thought to be a late feature described in patients recovering from respiratory failure. Cardiogenic shock and cardiac arrest contributes to 7-33% of deaths.(108, 114)
 - b. Mechanism: SARS-CoV-2 is thought to infect host cells through ACE2 to cause COVID-19, while also causing damage to the myocardium, although specific mechanisms are uncertain. (118)
 - c. Management: In the absence of high grade AV block or unstable bradycardia, cardiogenic shock, or acute kidney injury (AKI), guideline directed medical therapies should be continued in patients with heart failure as it can impact mortality.(119) Assessment of continuation of these therapies should be determined on a frequent basis depending on the patient's clinical status. Assessment of continuation of these therapies should be determined on a frequent basis depending on the patient's clinical status. The American College of Cardiology, Heart failure Society of America, American Heart association, and European Society of Cardiology have published statements at the time of this writing that recommends continuation of ACE-I/ARB therapy in patients with COVID-19.(108)
- 9. Cardiopulmonary Return to Exercise or Physical Activity Recommendations.**
- a. The timing and safety for resuming exercise, intense training, or physical conditioning in those with COVID 19 infection are unknown. Given the potential risk surrounding cardiovascular complications in COVID 19 including cardiomyopathy, arrhythmias, coronary syndromes, and thromboembolic events, the following return to duty recommendations are provided in Figure 10 with detailed information in **Appendix H.** (120-123)




Figure 10. Cardiopulmonary Return to Exercise or Physical Activity Recommendations after COVID-19. (ECG, electrocardiogram; Tnl, Troponin I; HsTn, high sensitivity troponin; TTE, transthoracic echocardiogram) a: Refer to Cardiovascular Disease Section under Management of Critical Illness and COVID 19 -Prevention of Complications. b: ECG findings that may indicate viral induced myocardial injury include pathological Q waves, ST segment depressions, (new) diffuse ST segment elevation, and T wave inversions that are outside of the normal parameters based on the Internal Recommendations For Electrocardiographic interpretation in athletes. c: Cardiac Biomarkers indicative of myocardial injury: >99th percent upper limit of normal levels for Troponin I or High Sensitivity Troponin I/ T. d: Transthoracic echocardiogram findings of cardiac injury- regional wall motion abnormalities,, abnormal systolic function with a reduced EF <45%.

Acute Kidney Injury (AKI)

1. When defined by the Kidney Disease: Improving Global Outcomes Guidelines (KDIGO) criteria,(124) AKI occurs in 61-68% of patients with COVID-19.(125, 126) Among patients with AKI in the ICU setting, a significant proportion (31-55%) require renal replacement therapy.(125, 126)
2. The etiology of AKI in COVID-19 is predominantly acute tubular necrosis in the setting of multi-organ failure and shock.(126) However, there have been unpublished reports of SARS-CoV-2 being isolated from urine and observed on kidney pathology. In conjunction with evidence that hematuria and proteinuria are common findings in COVID-19, this suggests that direct viral injury to the kidney may also play a role.(127)
3. The standard of care for critically ill patients with severe AKI is continuous RRT (CRRT). The dose of CRRT is the same as that recommended for other critically ill patients: 25mL/kg/hr.(124, 127)
4. If a MTF admits a large number of patients, it is likely that there will be a shortage of CRRT supplies. If this occurs, slow low efficiency dialysis (SLED) should be considered. SLED is a hybrid therapy that utilizes standard dialysis machines.
5. Regardless of the modality of RRT used, special attention should be paid to volume status and ultrafiltration, consistent with the goals of a restrictive fluid strategy.
6. The preferred location of a dialysis catheter is the right jugular vein, followed by a femoral vein, followed by the left jugular vein.(124) The subclavian vein should be avoided.
7. Patient with COVID-19 are hypercoagulable and will likely require anticoagulants to maintain filter patency. Regional anticoagulation with citrate is preferred, however this should only be done by centers that are already familiar with the technique given the risks of hypocalcemia and citrate toxicity. Second line anti-coagulation is heparin. This topic is reviewed extensively in section 5.3 of the Kidney Disease: Improving Global Outcomes Guidelines on AKI.(124) Other methods to improve filter patency are to increase blood flow (up to 400 mL/min), periodic 100mL flushes of the circuit, and pre-filter replacement fluid (if doing continuous veno-venous hemofiltration).

Hematology

1. Important pathophysiologic considerations concerning vasculature and blood in COVID-19:(37, 128-131)

Clinical Management of COVID-19, v4

- a. Endothelial cells abundantly express ACE2, the principal ligand for the SARS-CoV-2 Spike protein.
 - b. SARS-CoV-2 infects and damages endothelium. The endotheliopathy caused by SARS-CoV-2 is characterized by viral inclusions in endothelial cells, endothelial apoptosis and lymphocytic infiltration.
 - c. Damaged endothelium is incapable of maintaining an anticoagulant surface; microvascular and large vessel thrombosis is common in severe SARS-CoV-2 infection.
 - d. A recent study in ventilated ICU patients found thrombosis (PE or DVT) in 100% of patients receiving VTE prophylaxis (LMWH) and 56% of patients receiving full anticoagulation (anticoagulation treatment decisions made based on risk, not VTE diagnosis).
 - e. In severe SARS-CoV-2 infections, macrophage hyper-activation can occur and hemophagocytosis has been observed in spleen and lung. These findings are associated with elevated levels of IL-1B and IL-6, a so-called “cytokine storm.” Elevated inflammatory cytokine levels drive expression of other acute phase reactants including fibrinogen.
 - f. ARDS in general is associated with elevated levels of plasminogen activator inhibitors (PAI-1) and decreased fibrinolysis in lung tissue.
2. Key Hematologic Lab findings that may be associated with worsened prognosis in hospitalized patients:(5, 36, 132, 133)
- a. Lymphopenia (60% of hospitalized patients with ALC<1000; severe depletion of CD4+ lymphocytes associated with worse prognosis; lymphocyte recovery associated with viral clearance and improving clinical course)
 - b. Thrombocytopenia (most patients between 100-150; lower counts with severe disease)
 - c. Elevated D-dimers
 - d. Elevated fibrinogen (typically around 500 mg/dl)
 - e. Prolonged prothrombin time (generally mild, 1-2 seconds beyond normal range)
 - f. Hypercoagulability as measured by TEG or ROTEM (shorter K or CFT, elevated MA or MCF)
 - g. Hyperferritinemia (400-1500 ng/ml)
 - h. Elevated IL-6
3. Patient Management:
- a. Hematology laboratory testing to consider for known or suspected COVID-19 cases:(134)
 - i. CBC with differential (track lymphocyte count)
 - ii. Ferritin
 - iii. Type and Screen (needed if considering convalescent plasma treatment)
 - iv. D-dimer
 - v. TEG
 - vi. PT, aPTT
 - vii. Fibrinogen
 - viii. Anti-Xa activity
 - b. Anticoagulation considerations:(135-140) (adapted from Washington University – St. Louis)
 - i. All admitted patients should receive at a minimum VTE chemoprophylaxis (enoxaparin 40 mg sc daily). If possible, check anti-Xa daily 4hrs after third dose with goal 0.3-0.5. If at goal, no need to re-check; if not, adjust dose and monitor until at goal.
 - ii. In patients at higher risk of VTE or with more severe COVID-19 disease (evidence of coagulopathy with elevated D-dimers, prolonged PT, elevated fibrinogen, TEG hypercoagulability; intubated, prone and persistently hypoxic; MOF; requiring CVVH), it is reasonable to consider therapeutic anticoagulation or higher dose prophylaxis (e.g., enoxaparin 30 mg sc q12 hrs – “trauma dose” -- or enoxaparin 40 mg sc q12 hrs for patients with BMI>40).
 - iii. Persistent hypoxia should prompt evaluation for PE.
 - iv. VTE treatment (therapeutic anticoagulation) with enoxaparin should target anti-Xa of 0.6-1.0. Anticoagulation with unfractionated heparin should target anti-Xa of 0.3-0.7 (see dosing table).
 - v. Consider discharge prophylaxis for patients with moderate to severe COVID-19 not diagnosed with VTE (e.g., Apixaban 2.5 mg po q12 hrs for 30 days or Rivaroxaban 10mg PO q24 hours 30 days)
 - vi. In patients with VTE and/or persistent hypoxia (P/F < 150) despite maximum ventilator interventions (suggesting microvascular thrombosis), elevated fibrinogen (>500 mg/dl) and elevated d-dimer (> 6x

Clinical Management of COVID-19, v4

ULN), consider fibrinolytic therapy as a salvage regimen (tPA 50 mg bolus over 2 hours delivered with UFH full anticoagulation; re-bolus tPA 50 mg if no/transient improvement in P/F) [salvage regimen in use at BIDMC, Harvard Medical School; courtesy of Dr. Chris Barrett].

- c. Transfusion considerations: (141-144)
 - i. Convalescent plasma transfusion may accelerate viral clearance and improve clinical outcomes through passive antibody transfer, though this has not been definitively established.
 - ii. In general, plasma transfusion is considered a relatively low risk intervention. In addition to antibody transfer, plasma may have positive effects on vascular endothelium and can restore coagulation homeostasis. However; the non-immune effects of plasma transfusion in COVID-19 have not been established.
 - iii. Consider convalescent plasma transfusion in COVID-19 patients with moderate to severe disease. Obtain type and screen on admission to determine patient blood group.
 - iv. Refer to the Adjunctive Therapies Section for additional information about convalescent plasmas.
- d. **Appendix I** is a Weight-based Heparin Dosing Algorithm for venous thromboembolism

Nutrition

1. Nutrition care decisions are based on the patients' clinical presentation and the need to limit healthcare provider's exposure to patients, minimize contamination of equipment, and avoid transport.
2. Oral and enteral routes of nutrition are preferred. See **Appendix J** for Enteral Nutrition Pathway.
3. Ensure patients deficient in Vitamin D and Zinc are properly supplemented.(145-154)
4. Ensure patients get adequate amount of Vitamin A and Vitamin C either in their diet or other route of nutritional support.(155, 156)
5. Enteral Nutrition (EN) for COVID-19 Patients:
 - a. Consult a Registered Dietitian locally or via virtual health
 - b. Give early enteral nutrition (ideally within the first 24-36 hours of admission or within 12 hours of intubation), including patients on ECMO
 - c. Prefer gastric feeding for ease of placement and potential to use an existing NGT or OGT
 - d. Energy supply should target 15-20 kcal/kg actual body weight (ABW) for patients with body mass index (BMI) 18-29; target protein content is 1.2-2.0 g/kg daily. For patients with BMI 30-50, goal is 1-14 kcal/kg ABW/day and 22-25 kcal/kg ABW/day for patients with BMI >50.
 - e. Choose an nutrition formula based on facility availability and patient's medical presentation: https://www.nutritioncare.org/Guidelines_and_Clinical_Resources/EN_Formula_Guide/EN_Adult_Formulas/
 - f. Note: A standard high-protein (>20% protein) polymeric isosmotic enteral formula is recommended pending no renal insufficiency and normal GI function
 - g. Assess for risk of malnutrition/refeeding syndrome; if present, start at 25% of caloric goal (monitor serum phosphate, magnesium & potassium). Highest risk occurs during the first 72 hours of feeding.
 - h. Continuous infusion is recommended; start nutrition at a slow rate (10ml-20ml/hr) and advance to goal as tolerated (ideally within 3-7 days of initiation)
 - i. If patient is to be placed in the prone position, raise head of bed 10-25 degrees to decrease the risk of aspiration. Patients in prone position generally tolerate gastric feedings
 - j. Monitor fluid intake closely
 - k. Consider medications that provide calories and adjust tube feeding rate as needed: Propofol (1.1kcal/ml); Dextrose (3.4kcal/ml).
 - l. Labs: monitor electrolytes and glucose closely and triglycerides if patient is on propofol.
 - m. See The American Society for Parenteral and Enteral Nutrition's (ASPEN) Resources for Clinicians Caring for Patients with Coronavirus:(157) https://www.nutritioncare.org/Guidelines_and_Clinical_Resources/Resources_for_Clinicians_Caring_for_Patients_with_Coronavirus/
6. If unable to initiate EN due to failed EN trial with appropriate gastric tube placement, use of prokinetic agent, and/or post-pyloric tube placement, or EN is contraindicated (ileus, SBO, Mesenteric ischemia, high pressure respiratory pressure etc.), consult Registered Dietitian locally or via virtual health immediately for possible parenteral nutrition (PN) initiation. For patients with COVID-19, the threshold to utilize PN may be lower than other critically ill patients.

Other

1. Implement the following interventions in Table 3 below to prevent complications associated with critical illness. These interventions are limited to feasible recommendations and are based on Surviving Sepsis or other guidelines and have been adapted from the WHO guidelines for COVID-19.

Table 3. Prevention of Complications

Anticipated outcome	Interventions
Reduce days of invasive mechanical ventilation	<ul style="list-style-type: none"> • Use weaning protocols that include daily assessment for readiness to breathe spontaneously • Minimize continuous or intermittent sedation, targeting specific titration endpoints (light sedation unless contraindicated) or with daily interruption of continuous sedative infusions
Reduce incidence of ventilator-associated pneumonia	<ul style="list-style-type: none"> • Oral intubation is preferable to nasal intubation in adolescents and adults • Keep patient in semi-recumbent position (head of bed elevation 30–45°) • Use a closed suctioning system; periodically drain and discard condensate in tubing • Use a new ventilator circuit for each patient; once patient is ventilated, change circuit if it is soiled or damaged, but not routinely • Change heat moisture exchanger when it malfunctions, when soiled, or every 5–7 days
Reduce incidence of venous thromboembolism	<ul style="list-style-type: none"> • Use pharmacological prophylaxis (low molecular-weight heparin [preferred if available] or heparin 5000 units subcutaneously BID or TID) in adolescents and adults without contraindications. For those with contraindications, use mechanical prophylaxis (intermittent pneumatic compression devices)
Reduce incidence of catheter-related bloodstream infection	<ul style="list-style-type: none"> • Use a checklist with completion verified by a real-time observer as reminder of each step needed for sterile insertion and as a daily reminder to remove catheter if no longer needed
Reduce incidence of pressure ulcers	<ul style="list-style-type: none"> • Turn patient every 2 hours
Reduce incidence of stress ulcers and gastrointestinal (GI) bleeding	<ul style="list-style-type: none"> • Give early enteral nutrition (within 24–48 hours of admission) • Administer histamine-2 receptor blockers or proton-pump inhibitors in patients with risk factors for GI bleeding. Risk factors for GI bleeding include mechanical ventilation for ≥ 48 hours, coagulopathy, renal replacement therapy, liver disease, multiple comorbidities, and higher organ failure score
Reduce incidence of ICU-related weakness	<ul style="list-style-type: none"> • Actively mobilize the patient early in the course of illness when safe to do so

MANAGEMENT OF CRITICAL ILLNESS AND COVID-19: SEPTIC SHOCK & CARDIAC ARREST

Recognition of Septic Shock.

1. Recognize septic shock in adults when infection is suspected or confirmed AND vasopressors are needed to maintain mean arterial pressure (MAP) 60-65 mmHg despite adequate fluid resuscitation.(81, 158)
2. Recognize septic shock in children with any hypotension (systolic blood pressure [SBP] < 5th percentile or > 2 SD below normal for age) or two or more of the following: altered mental state; bradycardia or tachycardia (HR < 90 bpm or > 160 bpm in infants and HR < 70 bpm or > 150 bpm in children); prolonged capillary refill (> 2 sec) or feeble pulses; tachypnea; mottled or cold skin or petechial or purpuric rash; increased lactate; oliguria; hyperthermia or hypothermia.
3. Standard care includes early recognition and the following treatments within 1 hour of recognition: antimicrobial therapy, and initiation of fluid bolus and vasopressors for hypotension (Surviving Sepsis Guidelines). The use of central venous and arterial catheters should be based on resource availability and individual patient needs. Detailed guidelines from the Surviving Sepsis Campaign and WHO are available for the management of septic shock in adults and children.
4. Due to physiologic changes in pregnancy, standard risk scoring systems are less predictive for sepsis in pregnancy, although the Modified Early Obstetric Warning Score (MEOWS) has a sensitivity of 89% and a specificity of 79% in predicting morbidity in the obstetric population.(159, 160)

Septic Shock Resuscitation.

1. For septic shock in adults: give 250–500 mL crystalloid fluid as rapid bolus in first 15–30 minutes and reassess for signs of fluid overload after each bolus.(158)
2. For septic shock in children, give 10–20 mL/kg crystalloid fluid as a bolus as quickly as possible using a manual push and reassess for signs of fluid response after each bolus.(161)
3. **Avoid Excessive Fluid Resuscitation.** The cause of death from COVID-19 is most often ARDS and subsequent complications, which may be exacerbated by fluid administration. (3) Patients usually present with normal

Clinical Management of COVID-19, v4

lactate and blood pressure, but some patients do suffer from superimposed bacterial septic shock.

Conservative fluid therapy consistent with FACTT trial should be considered for patients with evidence of hypoperfusion and a without a history suggestive of hypovolemia (e.g. prolonged vomiting and diarrhea).(162) Consider use of POCUS to guide fluid resuscitation and prevent volume overload. If there is no response to fluid loading or signs of volume overload appear (e.g. jugular venous distension, crackles on lung auscultation, pulmonary edema on imaging, or hepatomegaly in children), then reduce or discontinue fluid administration. Clinical trials conducted in resource-limited studies comparing aggressive versus conservative fluid regimens suggest higher mortality in patients treated with aggressive fluid regimens.

4. Resuscitation endpoints include perfusion targets (e.g., MAP 60-65 mmHg in adults; urine output > 0.5 mL/kg/hr in adults or 1 mL/kg/hr in children; normalization of capillary refill; improved level of consciousness; and clearance of lactate).
5. In **pregnant women**, (>18 weeks gestation or when the uterus reaches the umbilicus) compression of the inferior vena cava can cause a decrease in venous return and cardiac preload and may result in hypotension and hypoperfusion. For this reason, pregnant women with sepsis and or septic shock should be placed in the left lateral decubitus position at 30 degrees to off-load the inferior vena cava. Respiratory failure and sepsis are managed similarly to non-pregnant adults.
6. Do **not** use hypotonic crystalloids, starches, or gelatins for resuscitation.
7. Vasopressors should be administered when shock persists during or after fluid resuscitation to maintain MAP goal 60-65 mmHg.
8. If central venous catheters are not available, vasopressors can be given through a peripheral IV, but use a large vein and closely monitor for signs of extravasation and local tissue necrosis. If extravasation occurs, stop infusion, aspirate as much as possible, and consider subcutaneous phentolamine. Vasopressors can also be administered through intraosseous needles.
9. If signs of poor perfusion and cardiac dysfunction persist despite achieving MAP target with fluids and vasopressors, consider an inotrope such as dobutamine.
10. Norepinephrine is considered first-line treatment in adult patients; epinephrine or vasopressin can be added to achieve the MAP target. Vasopressors are safe in pregnancy and MAP goal is >65 mmHg.
11. Intravenous hydrocortisone (200-300 mg total daily dose, administered in divided doses every 6-8 hours or as a continuous infusion after a 50-100 mg loading dose) is recommended for patients with persistent hypotension despite the use of two or more vasopressor agents.
12. Angiotensin II (Giapreza) is a vasopressor that may provide benefit in vasodilatory refractory shock as a third-line or fourth-line agent.
13. In children, epinephrine is considered the first-line vasopressor, while norepinephrine can be added if shock persists despite optimal dose of epinephrine.

Rapid Response and Code Blue.(117)

Closed chest massage and Bag Mask Ventilation are extremely aerosolizing. Each institution should consider, based on local prevalence and perceived risk, whether all Code Blues and RRTs on patients who don't have a known recent highly-reliable negative PCR should be performed with full PPE and treated as a suspected COVID-19 patient. **Appendix K** provides an example protocol, which was adapted at Brooke Army Medical Center (BAMC).

1. The American Heart Association (AHA), in collaboration with multiple medical specialty societies, released interim guidance that was published in *Circulation* on 09 Apr 2020 to help rescuers treat victims of cardiac arrest with suspected or confirmed COVID-19. Current cardiopulmonary (CPR) recommendations were reviewed in the context of the COVID-19 pandemic. In this context, the delicate balance is to provide timely and high-quality resuscitation to patients while simultaneously protecting rescuers.(117)
2. The AHA Interim Guidance provides general principles, specific strategies, and rationales for algorithm changes. It should be used to develop local "Protected Code Blue" and "Protected Rapid Response Team (RRT)" Protocols for medical emergencies that involve the resuscitation or clinical deterioration of COVID-19 suspected or confirmed patients. These policies and protocols should be peer-reviewed and based on the best available data and evidence, and should also be updated based on performance improvement data and experience. Refer to **Appendix L** for AHA ACLS and PALS protocols.

Clinical Management of COVID-19, v4

3. Protecting healthcare personnel (HCP) is a major priority in medical emergencies for suspected or confirmed COVID-19 patients. Although medical emergencies are time-sensitive situations, donning the appropriate PPE is extremely important as unintentional HCP exposure can result in detrimental effects to the workforce. Central strategies to protect HCPs during a medical emergency include efficient placement of appropriate PPE outside a patient's room, minimizing personnel in the room, and regular training.
4. Regular training should focus on the expectations, roles, and responsibilities for the individual participants in these medical emergency events, as outlined in **Appendix K**. Mock simulated scenarios should be regularly used to practice these clinical situations. Here are some hyperlinked video examples from Brooke Army Medical Center of simulated scenarios with a [COVID-19 RRT](http://amedapbamc0542.med.ds.osd.mil/VEMSWeb/VEMSHost.html?VBTemplate=Templates/UserLoginTemplate.xml) and [COVID-19 Code Blue](http://amedapbamc0542.med.ds.osd.mil/VEMSWeb/VEMSHost.html?VBTemplate=Templates/VideoInfoTemplate.xml&contentID=6475).
<http://amedapbamc0542.med.ds.osd.mil/VEMSWeb/VEMSHost.html?VBTemplate=Templates/UserLoginTemplate.xml>
<http://amedapbamc0542.med.ds.osd.mil/VEMSWeb/VEMSHost.html?VBTemplate=Templates/VideoInfoTemplate.xml&contentID=6475>
5. For a RRT or Code Blue on a suspected or confirmed COVID-19 patient, the following are important considerations and recommendations based on the principles from the AHA guidelines:
 - a. Donning of enhanced PPE in an expeditious fashion should be performed with a PPE Buddy to confirm the appropriate infection control procedures.
 - b. Consider having PPE readily available for rescuers, such as having a "go bag" or have it positioned on each ward or in the immediate vicinity of the crash cart.
 - c. Entry to a patient's room during a RRT or Code Blue should be minimized to HCP that are essential for delivery of appropriate patient care.
 - d. Close the door, when possible, to reduce the risk of airborne contamination of adjacent indoor space.
 - e. The patient should be assessed by the most senior medical staff available to determine appropriate management and disposition, unless deferred by the responsible staff.
 - f. If a patient starts to decompensate or is found unresponsive, the initial responder should prioritize the placement of a closely available surgical mask on the patient.
 - g. Chest compressions during cardiopulmonary resuscitation (CPR) is aerosol generating. Before commencing CPR, all medical personnel should wear airborne PPE, including PAPR if able. If available, an automated compressor device should be used to minimize personnel and exposure.
 - h. Appropriate equipment and supplies (viral filter, video laryngoscope, etc) should be prepositioned in the vicinity of the crash cart on COVID-19 ICUs and/or wards. Depending on local availability of resources, consider modifying the protocol for bringing the entire crash cart into the room. Due to the high risk of aerosol generation that occur during these clinical events, attempts should be made to minimize the degree and amount of door opening that occurs.
 - i. If not intubated, a non-rebreather mask should immediately be placed on the patient for passive oxygenation, covered by a surgical mask. SAFETY NOTE: ensure continuous oxygen delivery is temporarily removed for defibrillation to avoid airway fire. Depending on local protocol, a bag-valve mask (BVM) with a high efficiency particulate air (HEPA) filter may be considered if using a two-person technique to ensure a tight seal.
 - j. Minimize the likelihood of failed intubation attempts. Use an intubator with highest likelihood of first pass success. Pause chest compressions for intubation, and ideally time the pause with a pulse and rhythm check. Compared to direct laryngoscopy, video laryngoscopy is recommended as it may reduce intubator exposure to aerosolized particles; however, the intubator should use the technique with which she/he is most likely to have first-pass success.
 - k. If the patient is connected to a ventilator, minimize disconnections of the closed-circuit to reduce the potential for aerosolization. If a circuit disconnection must occur to switch to BVM with HEPA filter, recommend clamping the endotracheal tube to reduce aerosolization. If the patient is already mechanically ventilated with an advanced airway, consider maintaining a closed-circuit connection to reduce aerosolization. ***SAFETY NOTE: Use best clinical judgment and appropriate expertise for management of a patient already on mechanical ventilation at the time of cardiac arrest. Patients MUST be assessed for ventilator malfunction or airway obstruction as causative or contributing factor to cardiac arrest***. Here are the recommendations from the AHA Guidelines regarding the appropriate settings adjustments to allow for asynchronous ventilation, which replicates the bag-valve mask delivery of oxygen:

Clinical Management of COVID-19, v4

- i. Increase FiO₂ to 100% (1.0).
 - ii. Change mode to Pressure Control Ventilation mode that limits the amount to pressure to target a tidal volume of 6 ml/kg of ideal body weight.
 - iii. Adjust the trigger to “Off” to prevent the ventilator from auto-triggering with chest compressions and possibly prevent hyperventilation and air trapping.
 - iv. Adjust the set respiratory rate to 10 breaths per minute.
 - v. Assess the need and tolerance for adjusting PEEP to ensure appropriate oxygenation.
 - vi. Adjust alarms, and ensure that the endotracheal tube and ventilator circuit are secured appropriate to avoid unplanned repositioning, dislodgement, or full extubation.
 - vii. Ensure that a clamp and BVM with HEPA filter are readily available to allow an immediate switch to BVM with HEPA filter, if needed. If so, then clamp, disconnect from the ventilator circuit, connect the BVM with HEPA filter, and unclamp.
 - viii. SAFETY REMINDER: preplan and practice these adjustments with local expertise and HCP (i.e. Critical Care Physicians, Critical Care Nurses, Respiratory Therapists) to ensure appropriate understanding and avoid confusion during the actual resuscitation.
- l. Focus on potentially reversible conditions (H’s and T’s). For sudden hypoxia, use the mnemonic DOPE (Displacement of breathing tube, Obstruction, Pneumothorax, Equipment failure). Consider use of an available portable ultrasound. If a blood gas is obtained, utilize a portable analyzer or ensure appropriate infection control precautions if run outside of the room.
 - m. Avoid prolonged codes in patients with cardiac arrest. Discuss discontinuation at least after 20 minutes of a high-quality resuscitation attempt, taking into account the patient’s age, comorbidities, medical condition leading up to the event, and potential for reversal.
6. Refer to the AHA Guidelines for further discussion of specific considerations, including: pediatric, maternal, neonatal, prone positioning at time of arrest, out-of-hospital cardiac arrest, and modified AHA algorithms.
 7. The following table identifies best practices based on a “Minimum, Better, Best” model, as the COVID-19 outbreak could ultimately result in limited resources based on observational data from other countries. The goal is to achieve all elements of each category, as “Good” equates with the minimum standard-of-care while “Best” equates with the most ideal condition.

Table 4. Minimum-Better-Best Paradigm for Limited Code Blue

	Minimum	Better	Best
Advance Directives (Code status, goals of care)	Discuss & document with every patient’s medical power of attorney (MPOA) if patient unable to speak for self	Discuss & document with every patient; Involvement of Palliative Care for high risk	Develop a script for clinician that incorporates unique circumstances & ethical considerations if worsening pandemic.
Alert mechanism	Educate current Code Team members about who should respond to “Overhead Code Blue” to COVID-19 patients (used as backup or multiple codes at once)	Early activation	Directed announcement ONLY to COVID-19 Code Team
PPE / Precautions	Droplet for room; Minimize door opening	Airborne/Negative ISO; Infection Control Gatekeeper; Door remains closed	Regular practice with “Tactical PPE Donning” Checklist
Communication (via PAPRs or individuals outside room)	Whiteboard for written instructions; Closed-loop	Vocera; Speakerphone in room; Gatekeeper	Personal communication devices; Dedicated audiovisual devices
CPR	Rotate 2 individuals who don’t leave room	Rotate 2 individuals who don’t leave room and accomplish additional tasks based on pre-established priorities	Automated compressor device (e.g. LUCAS) to reduce risk to HCPs
IV access	Two standard functioning PIVs for all COVID-19 patients	Tibial IO (if needed)	Early placement of central access prior to arrest
ACLS Equipment	Dedicated Code Cart for COVID-19 ICU and wards; Accounting for Code Carts to ensure appropriate backups	For high-risk patients: consider early placement of defib pads in room or on patient, or prepositioning the Code Cart outside patient room	Specialized cart/kit containing appropriate meds, modular packs of equipment, and designated defibrillator; Dedicated COVID-19 ward: US, EKG machine, portable CXR

Clinical Management of COVID-19, v4

<p>Airway (see "Intubation" protocol)</p>	<p>Place NRB mask immediately on patient covered by a surgical mask</p>	<p>BVM with viral filter and ETCO₂ (2-person technique to ensure tight seal) Consider LMA (with viral filter) by trained and experienced personnel</p>	<p>Strict adherence to COVID-19 Intubation Protocol Protocol to identify high risk patients so that early intubation occurs BEFORE arrest</p>
<p>Simulation/Practice</p>	<p>Ongoing review and regular familiarization with this guidance; Development of didactic & "Mock COVID-19 Code Blue" scenarios</p>	<p>One-time practice with all members of the COVID-19 response teams</p>	<p>Regular practice and policy updates to all members of the COVID-19 response team</p>

Patient Transport.

1. If COVID-19 is widespread in the community, surgical masks should be considered for ALL patients irrespective of COVID-19 status.
2. The movement of patients with COVID-19 should be limited with all efforts made to ensure the patient is initially admitted to the appropriate location.
3. If patient transport is necessary:
 - a. Non-intubated patients should be transferred wearing a surgical mask over their oxygen delivery device which may include nasal prongs or a non-rebreather mask up to 15 L/min.
 - b. Staff should wear airborne PPE.
 - c. Once a patient is admitted to the ICU, transport outside of the ICU should be limited. If transport is required, then coordination should occur to ensure safety standards are maintained.
 - d. Hallways must be cleared where possible and only essential staff should accompany the patient. Staff not involved in the transfer should not come within 6 feet of the patient.
 - e. Intubated patients should have closed circuits, a HEPA filter in situ, and appropriate cuff pressure to reduce the aerosolization risks.

DoD Autopsies in Patients with COVID-19

1. To help researchers and clinicians understand COVID-19 and develop improved ways to treat severely affected COVID-19 patients with septic shock, Acute Respiratory Distress Syndrome (ARDS), myocardial dysfunction, and renal failure, Military Health System pathologists can provide critical support by performing autopsies, conducting diagnostic laboratory testing for the virus, performing critical clinical laboratory testing, and clearing blood products that facilitate safe patient care.
2. Previous communications from the Armed Forces Medical Examiner System (AFMES) and from the College of American Pathologists (CAP) discouraged post-mortem examinations due to safety concerns. Updated CDC recommendations for COVID-19 autopsies outline how to safely perform a postmortem examination, if done with trained staff and appropriate safety precautions in place, along with the appropriate consent from the decedent’s legally authorized representative.
3. When seeking consent from the decedent’s legally authorized representative, the clinical care team should include a pathologist (preferably the pathologist performing the autopsy) in the discussion. Among other things, the pathologist can explain the process, answer questions, and describe the contributions to scientific knowledge of this novel disease the autopsy would bring.
4. When conducting autopsies, pathologists need to follow CDC recommendations (<https://www.cdc.gov/coronavirus/2019-ncov/hcp/guidance-postmortem-specimens.html>). This includes the use of eye protection (goggles or face shields), techniques that minimize aerosols, environmental controls, and PPE. Consistent with CAP guidance, and with the concurrence of the Residency Program Director, pathologists should/may include pathologists-in-training.
5. The decision regarding whether to conduct an autopsy is at the discretion of the MTF Director in conjunction with guidance from the Chief of Pathology.

IMAGING OF COVID-19: RADIOLOGY DEPARTMENT GUIDANCE & IMAGING FINDINGS

Imaging findings have been widely reported in the context of COVID-19 and following initial guidance for non-urgent and elective procedures to be rescheduled, most institutions are now deciding when and how to safely

Clinical Management of COVID-19, v4

resume non-urgent, screening or elective imaging exams. Local policies for when and how to resume imaging are variable and must take into consideration many site-specific, regional and organizational factors. The American College of Radiology (ACR) has consolidated generalizable guidance for imaging department workflow, pandemic practice management, COVID-19 imaging findings and standardized reporting on its “ACR COVID-19 Clinical Resources for Radiologists” page which is updated regularly.(163)

Radiology Department Guidance Workflow.

1. A recent article by Davenport et al. addresses many considerations for radiology departments to safely resume routine care in the following categories: (164)
 - Safety measures
 - Respect local pandemic statistics
 - Risk-benefit decision making
 - Developing tiered plan for non-urgent exams
 - Accreditation and regulatory deferrals to avoid lapses
 - Address backlog of previously deferred exams
 - Manage fear
 - Develop local policies specific to academic practice environments

Use of Imaging for COVID-19

1. Whether to image a patient under investigation (PUI) for COVID-19 or previously diagnosed with COVID-19 depends on multiple factors including clinical symptoms, pre-test probability, potential for imaging results to alter management, and local resource availability. Various guidelines on imaging indications continue to be published regularly.
 - The ACR, Society for Thoracic Radiology (STR) and the American Society of Emergency Radiology (ASER) recommend that CT should not be used to screen or as a first-line test to diagnose COVID-19. (165)
 - Imaging should be reserved for cases where it will impact management or in order to evaluate for urgent/emergent alternative diagnoses. (166)
 - Multinational consensus statement from the Fleischner Society on the role of chest imaging (CXR and CT) for COVID-19 was published 7 April 2020 and provides specific imaging recommendations based on three clinical scenarios: 1) patients with mild features of COVID-19, 2) moderate-severe features of COVID-19 and 3) moderate-severe features of COVID-19 in a resource constrained environment.(167)
2. The reported sensitivity of Chest CT for COVID-19 ranges from 80-90% and the reported specificity ranges from 60-70%. (168, 169)
 - A normal chest CT does not mean a patient does not have COVID-19; a normal imaging study should not keep a patient from being quarantined if they meet other clinical criteria.
 - An abnormal CT is not specific for COVID-19 and it does not obviate the need for confirmatory laboratory testing. (170)
3. There is accumulating evidence of thromboembolic complications of COVID-19. In the event of acute clinical deterioration with suspected pulmonary embolism and/or rising D-dimer levels, CT pulmonary angiography should be considered. The National Institute for Public Health of the Netherlands recently published recommendations for imaging for pulmonary embolism or deep venous thrombosis (DVT) in COVID-19 patients.(171)
4. ACR recommends minimizing magnetic resonance imaging (MRI) use except where absolutely necessary. (ACR guidance on COVID-19 and MR use, <https://www.acr.org/Clinical-Resources/Radiology-Safety/MR-Safety/COVID-19-and-MR-Use>)
5. **Infection control and PPE:** When imaging is performed of patients who are positive or suspected positive for COVID-19, consider implementing the following infection control precautions. (166)
 - Portable imaging is preferred when possible, preferably using a portable x-ray machine dedicated for imaging COVID-19 suspected/positive patients. When possible, similar designation of other radiology equipment (e.g. ultrasound, CT and MRI) specifically for imaging COVID-19

Clinical Management of COVID-19, v4

suspected/positive patients should be made to limit cross contamination.

- Imaging should be performed nearest to the patient location to minimize exposure
 - Droplet precautions should be employed for all patients who are positive or suspected positive for COVID-19. Patients should be masked throughout the imaging exam and deep cleaning of all surfaces should be performed afterward by someone wearing proper PPE.
 - Airborne precautions are reserved for patients undergoing aerosol-generating procedures (e.g., bronchoscopy, transesophageal echocardiography, intubation, nebulization, or open suction).
 - Healthcare providers (technologist, nurse, etc.) should wear appropriate PPE (gloves, mask, eye-shield and possibly gown depending on the possibility of close or direct contact with the patient)
 - Record a census of other patients and staff present at the time of the patient visit, should the patient later test positive for COVID-19
6. When performing image-guided procedures on patients who are positive or suspected positive for COVID-19, consider implementing the following infection control precautions: (172)
- Store all PPE in secure locations with limited access, implement inventory controls, and clearly define PPE to be used based on patient status.
 - Identify a dedicated room to perform procedures on PUIs and COVID-19-positive patients. An air-negative room is strongly recommended if available.
 - Empty rooms designated for procedures on COVID-19-suspected/confirmed patients of all non-essential equipment and supplies to avoid contamination.
 - Create a staffing plan designed to preserve physician and staff availability if individuals become exposed and sick. Consider backup teams.
 - Minimize staff in the procedure room.
 - Develop clear plans for removing and disposing contaminated PPE.
 - Have a clear exit plan for COVID-19-suspected/confirmed patients to minimize staff exposure.
 - Ensure staff scrubs are changed and lead aprons are cleaned with EPA-approved disinfectants.

Thoracic Imaging Findings of COVID-19 on Chest Radiographs (CXR)

1. If imaging is part of a pre-hospital assessment of COVID-19 positive or PUI for COVID-19, portable x-ray is preferred (preferably using a dedicated portable x-ray machine to limit cross contamination).
2. In one study of 64 patients, baseline CXR had a sensitivity of 69%. (173)
3. Bilateral consolidation and ground glass opacities were the most common findings (59% and 41%, respectively) in a peripheral and lower lung distribution (51% and 63% respectively).
4. Severity of CXR findings peak at 10-12 days from date of symptom onset. (173)

Thoracic Imaging Findings of COVID-19 on Chest Computed Tomography (CT).

1. CT findings of COVID-19 overlap with findings of other viral pneumonias.
2. CT findings of COVID-19: (168, 174-177)
 - Extent - bilateral, multi-lobar
 - Distribution – peripheral and basilar or random
 - Characterization – rounded or peripheral ground glass opacities (GGO) without or with septal thickening (“crazy paving” pattern), consolidation, central low attenuation (reverse halo sign of organizing pneumonia)
3. Lymphadenopathy, pleural effusions and a nodular pattern are not common.
4. CT finding severity peak from 6-11 days after symptom onset. (178, 179)
5. Standardized reporting guidelines were developed and endorsed by the Radiological Society of North America (RSNA), the Society of Thoracic Radiology and the American College of Radiology. (180)
 - Consultation with clinical colleagues at each institution is suggested to establish a mutual approach.
 - If features of COVID-19 are discovered *incidentally* on exams performed for other indications, contact referring providers to discuss the possibility of viral infection and consider using the more general term “viral pneumonia” in the differential diagnosis. However, if after discussion COVID-19 is felt to be likely, then the authors suggest using one of the four structured reporting categories listed below.

6. Structured reporting categories for COVID-19 on chest CT.

- Typical appearance
 1. Findings: Peripheral, bilateral GGO with or without consolidation or visible septal lines (“crazy paving”); multifocal rounded GGO; reverse halo sign or other signs of organizing pneumonia (later in disease).
 2. Suggested reporting language: “Commonly reported imaging features of COVID-19 pneumonia are present. Other processes such as influenza pneumonia and organizing pneumonia, as can be seen with drug toxicity and connective tissue disease, can cause a similar imaging pattern.”
- Indeterminate appearance
 1. Findings: Absent typical features AND multifocal, diffuse, peri-hilar or unilateral GGO with or without consolidation lacking a specific distribution; lacking a rounded or peripheral characterization; few very small GGO non-rounded and non-peripheral.
 2. Suggested reporting language: “Imaging features can be seen with COVID-19 pneumonia, though are nonspecific and can occur with a variety of infectious and noninfectious processes.”
- Atypical appearance
 1. Findings: Absent typical or indeterminate features AND isolated lobar or segmental consolidation without GGO, discrete small nodules (centrilobular or “tree-in-bud”), lung cavitation, smooth interlobular septal thickening with pleural effusion.
 2. Suggested reporting language: “Imaging features atypical or uncommonly reported for COVID-19 pneumonia. Alternative diagnoses should be considered.”
- Negative for pneumonia
 1. Findings: No CT features to suggest pneumonia.
 2. Suggested reporting language: “No CT findings present to indicate pneumonia. (Note: CT may be negative in the early stages of COVID-19.)”

Cardiac Imaging Findings of COVID-19

1. In a study of 138 hospitalized patients positive for COVID-19, 16.7% of patients developed arrhythmia and 7.2% experienced an acute cardiac event.(10) Transthoracic and transesophageal echocardiography, typical first line imaging tools for the heart, require close contact with the patient and necessitate the use of high-level PPE. Please refer to the Cardiology Section in Prevention of Complications above for use of TTE/TEE.
2. To evaluate exclusion of left atrial appendage thrombus prior to cardioversion, please reference the Cardiology Section in the Prevention of Complications Section above.
3. The Society for Cardiovascular CT (SCCT) has released guidelines for the performance of coronary CT angiography based on elective indications, semi-urgent indications and urgent indications delineated on a dedicated website:(113) SCCT.org/page/COVID-19

Neuroimaging Findings of COVID-19

1. While the initial focus has been on the respiratory symptoms of coronavirus disease 2019 as drivers of morbidity and transmission, neuropsychiatric manifestations have been described as well. Perhaps most well-known is anosmia or ageusia, which occurs in the absence of nasal congestion or conductive pathology, and which has been attributed to neurotropic extension of SARS-CoV-2 along olfactory nerves, i.e. both neuroepithelial and endothelial tissue express ACE2 receptors. More nonspecific manifestations of neuro-COVID have included headache, paresthesias, and delirium (reported in 20-65% of SARS-CoV-2 patients). The underlying pathophysiology may be primary (direct viral invasion of CNS) and/or secondary (indirect effects of hypoxia or inflammatory cytokines) in nature, similar to HIV encephalopathy (181).
2. One of the earliest reports of a neuroimaging manifestation was a middle-aged airline worker with acute (hemorrhagic) necrotizing encephalopathy, which demonstrated symmetrical thalamic signal abnormalities on MRI.(182) Later case series identified diffuse confluent symmetrical white matter T2/DWI hyperintensities with microhemorrhages reminiscent of delayed posthypoxic leukoencephalopathy (183) and cortical gray matter with subcortical white matter signal abnormalities suspicious for autoimmune encephalitis in COVID-

Clinical Management of COVID-19, v4

19 patients with negative CSF RT-PCR.(184) Other autoimmune patterns have also been reported on neuroimaging of COVID-19, e.g. acute disseminated encephalomyelitis (ADEM) and Guillain-Barré syndrome (including Miller Fisher variant).

3. Although these case reports or series have called attention to the less common and more unusual neuroimaging findings of COVID-19, it should be noted that routine cerebrovascular diseases such as acute ischemic strokes, intracranial hemorrhage, and cerebral venous thrombosis are the most common neurological manifestations in hospitalized patients.(185) These cerebrovascular events reflect a combination of baseline risk factors plus endothelial injury with hypercoagulability in COVID-19 and require the usual stroke imaging evaluation (e.g. CTA) for guidance of treatment (e.g. thrombectomy for large vessel occlusion).

Abdominal Imaging Findings of COVID-19

1. In a recent single center, retrospective study of 412 inpatients with COVID-19, approximately one third had gastrointestinal symptoms. Of the patients who underwent abdominal imaging, bowel wall findings were common including bowel wall thickening, pneumatosis and portal venous gas. Possible etiologies include direct viral infection, small vessel thrombosis or nonocclusive mesenteric ischemia. Of patients who underwent ultrasound of the right upper quadrant, many had gallbladder sludge and distention, nonspecific evidence of cholestasis. (186)

ADJUNCTIVE THERAPIES FOR COVID-19: INVESTIGATIONAL APPROACHES

NIH COVID-19 Treatment Guidelines are available at: <https://www.covid19treatmentguidelines.nih.gov/whats-new/> These evidence-based recommendations focused on medical therapies are complementary to the PMG, and like it are intended to be updated regularly in response to emerging evidence.

Summary of treatment recommendations according to patient's disease status, as of 2 June 2020:

- **For uninfected persons and for asymptomatic/pre-symptomatic people with infection – No agents are recommended for use outside a trial**
- **For those with mild; moderate; severe; or critical illness – There are limited data to recommend for or against most specific antiviral or immunomodulatory therapies**

Recommendations on select specific agents:

- **For hospitalized patients with severe disease** (defined as having SpO₂≤94% on ambient air at sea level, requiring supplemental oxygen, mechanical ventilation, or ECMO), **NIH guidelines recommend the use of remdesivir (grade of recommendation: BI).**
- **There are insufficient data to recommend for or against the use of: convalescent plasma, hyperimmune immunoglobulin, or IL-6 / IL-1 inhibitors.**
- **Recommend against the use of: chloroquine/hydroxychloroquine, hydroxychloroquine + azithromycin, HIV protease inhibitors, other immunomodulators, except in a clinical trial.**

Note: *All therapies are investigational and none are proven as the literature is evolving quickly. No FDA unapproved medications should be routinely recommended for use outside of a clinical trial. There is no evidence for use of the following medications for outpatients or mildly ill patients. The American Society of Health-System Pharmacists (ASHP) website has a number of regularly updated resources at: <https://www.ashp.org/Pharmacy-Practice/Resource-Centers/Coronavirus>.*

Ethics of Clinical Research during a Pandemic: There is genuine uncertainty in the expert medical community over whether proposed off-label and investigational treatments are beneficial. Randomized, placebo-controlled trials (RCT) are the gold standard for determining if an experimental treatment can benefit patients. Some may question whether it is ethical to deprive patients of an agent that could potentially prevent or treat COVID-19, given the high mortality rate among critically ill patients and lack of known and available treatment options. A Committee of National Academies of Science, Engineering, and Medicine reviewed and conducted an analysis of the clinical trials conducted during the 2014–2015 Ebola virus disease outbreak in West Africa and found that the RCT was an ethical and appropriate design to use, even in the context of the Ebola

Guideline Only/Not a Substitute for Clinical Judgment

Clinical Management of COVID-19, v4

epidemic. The position of “ equipoise ”—genuine uncertainty in the expert medical community over whether a treatment will be beneficial—“ is the ethical basis for assigning only some participants to receive the agent. If the relative risks and benefits of an agent are unknown, participants who receive the experimental agent may receive a benefit or may be made worse off. Providing the experimental agent to all would expose all participants to potentially harmful effects.” (187)

Glucocorticoids.

1. The use of glucocorticoids is controversial and the risks must be weighed against the potential benefit. (188)
2. An early retrospective study from Wuhan, China seems to indicate possible benefit for patients with COVID-19 and severe ARDS; however, the small sample size and retrospective nature make it difficult to draw a conclusion.(189) The RECOVERY trial, although a much larger study with 2104 patients randomized to receive dexamethasone reported an improvement in 28-day mortality, has only released preliminary data as a press release, which has not yet been peer-reviewed.(190)
3. The Surviving Sepsis Campaign COVID-19 interim guidelines recommend considering methylprednisolone for intubated COVID-19 patients with ARDS.(81, 191) However, the NIH and IDSA guideline panels do not make this same recommendation, with NIH stating that there is insufficient evidence to recommend use and IDSA recommending routine methylprednisone only in the context of a clinical trial.(192)
4. In patients with severe ARDS (P_aO_2/FiO_2 ratio <100), it is reasonable to consider low dose glucocorticoid therapy (1 mg/kg methylprednisolone or 6 mg dexamethasone, especially in patients that have findings consistent with a secondary hemophagocytic lymphohistiocytosis (HLH) syndrome due to COVID-19. Common findings of HLH include: lymphopenia, splenomegaly, extremely elevated ferritin and CRP, fever. The presence of a secondary infection should also be assessed and treated if suspected.
5. Steroids may also be indicated for vasopressor-refractory shock, asthma, COPD exacerbation, or for antenatal therapy at risk for preterm birth from 24-34 weeks of gestation (see Pregnancy Section).

Remdesivir.

1. Remdesivir is an investigational intravenous drug with broad activity against RNA viruses that inhibits replication through premature termination of RNA transcription. Remdesivir has *in vitro* activity against SARS-CoV-2 and *in vitro* and *in vivo* activity against related beta-coronaviruses. It has shown promise *in vitro* and in animal models for coronavirus infections.(193-195)
2. Data from uncontrolled compassionate use case reports has suggested potential benefit; safety and efficacy data from RCTs conducted in China found that remdesivir use was not associated with the primary endpoint, a difference in time to clinical improvement. A statistically non-significant faster time to clinical improvement compared to placebo was seen among patients with symptom duration of 10 days or less; however, this study was terminated early due to low enrollment, and may have been underpowered.(196)
3. Data from the Adaptive COVID-19 Treatment Trial (ACTT) led by the National Institute of Allergy and Infectious Diseases (NIAID) (<https://doi.org/10.1056/nejmoa2007764>) indicate that patients who received remdesivir had a significantly faster time to recovery compared to placebo (11 days versus 15 days, $p<0.001$) with a trend towards improved survival.(197) In the SIMPLE trial, a Gilead-sponsored Phase 3 randomized open-label trial of remdesivir comparing 5 days to 10 days of therapy for severely ill patients not requiring mechanical ventilation, the company reported that the two regimens did not appear to differ in terms of efficacy.(198)
4. Exclusion criteria in the RCTs included increased ALT or AST levels (>5 times the upper limit of normal) and impaired creatinine clearance ($CrCl<30$), which clinicians should consider regarding applicability of the preliminary findings to the care of individual patients.
5. On 01 May 2020, the FDA issued an emergency use authorization (EUA) for the use of remdesivir. At this time, access to remdesivir under the EUA remains limited, but the MHS has received a donated supply of product that is being distributed through US Army Medical Materiel Development Activity (USAMMDA). The Health Affairs (HA) Clinical Practice Guideline Workgroup developed initial site approval criteria and USAMMDA pre-positioned medication supplies at those sites. Sites must have access to Intensive Care Unit (ICU) beds and consultation with a Critical Care or Infectious Diseases specialist (either in person or via telehealth) available. Per the HA Work Group, prescribing is restricted to Critical Care or Infectious Disease specialists. Patients must be hospitalized with IV access and have a suspected or laboratory confirmed SARS-CoV-2 infection and oxygen

Clinical Management of COVID-19, v4

saturation (SpO₂) of < 94% on room air or require supplemental oxygen or mechanical ventilation. Within the MHS, the USAMMDA Force Health Protection Division has also established an expanded access program (“Intermediate-Size Patient Population Expanded Access Protocol for Treatment of Coronavirus Disease 2019 (COVID-19) with Remdesivir”). This program has a limited number of treatment courses of remdesivir for active duty service members both within CONUS and OCONUS, as well as for federal civilian and contract employees deployed OCONUS while in support of operational forces. There may be limited availability of remdesivir for other MHS beneficiaries (e.g., retirees, dependents), which may be reviewed with USAMMDA. Access to the protocol requires a site investigational new drug (IND) submission to USAMMDA for patients meeting inclusion criteria. Clinicians can contact USAMMDA FHP Division to determine eligibility to receive product using their 24-hour international telephone: +1-301-401-2768.

Chloroquine (CQ) and Hydroxychloroquine (HCQ).

1. Previously available for use based on small reports from France that treatment with HCQ alone and HCQ plus azithromycin were associated with marked reduction in time to clearance of SARS-CoV-2 RNA, although these findings have not been substantiated in a subsequent report.(199, 200)
2. Based on these reported findings and other anecdotes, FDA issues an Emergency Use Authorization (EUA) for use of HCQ and CQ in COVID-19 patients on 28 Mar 2020.
3. Clinical data do not support the efficacy of CQ or HCQ for either treatment nor for post-exposure prophylaxis. Potential toxicities include QTc prolongation, risk for arrhythmias, retinal pigmentation, and vision loss.
4. The FDA revoked the EUA for HCQ on 15 Jun 2020. **Use of HCQ or CQ for the treatment of COVID-19 is not recommended outside of a clinical trial.** (<https://www.fda.gov/media/138945/download>)

Lopinavir/Ritonavir.

1. Coronavirus cellular infectivity and replication are dependent on virally-encoded and cellular protease activity. Clinically used protease inhibitors effective for HIV and HCV infection have been examined for potential utility in treatment of SARS, MERS, and COVID-19, but are currently not recommended.
2. On 18 March 2020, RCT results were reported that found no benefit in patients who received lopinavir/ritonavir compared to standard care for treatment of severe disease.(201-203)
3. Do not use in combination with amiodarone (fatal arrhythmia), quetiapine (severe coma), or simvastatin (rhabdomyolysis).

Host-directed anti-inflammatory strategies. ARDS and sepsis, life-threatening downstream complications of COVID-19, and many other infectious and non-infectious conditions, remain significant unmet therapeutic gaps. Historically, numerous anti-inflammatory and anti-cytokine agents, as well as many other drug candidates, have been tested and failed to meaningfully affect morbidity and mortality in ARDS, sepsis and/or septic shock.

Anti-IL6 monoclonal antibodies.

1. A variety of therapies are being administered to severely ill patients in China and elsewhere. One that is receiving substantial attention currently is an anti-IL6 receptor humanized monoclonal antibody, tocilizumab (Actemra®), which was added to the treatment guidelines published by China’s National Health Commission (4 Mar 20) to treat serious coronavirus patients with lung damage.
2. Tocilizumab and sarilumab are licensed in US for treatment of giant cell arteritis, rheumatoid arthritis, and cytokine release syndrome following CAR-T therapy. They carry a black box warning for risk of severe, potentially fatal, infections.
3. No high-quality evidence currently exists to support use. Some reports from China have suggested elevated IL6 levels are associated with severe disease in COVID-19 infection, though other reports have not found the same association. Tocilizumab has been used in Italy according to anecdotal reports and an unpublished uncontrolled case series from China treated 21 hypoxemic patients with tocilizumab 400 mg IV x1 and reported improvement in respiratory parameters.(204, 205)
4. Manufacturer-supported US randomized controlled trials of tocilizumab and sarilumab are underway.

Convalescent Plasma.

1. Convalescent plasma from patients who have recovered from SARS CoV-2 infection has been proposed as

Clinical Management of COVID-19, v4

a potential therapy for patients with severe COVID-19.(206) In an uncontrolled case series in China, 5 patients were given plasma from patients that had recovered from COVID-19 with improvement in clinical status.(207) In a randomized open-label clinical trial in China, 27/52 (51.9%) plasma recipients achieved clinical improvement in 28 days versus 22/51 (43.1%) of controls who received standard treatment. Interpretation is limited by early termination of the trial (due to decreased Wuhan COVID-19 cases) which may have been underpowered to detect a clinically significant difference.(144) In a matched control study out of Mt. Sinai, 12.8% of 39 plasma recipients versus 24.4% of 1:4 matched control patients died. In a covariates-adjusted Cox model, convalescent plasma transfusion was significantly associated with improved survival in non-intubated patients but not in intubated patients.(208)

2. As of 3 April, the FDA has authorized the use of convalescent plasma to treat “serious or life threatening” COVID-19 disease under Investigational New Drug (IND) protocols.(209)
 - a. Severe disease is defined by the FDA as follows: dyspnea, RR \geq 30 breaths/min, SpO₂ \leq 93% on RA, PaO₂:FiO₂ ratio of <300, or increases in lung infiltrations by >50% within 24-48 hours.
 - b. Life threatening disease is defined by the FDA as follows: respiratory failure, septic shock, or multiple organ dysfunction or failure.
3. A DOD Expanded Access IND protocol for convalescent plasma sponsored by Army OTSG and executed through USAMMDA FHP Division was approved on 20 May by ASD(HA) Hon. McCaffery. Patients, regardless of age or pregnancy status or beneficiary status, admitted to DoD facilities with confirmed COVID-19 and respiratory compromise (e.g., dyspnea, supplemental O₂ requirement) are eligible for treatment with convalescent plasma. CONUS and OCONUS (including deployed) providers should contact USAMMDA FHP at usarmy.detrick.medcom-usamma.mbx.force-health-protection@mail.mil or (301) 401-2768 to establish their facility as a treatment site. Plasma procurement will be coordinated by the Armed Services Blood Program.
4. DASH (HRP&O), DHA, and R2O2 coordinated with Mayo Clinic to establish a process by which DoD Military Treatment Facilities (MTF) can participate in Mayo’s Expanded Access Program (EAP) in a manner compliant with DoD requirements and simultaneously adherent to the procedures established by the Mayo Clinic for all participating sites.
5. For CONUS facilities not registered with either the DoD EAP or the Mayo EAP, clinicians can request single patient Emergency INDs (E-IND) from FDA. To obtain a single patient E-IND, the provider must determine that the probable risk to the person from the investigational drug is not greater than the probable risk from the disease or condition 21 CFR 312.310(a). For requests between 8am EST and 8pm EST (Mon-Sun), the requesting physician may contact FDA by completing Form FDA 3926 (<https://www.fda.gov/media/98616/download>) and submitting the form by email to CBER_eIND_Covid-19@FDA.HHS.gov. For E-IND requests submitted via email during this time frame, FDA will respond within 4 hours. Further guidance on E-INDs can be found in the document entitled, “How to Request and Implement a Single Patient Emergency Investigational New Drug Application,” prepared by Emily Badraslioglu, Office of Regulated Activities, USMRDC.

Several additional agents are under investigation and information is expected to emerge rapidly. Discernment of benefits and harms from novel therapies will require diligent attention to quality of evidence reported.

CARING FOR SPECIAL POPULATIONS: Pregnancy and Lactation, Infants, Children, and the Elderly

OVERVIEW

- It is not currently known if pregnant women have a greater chance of getting sick from COVID-19 than the general public, nor whether they are more likely to have serious illness as a result. Based on available information, **pregnant women seem to have the same risk as adults who are not pregnant.**
- Healthcare providers should be aware of the physiologic changes associated with pregnancy. Pregnant women have changes in their bodies that may increase their risk of some infections. Pregnant women have had a higher risk of severe illness when infected with viruses from the same family as COVID-19 and other viral respiratory infections, such as influenza.

Clinical Management of COVID-19, v4

- Healthcare providers treating pregnant women should be aware of the most current guidance on Pregnancy/Lactation guidance as prescribed by the Centers for Disease Control and Prevention (CDC), American College of Obstetricians and Gynecologists (ACOG) and Society for Maternal-Fetal Medicine (SMFM), among others.
- Visitors are limited to one (healthy) support person during the entire admission.
- Cross-collaboration with surgical services healthcare providers and communities and pediatric and neonatal healthcare providers and communities is essential to ensuring positive outcomes.

Caring for Pregnant Women during the COVID-19 Pandemic

1. Based on limited data, pregnant women do not appear to be at higher risk for severe disease. Emerging reports from the United States suggest that pregnant women may be at higher risk of being an asymptomatic carrier and having an atypical presentation with severe respiratory morbidity and preterm labor.(22-25) Clinical findings in reported cases were similar in cases of non-pregnant adults. Pregnant women experience immunologic and physiologic changes that make them more susceptible to viral respiratory infections.(23) Pregnant women are at greater risk for severe illness, morbidity, or mortality compared with the general population, as is observed with other related coronavirus infections.(24, 25) Pregnant women should receive the same care as those not pregnant in regards to screening, radiology studies, laboratory evaluations and critical care.
2. **Pregnancy complications:** Pregnancy in the setting of a COVID-19 infection is associated with higher rates of miscarriage (39.1%), preterm birth less than 37 weeks (24.3%), preeclampsia (16.2%), cesarean delivery (84%), increased incidence of neonatal admission (57.2%) and perinatal death (11.1%) Some cases of preterm birth were iatrogenic and not due to spontaneous preterm labor. (23, 24)
3. Pregnancy care should be considered non-elective during the COVID-19 pandemic.
4. Providers are encouraged to encourage patient enrollment of pregnant patients confirmed with COVID-19 in the Pregnancy Coronavirus Outcomes Registry (PRIORITY) (<https://priority.ucsf.edu/>).
5. Health care providers should be familiar with the physiologic changes of pregnancy that make pregnant women more susceptible to some respiratory infections.
 - a. Immune modulation of pregnancy
 - b. Pregnant women are more susceptible to respiratory failure and can decompensate quickly (especially in the third trimester) due to 20% decrease in functional residual capacity.
 - c. Respiratory changes: Pregnancy is a metabolically compensated respiratory alkalosis
 - i. Normal pregnancy ABG pH 7.4-7.47
 - ii. Normal pregnancy PaO₂ 75-106 mm Hg (PaO₂ increases by 30 mm Hg)
 - iii. **Normal pregnancy PaCO₂ 26-32 mm Hg** (PaCO₂ decreases by 30 mmHg)
 - iv. Normal pregnancy HCO₃ 18-21
 - d. **A PaCO₂ of 35 to 45 is ABNORMAL in pregnancy, and signifies impaired ventilation and impending respiratory compromise.**
 - e. Critical care considerations for pregnant women; online training available at <https://www.smfm.org/critical-care/cases/new-2019>
6. **Screening Guidelines:** ACOG and SMFM algorithm for outpatient assessment and management of pregnant women with suspected or confirmed COVID-19: <https://www.acog.org/-/media/project/acog/acogorg/files/pdfs/clinical-guidance/practice-advisory/covid-19-algorithm.pdf?la=en&hash=2D9E7F62C97F8231561616FFDCA3B1A6>
7. A system should be in place for pregnant women who are tested for COVID 19 to be reported to their OB Providers. This will allow OB providers to make critical delivery, care planning recommendations and decisions related to PPE recommendations, as all obstetric patients will require inpatient admission for delivery and initial postpartum period (1-4 days).
8. **Risk of vertical transmission:** Case series to date suggest no evidence of vertical transmission, similar to other viral respiratory illnesses, such as influenza.(23, 24)
9. **Changes to routine OB care** during COVID-19 pandemic: To decrease opportunities of exposure to coronavirus, OB providers should be taking steps to reduce patient encounters and optimize telehealth visits and home blood pressure monitoring. Guidance for practice has been published and we recommend developing plans at each MTF to standardize changes in Prenatal Care. (210)
10. **Inpatient Obstetric Staffing:** To ensure the availability of healthy providers and nurses to support ongoing

Clinical Management of COVID-19, v4

needs of necessary care, consider workplace segregation, which will ensure service continuity and social distancing of healthcare workers, infection control and facilitate contact tracing. This is especially important for obstetric and newborn service lines which must continue to provide necessary prenatal, intrapartum and neonatal/postpartum care.

11. Care for the pregnant patient with PUI or COVID 19

- a. **Admission:** Patients with suspected or confirmed COVID-19 should be admitted to a unit capable of caring for the respiratory needs of the patient as well as provide appropriate fetal monitoring as clinically indicated. Patient should be in isolation per hospital and CDC guidance.
 - i. Outpatient monitoring with a 14 day self-quarantine can be considered for pregnant patients with COVID 19 who have mild symptoms or are asymptomatic.
 1. Patients should be monitored closely by their health care provider for worsening symptoms. Patients should perform daily self-assessments and educations of symptoms for worsening condition.
 - a. Worsening shortness of breath
 - b. Tachypnea
 - c. Unremitting fever despite acetaminophen
 - d. Inability to tolerate oral hydration or needed medication
 - e. Oxygen saturation less 95% at rest or with exertion (if home pulse oximetry is available)
 - f. Persistent pleuritic chest pain
 - g. New onset confusion or lethargy
 - h. Cyanotic lips, face, or fingertips
 - i. Obstetrical complaints such as preterm contractions, vaginal bleeding or decreased fetal movement.
 - ii. Inpatient monitoring may be needed for the following categories of patients.
 1. Pregnant COVID 19 patients with moderate to severe signs and symptoms or oxygen saturation less than 95%
 2. Pregnant COVID 19 patients with comorbid conditions: uncontrolled HTN, inadequately controlled gestational or pre-gestational diabetes, chronic renal disease, chronic cardiopulmonary disease, concurrent pulmonary disease or immunosuppressive state (intrinsic or medication related)
 3. Pregnant COVID 19 patients with fevers greater than 39 degrees Celsius despite acetaminophen, raising concern for secondary hematophagocytic lymphohistiocytosis (sHLH)
 4. Pregnant COVID 19 patient with significant dehydration
 5. Pregnant COVID 19 patient with concurrent obstetric concerns. Society for Maternal-Fetal Medicine Management Considerations for Pregnant Patients With COVID-19 Developed with guidance from Torre Halscott, MD, MS and Jason Vaught, MD 4.30.20
https://s3.amazonaws.com/cdn.smfm.org/media/2336/SMFM_COVID_Management_of_COVID_pos_preg_patients_4-30-20_final.pdf.
 - b. COVID-19 may be associated with a transaminitis and thrombocytopenia, this is an important consideration when assessing women with a hypertensive disorder to determine if she has features of preeclampsia or HELLP syndrome (hemolysis elevated liver enzymes low platelet count).
 - c. **Guidance for treatment:** Any patient warranting pharmacologic treatment should be considered for inpatient monitoring. At this time all pharmacologic agents are considered investigational and drug efficacy in COVID 19 remains unclear. Supportive therapy should be administered. Aggressive infection control, testing for COVID-19, testing for co-infection, oxygen therapy as needed, avoidance of fluid overload, empiric antibiotics (due to risk of superimposed bacterial risk), fetal and uterine contraction monitoring for viable pregnancies, early mechanical ventilation for progressive respiratory failure, individualized delivery planning, Maternal Fetal Medicine (MFM) consultation, Pulmonology, Critical Care and Infectious disease involvement as indicated. Team based management is recommended. Consider early transfer to higher level facility if unable to provide services at MTF.(211)
 - i. Ongoing clinical trials are investigating several pharmacologic treatment strategies in non-pregnant populations. Including hydroxychloroquine, azithromycin, remdesivir, tocilizumab,

Clinical Management of COVID-19, v4

Bacillus Calmette-Guerin vaccine and convalescent plasma. None of these therapies are contraindicated in pregnancy. To date pregnancy remains an exclusion criteria for clinical trials of many therapies. Obstetric providers can advocate for compassionate use protocols and inclusion at their institutions.

- ii. Chloroquine and Hydroxychloroquine are well tolerated in pregnancy and human data in exposed pregnancies do not suggest harm.(212) In patients receiving hydroxychloroquine an electrocardiogram to assess baseline QT segment prolongation and arrhythmias should be considered. As well as a detailed medication history to assess for other medications that have QT prolongation affects.
 - iii. Remdesivir has not been studied in pregnancy and no human or animal data could be found. (212)
12. **Imaging:** Necessary radiographic studies should not be withheld from a pregnant patient. Fetal risk of anomalies, growth restriction or abortion have not been reported with radiation exposure of less than 50 mGy, a level above the range of exposure for most diagnostic procedures.
13. **Antenatal surveillance:** Gestational age appropriate fetal monitoring should be part of the initial assessment of any women with respiratory symptoms. Continuous fetal monitoring in the setting of severe illness should be considered only when delivery would not compromise maternal health, or as another noninvasive measure of maternal status. For women who recover from an acute infection, antepartum testing later in the pregnancy is not needed.
14. **Ultrasound:** Consider a detailed level 2 anatomic survey for women following recovery from a first trimester infection and a fetal growth assessment in the third trimester for women who recover from an infection later in pregnancy (later second trimester and third trimester infections). Healthcare providers should be aware of the ISUOG Safety Statement: Safe performance of obstetric and gynecological scans and equipment cleaning in the context of COVID-19 (<https://www.isuog.org/resource/isuog-safety-committee-position-statement-safe-performance-of-obstetric-and-gynecological-scans-and-equipment-cleaning-in-the-context-of-covid-19.html>). **Appendix M** has detailed instructions on preparation and cleaning of ultrasound rooms for patients.
15. **Delivery planning for the COVID-19 Patient:** Timing of delivery, in most cases, should not be dictated by maternal COVID-19 infection. For women infected early in pregnancy who recover, no alteration to the usual timing of delivery is necessary. For women infected in the third trimester who recover, it is reasonable to attempt to postpone delivery (if no other medical indications arise) either until a negative COVID-19 testing result is obtained or quarantine status is lifted in an attempt to avoid transmission to the neonate. ***In general, COVID-19 infection itself is not an indication for delivery.*** Recommend health care team wear appropriate PPE during delivery and delivery should occur in a negative pressure room. Skin to skin care following delivery is not recommended. In cases of severe maternal infection with a term infant, care teams may consider avoiding delayed cord clamping to minimize the risk of transmission to the neonate.
16. **Protocols for inpatient care of the COVID-19 Pregnant Patient**
- a. Vital sign assessment: depends on the severity of the illness. For patients with mild symptoms requiring inpatient management, vitals signs every 4-8 hours and as needed. For patient with severe disease vital signs every 2-4 hours is appropriate. For patients with critical illness continuous pulse oximetry and telemetry should be utilized. Noninvasive and invasive cardiovascular monitoring as indicated and vital signs and respiratory support as needed and at least every 1-2 hours.
 - b. **Fetal monitoring:** at > 24 weeks, electronic fetal monitoring for antenatal surveillance at least daily. Recommend additional fetal monitoring with any change in the maternal status if a cesarean at bedside is feasible. The fetus can be a sixth vital sign reflecting early deterioration in maternal status.
 - c. Recommend maintaining maternal O₂ saturations at > 95%.
 - d. Early warning signs of worsening condition: increased sensation of dyspnea or work of breathing. Inability to maintain adequate oxygen saturation. Persistent or more frequent fevers. Worsening myalgias.
 - e. ICU admission criteria: Presence of any of the following
 - i. Inability to maintain oxygen saturation > 95% with supplemental oxygen or rapidly escalating supplemental oxygen requirement.
 - ii. Hypotension (MAP > 65) despite appropriate fluid resuscitation (500-1000 mL bolus of crystalloid fluids).

Clinical Management of COVID-19, v4

- iii. Evidence of new end organ dysfunction (altered mental status, renal insufficiency, hepatic insufficiency, cardiac dysfunction, etc).
 - f. Pregnancy has a natural respiratory alkalosis with a normal PCO₂ of 28-32.
 - g. Therapy for ARDS involves low tidal volumes and permissive hypercapnia (PCO₂ > 60). Data on permissive hypercapnia in pregnancy are limited, but there do not appear to be adverse fetal effects.
 - h. It may be necessary to increase tidal volume and/or PEEP to meet goal PaCO₂ and oxygenation targets while remaining mindful not to allow alveolar plateau pressures to exceed 35 cm H₂O.
 - i. Prone ventilation has been found to improve oxygenation in the setting of ARDS. If the patient would benefit from prone ventilation it should be performed and is safe.
 - j. In the third trimester, increased PEEP may be required for pregnant moms on mechanical ventilation.
 - k. Venovenous ECMO is a proven life-saving salvage therapy for severe reversible respiratory failure, and its benefit among critically ill pregnant women has been reported.
 - l. Goal BP should be < 160/110.
 - m. Patient should be positioned with left lateral tilt (if no other position is mandated for their treatment, for example, prone position) to relieve pressure from the gravid uterus on venous return.
 - n. Therapeutic anticoagulation in critically ill pregnant patients: antepartum and postpartum
 - i. Prophylactic heparin or low-molecular weight heparin if there are no contraindications to use should be considered.(213)
 - ii. There is limited data on the use of therapeutic anticoagulation for severe COVID 19 disease.
 - iii. For therapeutic anticoagulation without confirmed thrombosis in a critically ill pregnant patient, unfractionated heparin should be considered due to its short half-life and reversibility with protamine sulfate. Unfractionated heparin should be considered for prophylaxis in patients at high risk for preterm birth due to its potential reversibility.
 - o. **Antibiotics:** If co-infection is suspected cultures should be obtained when possible and appropriate antibiotics should be started within 45 minutes of diagnosis. Ceftriaxone plus azithromycin or ceftriaxone alone are commonly used and are not contraindicated in pregnancy.
 - i. For patients with severe disease or who have risk factors for hospital acquired, ventilator acquired or drug resistant types of pneumonia, broad spectrum agents should be employed such as cefepime, meropenem, piperacillin-tazobactam, linezolid, and vancomycin all are acceptable for use in pregnancy.
 - p. Judicious use of corticosteroids for fetal maturation, balancing gestational age with potential risks of worsening maternal morbidity. Discussions with critical care team and neonatology.
- 17. Delivery Planning in ICU**
- a. Equipment for emergency cesarean delivery should be at bedside, with neonatal resuscitative equipment including warmer.
 - b. Hemorrhage Code Purple cart stocked with medications and devices should be in the ICU. Medications should readily available include methergine, hemabate, Tranexamic acid (**TXA**) and misoprostol.
 - c. Use of terbutaline should be reviewed with critical care team, depending on patient's clinical status due to the risk of tachycardia.
 - d. Establish effective means of communication with Nursing, ICU, anesthesia, neonatal, and obstetrical teams.
 - e. If emergent delivery is planned, this may be performed at bedside in the ICU, or in a main operating room.
 - f. Timing – consideration should be given to delivery > 34 weeks for critically ill maternal patient. Delivery consideration should be weighed carefully the risks and benefits. Decision for delivery requires close communication between the maternal fetal medicine and critical care team.
 - i. In the third trimester, the pressure of the uterus can decrease expiratory reserve volume, inspiratory reserve volume, and functional residual capacity, which can increase the risk of severe hypoxemia in pregnant patients, especially those who are critically ill.(214)
18. Intrapartum care if a pregnant patient at term in critical condition goes into labor, precautions as above should be initiated. Assisted second stage (OB forceps/Vacuum) is likely to be necessary.
19. A dedicated obstetrician should be present at the time of delivery, and infant placed in isolation after delivery given the unknown risks of transmission.
20. Prevention of postpartum hemorrhage as detailed above.

Clinical Management of COVID-19, v4

21. Breast pumping encouraged after review of maternal medications.

22. **Obstetric medications**

- a. Indomethacin – in the setting of indications for tocolysis, nifedipine may be considered as an alternative, given the uncertainty regarding NSAID impact on COVID-19.
- b. Betamethasone/Dexamethasone for fetal maturation – given the association between steroids and worsening morbidity of viral pneumonia, specifically COVID-19, steroids for fetal maturation should be used judiciously. **AVOID late preterm steroids 34-46 weeks** in COVID-19+/PUI patients.
- c. Magnesium sulfate is recommended for fetal neuroprotection for anticipated preterm delivery <32 weeks or for seizure prophylaxis for Preeclampsia with severe features. Given potential respiratory complications, **use judiciously in the setting of severe respiratory symptoms**. Magnesium sulfate may be used in patients with mild-moderate symptoms, may consider single 4 gm bolus.

Table 5. Use of Common Obstetric Medications:

Gestational Age	< 32 weeks		32-34 weeks		34-36 weeks
Respiratory Sx	Mild-Mod	Severe	Mild-Mod	Severe	Any
Steroids for fetal maturation /rescue steroids	Use	Discuss risks/benefits with multi-D team (ID, Critical care, Neonatology)	Consider	Avoid	Avoid
Indomethacin	May consider	Use nifedipine instead	Use nifedipine instead	Use nifedipine instead	Consider nifedipine instead
Magnesium sulfate neuroprotection	Use	Discuss risks/benefits with multi-D team (ID, Critical care, Neonatology)			

23. **Cardiac arrest:** in pregnancy should be managed similar to cardiac arrest in non-pregnant adults. If pregnancy is ≥ 20 weeks (uterus at or above the umbilicus), significant aortocaval compression exists. Left uterine displacement is recommended during high-quality CPR, with resuscitative cesarean delivery (perimortem cesarean delivery) if ROSC not achieved by 4-5 minutes. Resuscitative cesarean delivery should be performed at the bedside (do not move to the OR).(215)

24. **Intrapartum Care during the COVID-19 Pandemic:**

- a. Screen all patients and support person(s) according to ACOG SMFM algorithm upon presentation to L&D.
- b. We also suggest asking all patients and support person(s) about exposures (close contact) to COVID positive patients and if they themselves have been tested in the past 14 days for COVID-19.
- c. Recommend a designated staff member at the front of the unit to verbally screen for URI symptoms, diagnosis of COVID-19 or PUI within the past 2 weeks.
- d. Any patient with fever, cough, or respiratory symptoms (+/- fever) should put on a surgical mask and be evaluated by a nurse or provider (and put in a room).
- e. If a patient screens positive to any of the above prior to a scheduled delivery (IOL or CD), evaluate to determine if re-scheduling in 2-3 days is feasible to allow for results of COVID-19 testing
- f. For COVID-19 positive patients with mild or moderate symptoms not requiring immediate care, it is important to recognize that the severity of disease peaks in the second week, so planning delivery prior to that time is optimal.
- g. Risk of vertical transmission – no documented evidence of vertical transmission based on limited case series
- h. Avoid oxygen for fetal resuscitation (this intervention has not been shown to be beneficial and may increase the risk of aerosolization).
- i. If a birth partner (support person) has a fever, cough, or respiratory symptoms (+/- fever) (or confirmed COVID-19 positive or PUI), they should not come to L&D, and will not be admitted to L&D as a support person.
- j. Support persons of a COVID-19 positive or PUI mother should wear a mask during their hospital stay, and are restricted to the patient room. They are not allowed to visit hospital areas outside the patient room. They should use the bathroom in the patient room, and should have all meals brought to the room.

Clinical Management of COVID-19, v4

- k. Routine preoperative labs for scheduled cases should be drawn the day of procedure to minimize trips to the hospital.
 - l. **Intrapartum fever** – should be evaluated in the usual fashion with consideration for both obstetric and non-obstetric causes. Recommend empiric treatment for the clinically suspected cause (e.g. chorioamnionitis), with increased vigilance and consideration of rapid COVID-19 testing, as early experience has shown the possibility of asymptomatic pregnant patients to develop symptoms postpartum.
 - m. **Cesarean section:** As for all patients, cesarean section should be reserved for maternal and fetal indications. Consider conversions of operating rooms to negative pressure rooms (conversion to negative pressure ante-rooms or neutral pressure ORs are alternatives) for COVID positive or PUI. Such conversions may not be possible in all facilities, and with proper PPE and patient transfer protocols, cesarean deliveries can still be safely performed in a positive-flow OR. In general, negative pressure ORs should not have open surgical equipment (as is often done for designated emergent cesarean delivery rooms). Teams should coordinate with local infection control teams to inform these decisions. Consider universal airborne PPE use (including N95 masks) for all surgical procedures for COVID+/PUI patients during labor and delivery due to high risk for aerosolizing procedures (intubation).
25. **Support Person:** If a birth partner (support person) has a fever, cough, or respiratory symptoms (+/- fever) (or confirmed COVID-19 positive or PUI), they should not come to L&D, and will not be admitted to L&D as a support person.
- a. Visitors are limited to one (healthy) support person during the entire admission. (210)
 - b. Support persons of a COVID-19 positive or PUI mother should wear a mask during their hospital stay, and are restricted to the patient room (should not visit hospital areas outside patient room). They should use the bathroom in the patient room, and should have all meals brought to the room.
26. **Inductions of Labor:**
- a. Induction of labor with medical indications in asymptomatic women should **NOT** be postponed or rescheduled. This includes 39-week inductions after patient counseling. However, in cases of extreme healthcare burden, it may be appropriate to consider postponing or rescheduling inductions. For example, in a region early in a COVID-19 emergency, it may be prudent to get patients delivered prior to high COVID-19 burden in the hospital.
 - b. Consider outpatient cervical ripening with Foley in low-risk women to limit hospital time.
 - c. Management of the first stage of labor is not generally altered. Oral restriction of fluid and solid food in the first stage of labor is not recommended, oral water and clear fluids can be encouraged as tolerated in labor. If oral restriction, IVF at 250 mL/hr. containing dextrose, with upright positions in the first stage of labor for women without epidural. If walking, must stay in the room. Oxytocin augmentation is recommended to shorten time in labor if slowed progress, with early amniotomy.
 - d. Intrapartum oxygen therapy has no fetal benefit and may cause harm, **recommend NOT utilizing oxygen therapy for fetal resuscitation.** Given the high rate of asymptomatic carriers, this principle applies to all patients on L&D regardless of the patient's COVID-19 status. Supplemental oxygen may be administered for maternal indications, cover nasal cannula with a surgical mask.
27. **Second Stage (Pushing to delivery):** Pushing should not be delayed for any delivery as it prolongs time to delivery and increases chorioamnionitis and postpartum hemorrhage.
28. **Third Stage (Delivery of baby to delivery of placenta):** There are concerns about limited blood resources during the COVID-19 pandemic. The below recommendations apply to all deliveries to further minimize use of blood products at delivery.
- a. Recommend optimizing antenatal hemoglobin prior to delivery to minimize the need for blood transfusion at delivery.
 - b. Consider 400 mcg misoprostol buccally with delivery (to decrease risk of PPH).
29. **PPE Considerations during COVID-19 Pandemic for Pregnancy:**
- a. Screen positive patients (symptoms or prior COVID-19 diagnosis) or PUI:
 - i. PPE during admission: Surgical mask for all patients with symptoms or COVID-19+/PUI. Airborne precautions: N95 masks and droplet PPE (Gown, gloves, mask/face shield) for all HCP.
 - b. Screen negative patients (no symptoms or prior COVID-19 diagnosis):

Clinical Management of COVID-19, v4

- i. PPE during delivery: Surgical mask and droplet PPE (Gown, gloves, mask/face shield) should be used during all patients in the second stage. N95 Mask could be considered for the surgical team for any cesarean section as there is the potential risk of requiring intubation during the surgery. Provider discretion and individual MTF PPE availability can be considered.(210)

Table 6. Suggested PPE During Obstetric Care (210)

Care situation	Surgical mask	Droplet PPE	N-95 mask or PAPR
		(gown, gloves, surgical mask/ face shield)	
Patient (cloth mask acceptable if no resp sx)	X		
Provider during routine encounters	X		
Provider during patient encounters with URI symptoms		X	
Provider during patient encounters with suspected or confirmed COVID-19		X	X

- c. Women who are COVID-19+ or PUI should wear a surgical mask at all times as clinically able.
 - d. Women who are COVID-19+ or PUI should be placed in an isolation/private room. Airborne infection isolation rooms (negative pressure rooms), if available, can be used if performance of aerosolizing procedures is anticipated. In general, isolation rooms with droplet precautions are recommended.
 - e. Staff PPE:
 - i. Proper donning and doffing of PPE takes time. Training in the use of PPE should emphasize safety of healthcare workers, recognizing that clinical response times may be slowed by these precautions.
 - ii. Proper donning and doffing procedures should be reviewed and practiced frequently; Recommend simulated patient transfers (e.g. from L&D to OR).
 - iii. Recommend posting diagrams and checklists in areas where donning and doffing will occur.
 - iv. For HCP that do not fit N95 masks, PAPR should be used. For staff in the operating the OR, the PAPR with shroud must be used, followed by sterile gown over the shroud. This ensures proper venting of the PAPR out the bottom of the surgical mask to ensure sterility of the field.
 - v. Have an observer witness donning/doffing when possible.
 - f. Anticipate emergencies as best as possible; plan ahead and proactively intervene for situations that could result in emergent cesarean delivery (e.g. Category II FHR), early peds notification.
 - g. Collaborate closely with Surgical Services to support additional operating room/staffing capabilities.
 - h. Define patient OR plan on admission (COVID-19 or not)
 - i. Coordination with Pediatrics and Neonatology upon admission for any mother COVID-19+ or PUI
- 30. Considerations for Support Person/VISITORS to L&D and ANTEPARTUM/POSTPARTUM**
- a. One designated (healthy) support person during the entire admission, easily identifiable by L&D staff. Consider a colored wrist band for identification. Support person should be screened as above, wear a mask, and remain restricted to the patient room for mothers that are COVID-19 positive or PUI.
 - b. No children < 16 years permitted.
 - c. Additional visitors for end-of life situations or bereavement (e.g. IUFD) may be considered/evaluated on a case-by-case basis.
 - d. All efforts should be made to limit the movement of COVID-19 positive/PUI women from one care area to another. Consider postpartum care in the same room as delivery if possible.
 - e. If increased prevalence of disease and community transmission is present, individual MTFs could consider a no visitation policy to minimize potential exposure of staff and patients.
- 31. Anesthesia Considerations for Intrapartum Care (Refer to Implications for Surgical Care Section)**
- a. Recommend early epidural to minimize need for general anesthesia in the event of an emergent cesarean.
 - b. COVID-19 is not a contraindication to neuraxial anesthesia.
 - c. Anticipate emergencies as best as possible; plan ahead and proactively intervene for situations that could result in emergent cesarean delivery (e.g. Category II FHR tracing).
 - d. Recommend limiting exposure of trainees to COVID+/PUI, with experienced staff providing care.
 - e. Suspend nitrous oxide programs on L&D due to possible aerosolization.

32. Postpartum Care

Clinical Management of COVID-19, v4

- a. Women should be notified that in order to limit the risk of infection to themselves, staff and other patients, mothers and infants should be discharged in an expedited and safe fashion. Vaginal deliveries – goal of discharge on postpartum day 1 (same day for select women). Cesarean deliveries – goal of discharge on postpartum day 2. Home blood pressure monitoring devices may be needed.
 - b. All postpartum visits, including wound checks, should be via telehealth. Can optimize by uploading photos through EMR/patient portals.
33. **Pregnant patient work restrictions:** Delivery is a unique scenario in the COVID-19 pandemic. Hospital admissions for delivery are anticipated around the patient’s due-date. In anticipation of hospital admission for delivery, if feasible and mission permitting, consider having pregnant women work from home at 37 weeks (2 weeks prior to 39 weeks or 2 weeks prior to anticipated delivery), and practice strict social isolation during this time. (210) Strict social isolation is encouraged for the entire family unit. The goal is to limit risk of exposure around the time of delivery. Depending on mission requirements and increasing disease burdens, such accommodations may not be possible but should be considered.(210)
34. **Pregnant health care workers:** Facilities consider limiting exposure of pregnant HCP to patients with confirmed or suspected COVID-19 infection, especially during higher-risk procedures such as aerosol generating procedures (intubation, extubation, BiPAP, high flow nasal cannula, nebulized medications) if feasible based on staffing availability. With ongoing stresses in the MHS and increasing disease burdens, such accommodations may not be possible.

Caring for Infants and Mothers with COVID-19: IPC and Breastfeeding

1. Current evidence is inconclusive about in utero transmission of SARS-CoV-2 from mothers with COVID-19 to their newborns. Transmission of SARS-CoV-2 can occur after birth via contact with infectious respiratory secretions. Data suggests that infants may be at higher risk for severe illness compared with older children. (216, 217)
2. To reduce the risk of post-natal transmission from mother to infant, the CDC and American Academy of Pediatrics recommends consideration of temporarily separating a symptomatic PUI or COVID-19 positive mothers from her infant (e.g. separate rooms). In the absence of more definitive data, this decision should reflect an individualized risk - benefit consideration for the mother and infant, cognizant of the potential for delayed maternal-child bonding and impaired breastfeeding. This will require an additional healthy (non-infected) adult to care for the infant while separated from mother.
3. COVID-19 positive postpartum mothers as well as postpartum PUIs will be counseled about the risks and benefits of colocation vs. separation.
 - a. If a postpartum PUI mother elects to be separated from infant and then her test is negative for COVID-19, the mother and infant can be reunited and ‘room in.’
 - b. Postpartum COVID-19+ or PUI mothers who elect to co-locate (also referred to as ‘rooming in’) with their infants should be instructed to wear a facemask at all times. They will also practice hand hygiene before each feeding and wear gloves during infant contact. They will also be encouraged to wash any skin that may come in contact with the infant (e.g. breasts, chest, arms, etc.). They will be encouraged to limit other close contact with the infant(s) and a separate non-infected caregiver should be present to help care for the infant. This separate non-infected caregiver should perform a majority of the infant’s care. While not breastfeeding, infants should be kept > 6 feet away from the mother within the room, per CDC guidance.

Lactation: Breastfeeding, Pumping, or Expressed Breast Milk (218)

1. Breast milk is the best source of nutrition for most infants. It is unknown if mothers with COVID 19 can transmit the virus via breast milk.
2. Postpartum patients with COVID-19 who are pumping should be provided with a dedicated breast pump while inpatient and follow CDC guidelines on equipment use and feeding. Mothers should wash hands and breasts before pumping and wear a mask.
3. Recommended procedure to follow while pumping milk:
 - a. Wipe the surface where syringes/bottles will be placed after collection with a germicidal disposable wipe, and cover surface with clean paper towel or cloth.
 - b. Mother collects breast milk by hand or by pump into clean syringes or bottles then ensures

Clinical Management of COVID-19, v4

syringe/bottle cap is secured. The outside of the container will be wiped with a germicidal disposable wipe. A label is then placed to identify date, time, and patient.

- c. Transport and storage of breast milk from mother's room to common refrigerated storage area should follow strict infection control procedures per hospital policy.

Infants

1. Infants born to mothers with confirmed COVID-19 should be considered PUIs.
2. All infants born to mothers with suspected or confirmed COVID-19 should, if resources allow, be resuscitated in a separate adjacent room utilizing airborne PPE due to potential for aerosolization during the 2nd stage of labor and potential need for intubation of the infant. The infant should be bathed as soon as clinical condition allows.
3. The CDC recommends testing for all neonates born to women with confirmed or suspected COVID-19. Infants should be tested at ~24 hours. If initial test results are negative, testing should be repeated at 48 hours. If infant is asymptomatic and expected to be discharged at <48 hours, a single test performed when the infant is >24 hours is acceptable.(217)
4. All elective procedures to include circumcision should be deferred while infant is a PUI.
5. If hearing tests can be performed outpatient, it is acceptable to defer until COVID-19 testing is negative. If it is not easily available outpatient, ensure proper disinfection measures are used when cleaning equipment.

Neonatal Intensive Care Unit (219)

1. Recommend any infant who has symptoms that meet criteria for NICU admission be assessed by the NICU team and admitted to a COVID-19 cohort pod or other segregated section of the unit.
2. Healthcare workers should wear full PPE including N95 (or PAPR), eye shields, gown, hair cover, and gloves should be worn when caring handling the PUI infant.
 - a. In situations where there is limited PPE available, N95 (or PAPR) use should be prioritized for use in infants requiring CPAP, SiPAP, use of high-flow nasal cannula with flow rate >2LPM, or undergoing aerosolizing procedures such as intubation.

Newborn Visitation

1. No visitors experiencing cough, fever, or shortness of breath should be allowed in any care setting.
2. All visitors should wear a facemask and adhere to local infection control policies.
3. For NICU: Visitation should be limited to the mother and one support person. COVID-19 positive persons or their household contacts should not be allowed to visit until they meet the following requirements:
 - a. Resolution of fever without antipyretics for 72 hours with improving respiratory symptoms; **AND**
 - b. At least 10 days have passed since symptoms first appeared or since the positive test; **OR**
 - c. Negative results of a molecular assay for detection of SARS-CoV-2 from at least 2 consecutive nasopharyngeal swab specimens collected at least 24 hours apart.
 - d. Entrance to other family support personnel should be determined on a case by case basis.
4. For Labor and Delivery, Post-partum / Newborn Nursery: each COVID-19 positive or PUI postpartum mother may be allowed to have one support person with her who must remain with her throughout the admission. This support person should be isolated to the post-partum room and not traveling elsewhere in the hospital.
 - a. If the mother chooses to co-locate with the infant, the support person should help with infant care.
 - b. If the mother chooses to be separated from her infant, the support person may help with the infant's care when they are brought to the room.
5. AAP recommends that well newborns, defined as negative molecular testing and asymptomatic, can receive circumcision. Newborns who are PUIs are not eligible for elective circumcision.

Newborn Discharge

1. After hospital discharge, a mother with COVID-19 is advised to maintain a distance of at least 6 feet from the newborn, and when in closer proximity, to use a mask and hand-hygiene for newborn care until:
 - a. She is afebrile for 72 hours without use of antipyretics; **AND**
 - b. At least 10 days have passed since symptoms first appeared; **OR**

Clinical Management of COVID-19, v4

- c. She has negative results of a SARS-CoV-2 test from at least two consecutive specimens collected ≥ 24 hours apart.
2. A mother with COVID-19 whose newborn requires ongoing hospital care should maintain separation until:
 - a. She is afebrile for 72 hours without use of antipyretics, with improving respiratory symptoms; **AND**
 - b. At least 10 days have passed since her symptoms first appeared or since the positive test; **OR**
 - c. Negative results are obtained from at least two consecutive SARS-CoV-2 nasopharyngeal swab test collected ≥ 24 hours apart.

CDC Guidance: <https://www.cdc.gov/coronavirus/2019-ncov/hcp/caring-for-newborns.html>

AAP Guidance: <https://services.aap.org/en/pages/2019-novel-coronavirus-covid-19-infections/fags-management-of-infants-born-to-covid-19-mothers/>

Caring for Children with COVID-19

1. Children (0-18 years) currently make up less than 2% of the laboratory-confirmed COVID-19 cases worldwide, though this may represent an ascertainment bias of symptomatic cases or known contacts.
2. Approximately 90% of children with COVID-19 remain asymptomatic or have mildly symptomatic disease.
3. Hospitalization rates in the US are higher for children 0-4 years-old as compared to 5-17 years-old but both are still markedly below adults. This is consistent with data from China that showed children < 5 years-old having more severe disease with the highest rates in those < 1 year-old. There are case reports in the US of febrile infants < 3 months-old who required hospitalization for fever evaluation, all of which was negative except positive for SARS-CoV-2.(31, 220-222)
4. One report found 23% of children with symptomatic COVID-19 had ≥ 1 underlying condition with the most common including chronic lung disease (including asthma) and cardiovascular disease.(223) Obesity is an emerging comorbidity amongst those hospitalized.(224)
5. The mortality rate appears to be extremely low worldwide; total pediatric deaths related to COVID in the single digits per each country.(225, 226)
6. Respiratory virus co-infections and secondary bacterial infections are possible. One study found 20.7% of COVID+ patients were also infected with one or more respiratory pathogens and results did not differ significantly in age (child vs. adult). The most common co-infections were rhinovirus/enterovirus (6.8%), RSV (5.2%) and non-SARS-CoV-2 coronaviridae (4.3%).(227)
7. Pediatric symptoms, if present, are similar to common viral upper respiratory infections, which differs from adults, who tend to have lower respiratory symptoms as most prominent. Fever is the most common symptom but often brief, lasting only 1-2 days. GI symptoms such as diarrhea have been reported but are less frequent than in adults. Severe symptoms include hypoxia and respiratory distress.(28, 31, 223) The common age-specific presentations are as following:
 - a. Neonates: temperature instability, lethargy, poor feeding, shortness of breath
 - b. Infants: fever, rhinorrhea, congestion, cough (similar to viral pneumonia or bronchiolitis)
 - c. Children: +/- fever, URI symptoms, GI symptoms such as diarrhea
8. Recent reports have identified children presenting with symptoms of a multisystem inflammatory syndrome with features overlapping Kawasaki disease, toxic shock syndromes and myocarditis (see below for details).
9. Most laboratory results, to include inflammatory markers (ESR, CRP), chemistries, kidney and hepatic function are normal. White blood cell count is typically normal but may be low. Procalcitonin may be elevated but might suggest co-infection.(79)
10. When imaging is abnormal in children with COVID-19, CXR reveals non-specific increased lung markings or patchy infiltrates, and chest CT reveals glass opacities and halo signs.(79)
11. Pediatric patients can be considered mild or moderate disease if there is no new supplemental oxygen requirement or no increased requirement for patients who require supplemental oxygen at baseline. A majority of these patients will self-resolve without intervention.
12. Those who require hospitalization should receive supportive care to include critical care interventions as required.
 - a. Respiratory Support
 - i. There is no current evidence to alter treatment of severe respiratory failure from standard pediatric ARDSNet guidelines (2005), including indications for intubation and use of non-invasive respiratory

Clinical Management of COVID-19, v4

support although increased use of prone positioning is recommended if tolerated.

- ii. Use of viral filters on circuits are necessary including for side sampling ET_{CO₂}; consider effects on flow dynamics of added resistance for smaller patients.
- iii. Judicious sedation and/or neuromuscular blockade for intubated patients should be considered given risk of rapid decompensation and self-extubation with delayed provider response time while donning PPE.
- iv. Similar CT findings as in adults are expected for severe cases although often unhelpful to guide clinical practice.
- v. For mild/moderate patients requiring nasal cannula/mask, goal is to target SpO₂ >94% during resuscitation, and >90% once stable. For flows over 3 L/min use of a heated/humidified circuit (HHFNC) or non-invasive positive pressure ventilation (CPAP or BiPAP) are well tolerated, but its use will increase aerosolization.

b. Shock

- i. Recognize the multisystem inflammatory syndrome in children (MIS-C). Note this may also occur in young adults. Further discussion is below.
- ii. Recognize septic shock in children with any hypotension (systolic blood pressure [SBP] < 5th percentile or > 2 SD below normal for age) or two or more of the following: altered mental state; bradycardia or tachycardia (HR < 90 bpm or > 160 bpm in infants and HR < 70 bpm or > 150 bpm in children); prolonged capillary refill (> 2 sec) or weak pulses; tachypnea; mottled or cold skin or petechial or purpuric rash; increased lactate; oliguria; hyperthermia or hypothermia. Mental status is often preserved in older children and adolescents.
- iii. For septic shock in children, give 10–20 mL/kg crystalloid fluid as a bolus as quickly as possible using a manual push and reassess for signs of fluid after each bolus. Evaluate for cardiogenic shock in unresponsive/worsening patients and use of vasoactive medications with the development of hepatomegaly, pulmonary edema or elevated CVP.(161)
- iv. Resuscitation endpoints include perfusion targets (e.g. urine output > 1 mL/kg/hr in children, improved level of consciousness and perfusion, resolving lactate or improvements in clinical indicators (as measured by advanced monitoring: CVP, cvSaO₂, cardiac index etc.).
- v. In children, consider epinephrine for first-line treatment, while norepinephrine can be added if shock persists or primarily ‘warm’ shock. Milrinone is appropriate for use in diagnosed impaired cardiac contractility, if patient is no longer hypotensive.(161)

c. Antivirals Use – for severe patients only

- i. Enrollment in clinical trials or compassionate use of experimental therapies to include antivirals, should be considered for children with severe disease on a case-by-case basis with appropriate monitoring and in consultation with Peds ID when possible.(228)
- ii. Remdesivir is available from the manufacturer for children <18 years-old as compassionate use and for children ≥ 12-years old as part of randomized clinical trial (see Adjunctive Therapies section above for more information).
 - Dosing recommendations from Pediatric Infectious Disease Society (verify with manufacturer)
 - <40kg: 5mg/kg IV loading dose on day 1; followed by 2.5mg/kg IV Q24hr
 - ≥40kg: 200mg IV loading dose on day 1; followed by 100mg IV Q24hr
 - Recommended duration: up to 10 days with 5-day duration favored for fast responders
- iii. There is emerging data and concerns about safety and efficacy of hydroxychloroquine for treatment of COVID-19. Use outside of approved clinical trials is not recommended.

13. There is no evidence to suggest that prophylaxis is necessary or effective for the majority of children.

14. Children appear to efficiently shed the virus, even if asymptomatic. Average duration of shedding is 11 days for asymptomatic and 17 days for symptomatic.(65, 229)

15. Given the prolonged duration of shedding of respiratory viruses in children, during periods of community transmission of SARS-CoV-2, it may be prudent to assume symptomatic children are infected, unless proven otherwise from an infection control standpoint - an issue particularly relevant to caregivers from vulnerable risk populations.

Multisystem Inflammatory Syndrome in Children (MIS-C) associated with COVID-19

1. MIS-C is a novel syndrome that appears temporally associated with COVID-19. It consists of fevers and a variety of clinical findings similar to incomplete/atypical Kawasaki, and high inflammatory markers. Up to 40% are PCR positive for SARS-CoV-2, but more than 80% have positive serology for SARS-CoV-2 indicating preceding infection. The peak incidence of this syndrome occurs 4-8 weeks after peak COVID disease in local community, suggesting a post-infectious etiology. The majority of patients have some cardiac involvement requiring ICU-level care. Due to severity of disease, it is of utmost importance for early recognition and transfer to tertiary care center. Awareness and communication of the risk of this disease in child and young adult populations. See **Appendix N** for DHA summary communication regarding MIS-C.
2. CDC Case Definition for MIS-C
 - a. An individual aged < 21 years (editors note: older patients have been reported) presenting with all the following:
 - i. Fever 38°C for ≥24hrs or report of subjective fever lasting ≥24hrs **AND**
 - ii. Laboratory evidence of inflammation* **AND**
 1. * including, but not limited to one or more of the following: elevated CRP, ESR, fibrinogen, procalcitonin, d-dimer, ferritin, LDH, IL-6, elevated neutrophils, reduced lymphocytes and low albumin
 - iii. Evidence of clinically **severe illness** requiring hospitalization, with multisystem (>2) organ involvement (cardiac, renal, respiratory, hematologic, gastrointestinal, dermatologic or neurological) **AND**
 - iv. No alternative plausible diagnoses **AND**
 - v. Positive current or recent SARS-CoV-2 infection by RT-PCR, serology, or antigen test; or COVID-19 exposure within the 4 weeks prior to the onset of symptoms
3. The spectrum of illness in MIS-C is emerging, but at present the clinical presentation can include: (230-236)
 - a. Fever (100%)
 - b. Hypotension (50-100%)
 - c. Diarrhea (60-100%)
 - d. Conjunctivitis (20-76%)
 - e. Mucosal changes such as pharyngeal erythema, fissured/cracked lips (40-54%)
 - f. Rash (50-57%)
 - g. Neuro symptoms such as headache, irritability, stiff neck, vision changes (43-58%)
 - h. Swelling of hands/feet (35-50%)
 - i. Other: lymphadenopathy, myalgias, fatigue, cough, dyspnea
4. Patients with signs and symptoms compatible with MIS-C, should be evaluated for admission in an emergency department setting. Initial evaluation should include:
 - a. Stabilization using PALS algorithms to optimize hemodynamics and respiratory support
 - i. Judicious fluid management due to potential cardiac involvement
 - ii. Attention to COVID infection prevention & control measures.
 - b. EKG, chest x-ray, and bedside cardiac ultrasound to evaluate function (if available)
 - c. Initial Labs (performed at initial point of care)
 - i. SARS-CoV-2 RT-PCR (NP specimen preferred)
 - ii. Respiratory pathogen PCR (e.g. Biofire filmarray)
 - iii. Rapid strep and throat culture
 - iv. Blood culture
 - v. Urinalysis with urine culture
 - vi. CBC with differential
 - vii. CMP, Magnesium, Phosphorus
 - viii. CRP, ESR, Procalcitonin
 - ix. Ferritin, LDH
 - d. Transfer child to a military or civilian **tertiary medical center** with PICU and peds specialists (including rheumatology, immunology, cardiology, infectious disease and heme/onc) if any of the following:

Clinical Management of COVID-19, v4

- i. Clinical Concern for Kawasaki disease or MIS-C
- ii. Child is ill-appearing or has PEWS score >4
- iii. Hypoxia or shock present
- iv. Abnormal EKG or point of care echo
- e. Once at a tertiary medical center w/ pediatric subspecialists and concern for MIS-C, obtain echocardiogram and consider additional labs:
 - i. SARS-CoV-2 serology (if not yet confirmed)
 - ii. Troponin (high sensitivity), Pro-B Natriuretic Peptide (pro-BNP), CK
 - iii. Coagulation panel to include D-dimer; von Willebrand Factor Antigen
 - iv. Triglyceride, Amylase
 - v. IL-6, Cytokine panel (typically a send out; can use Cincinnati Children's)
 - vi. CSF studies if signs/symptoms of meningitis
 - vii. Rickettsial serologies if exposure to endemic regions
 - viii. Save 5-7mL blood in EDTA tube and 5mL serum prior to any immune modulating treatment such as IVIG for future diagnostic studies.
- f. Initial treatment decisions should be made with a multidisciplinary approach as the most effective standard of care has not yet been determined and there is clinical overlap with other infectious and inflammatory conditions.
 - i. General
 1. Judicious fluid management due to potential cardiac dysfunction
 2. Close hemodynamic and electrolyte monitoring.
 - a. to prevent dysrhythmia, goal Mag = 2, Phos = 4
 - ii. Empiric antibiotics – early discontinuation if no evidence of bacterial infection
 1. Clindamycin and Ceftriaxone for Toxic Shock Syndrome
 2. +/- Vancomycin if concerns about MRSA
 3. +/- Doxycycline if concerns for *Rickettsia*
 - iii. Anti-Inflammatory – based on input from Peds Cardiology, Rheumatology, Immunology
 1. IVIG 2g/kg over 8-12hrs; ideally within 7-10 days of fever onset
 2. High-dose aspirin (20mg/kg Q6hr)
 3. additional immune blockade with agents such as methylprednisolone or biopharmaceuticals such as anakinra with expert consultation.
 - iv. Anticoagulation
 1. Enoxaparin (Lovenox) with Peds Heme/Onc or Critical Care consultation

Sustaining Pediatric Preventive Medicine Services During the Pandemic

The Centers for Disease Control and Prevention (CDC) and the American Academy of Pediatrics (AAP) both strongly support the sustainment of well child and preventive health care during the pandemic (citations: AAP COVID-19 website; CDC Information for Pediatric Healthcare Providers website). Of highest priority are the prevention of secondary outbreaks of vaccine-preventable illnesses, newborn follow-ups, and developmental surveillance. Local MTFs should plan for and be prepared to implement pediatric care in accordance with CDC and AAP recommendations, in the setting of DHA guidance based upon local Health Protection Condition (HPCON) levels (citation: Memorandum Resuming Full Operations_ADHCA signed). These plans may include utilization of telehealth options, adjustments to patient flow, prioritization to continue to immunize the highest risk groups, and establishing database tracking mechanisms to proactively communicate with families to ensure that immunizations and overdue care is scheduled and provided for as soon as possible as local conditions and HPCON levels allow.

References:

CDC Information for Pediatric Healthcare Providers, May 2020 [Available from: <https://www.cdc.gov/coronavirus/2019-ncov/hcp/pediatric-hcp.html>]

American Academy of Pediatrics Guidance on Providing Pediatric Well-Care During Covid-19, May 2020 [Available from: <https://services.aap.org/en/pages/2019-novel-coronavirus-covid-19-infections/guidance-on-providing-pediatric-well-care-during-covid-19/>]

Memorandum Resuming Full Operations_ADHCA signed, May 2020 [Available from: <https://www.milsuite.mil/book/docs/DOC-810764>]

Caring for Older Persons with COVID-19

1. COVID-19 can result in severe disease and death among older adults. In the United States, 8 out of 10 deaths have been in adults above age 65. Mortality rates in patients > 85 have ranged 10-27%, and 4-11% among patients 65-84 years.(237)
2. Older adults, especially those that are frail and have multiple comorbidities, may not present with the typical syndrome of fever, fatigue, or cough. Atypical presentation of disease includes tachypnea, delirium, malaise, myalgias, and diarrhea early in the disease course; fever was not as prominent in several cases.(238)
3. Have a high index of suspicion for COVID-19 in those patients not at their baseline, especially those residing in long term care facilities who present with respiratory difficulties, changes in vital signs other than temperature or other signs of infection or sepsis.
4. Ensure that care for the older adult and severely ill is in keeping with their goals of care, advance directives and patient and family wishes.
5. Conversations regarding goals of care should continue to be part of routine care.
6. Patients should be informed about their condition and their prognosis (if desired), in a way easy to understand.
7. If the patient is unable to communicate meaningfully, ensure that a surrogate decision maker or health care agent has been identified in accordance with state law based on facility location.
8. All providers should provide basic symptom management, perform routine discussions about goals of care and code status in seriously ill patients. If complex symptom management or difficult discussions surrounding goals of care or code status arise, consult a palliative medicine subspecialist if available at your institution.
9. Symptom management: Aggressive control of symptoms such as pain, dyspnea, or other symptoms relieves unnecessary suffering, which is crucial for all patients regardless of age, function, comorbidities and prognosis.
 - a. *Pain*
 - Acetaminophen should be used first, typically 500mg every 6 hours as needed.
 - If acetaminophen is insufficient, and other modalities such as topical agents are ineffective, start an opioid for moderate to severe pain (drug, dose, route, and frequency should be individualized and based on symptom severity, kidney/liver function and prior opioid exposure: See Table 7). Consider local supply in drug selection to mitigate risk of drug shortage.
 - Start a stimulant laxative, such as Senna 8.6mg PO daily, if prescribing an opioid to prevent constipation. Titrate to effect. Escalate bowel regimen as needed, with a goal of one soft bowel movement at minimum every other day.
 - b. *Dyspnea*
 - If providing supportive care and supplemental oxygen is ineffective for management of severe dyspnea, a low-dose opioid may be used to help alleviate symptoms.
10. Communication challenges may be exacerbated by the use of PPE. In patients with sensory impairments it is important to remember to eliminate or minimize background noise, state information slowly, and avoid yelling. It may be helpful to display information in writing. Hearing aids/glasses should be worn if available.
11. Older adults, especially those with cognitive impairment, when ill, hospitalized, or placed in a new environment may become anxious, agitated or less interactive. Delirium, a diagnosis not exclusive to older adults, manifests as acute onset inattention, disorganized thinking and an altered level of consciousness. Delirium may be seen any patient, especially those with severe infection, and those requiring mechanical ventilation. Hyperactive delirium (delirium with agitation) may make management and risk mitigation challenging in those diagnosed with COVID-19.
 - a. Early recognition and management of delirium is important. Regular delirium screening should occur using validated methods such as the Confusion Assessment Method, bCAM, or the 4AT (www.the4AT.com). (239, 240)
 - b. Risk factors for delirium include older age, sensory impairment (vision and hearing), history of dementia, patients admitted from long-term care units, and those with serious infection.(241)
 - c. Management of Delirium: (242, 243)
 - Prevention of delirium is the best strategy. Strategies include maintaining normal circadian rhythms, exposure to natural light, regular reorientation, mobilization, treating pain, fever, and nausea, maintaining oxygenation, avoiding constipation and urinary retention, and performing medication

Clinical Management of COVID-19, v4

reconciliation to minimize potentially inappropriate medications. Ensure basic needs are met for food and water.

- Standard non-pharmacological approaches such as frequent reorientation, family at bedside, hospital environmental manipulation (maintenance of day/night cycle, appropriate use of TV and lights), calming music, calls from family, and professional sitters should be employed but may not be feasible in an isolation setting.
- In patients with hyperactive delirium, try nonpharmacological techniques first.
- Current evidence does not support routine use of antipsychotics in management of delirium.(244)
- If severe agitation occurs, and nonpharmacological approaches have not been effective or more rapid control is needed for the safety of the patient or others, antipsychotics may be used but are off-label. When using an antipsychotic, use the lowest effective dose for the shortest amount of time. Of note, all antipsychotics carry a FDA Black Box warning due to an increase in mortality when used in patients with dementia. The patient should be monitored closely for side effects such as QTc prolongation and over sedation.

- Some examples of antipsychotics are Quetiapine 25mg - 50mg PO, Olanzapine 2.5mg - 5mg PO/IM, and Haldol 0.25mg -1mg IV

- d. Cautious use of antipsychotic medication is needed especially in patients with movement disorders such as Parkinson’s disease and Lewy Body Dementia as this class of medication may exacerbate extrapyramidal symptoms. Quetiapine is preferred if antipsychotic medications are needed in patients with movement disorders given its lower risk of extrapyramidal symptoms.(245) Any patient is at risk for acute dystonic reaction to antipsychotic medications.

12. Many older adults will recover from their illness, and it is important to not forget other complications such as hospital-associated deconditioning, falls and wounds. Standard of care should be provided for these other common complications alongside supportive care for COVID-19. Prompt mobilization and therapy should be started, when able, in accordance with infection control practices. Focusing on other treatable conditions should continue alongside supportive care for COVID-19.

PALLIATIVE MEDICINE DURING THE COVID-19 PANDEMIC

Palliative medicine can assist at all stages of contingency/crisis planning. Prepare for increased use of symptom management resources including opioids (morphine IV and PO, hydromorphone IV and PO, oxycodone PO, fentanyl IV and transdermal), and benzodiazepines. Consider dedicated space for end-of-life care beds. Where possible, symptom management resources should be de-conflicted with highly utilized intensive care medications use to prevent and adapt to shortages.

Goals of Care Discussions

(Adapted from vitaltalk.org COVID-19 Open Source Resources. www.vitaltalk.org)

1. Eliciting a patient’s goals of care is integral to providing the best and most appropriate medical care and can improve resource allocation during a time of scarcity. Engage patients proactively in goals of care discussions informed by personal values and clinical context.
2. Treat patients and their families with respect and compassion. Quickly and effectively elicit a patient’s concerns, values, and preferences with a few key statements. Table 7 offers suggestions and examples to help guide your conversations.

Table 7. Difficult Conversations and Scripts for Communicating with Patients and Families

What the patient/family says	What you may say
Admitting a Patient	
How bad is this?	<ul style="list-style-type: none">• From the information I have now and from my exam, your situation is serious enough that you should be in the hospital. We will know more in the coming hours to days, and we will update you. Who else should know about your/their situation and how will they know?
Is my grandfather going to make it?	<ul style="list-style-type: none">• I imagine you are scared. Here’s what I can say: because he is 90, and is already dealing with other illnesses, I worry that he is at risk of dying if this worsens in the hospital. While it is too soon to say for certain, what worries you most about that?
Are you saying that no one can visit me?	<ul style="list-style-type: none">• I know it is hard to not have visitors. The risk of spreading the virus to other vulnerable people is so high that they and those they contact will be in more danger if they come into the hospital. I wish things were different.

How can you not let me in for a visit?	<ul style="list-style-type: none"> The risk of spreading the virus is so high that I am sorry to say we cannot allow visitors. We can help you be in contact electronically. <i>I wish I could let you visit, because I know it's important, but it is not possible now.</i>
When things aren't going well, goals of care discussion, code status discussions	
I want everything possible. I want to live.	<ul style="list-style-type: none"> We are doing everything we can. This is a tough and scary situation for many of us. Could we step back for a moment so I can learn more about you? <i>What do I need to know about you to do a better job taking care of you?</i>
I don't think my grandfather would have wanted this.	<ul style="list-style-type: none"> Well, let's pause and talk about your concern. <i>Can you tell me what we should know to take the best care of him?</i>
I don't want to end up being a vegetable or on a machine.	<ul style="list-style-type: none"> Thank you, it is very important for me to know that. <i>Can you say more about what you mean?</i>
I am not sure what my grandfather wanted – we never spoke about it.	<ul style="list-style-type: none"> You know, many people find themselves in the same boat. This is a hard situation. To be honest, given his overall condition now, I worry that further treatments may not be successful in preventing him from dying. <i>In a situation like that, I have recommended that we allow a natural death.</i> That could be hard to hear. What do you think?
When coping needs to be boosted, or emotions are running high	
I'm scared.	<ul style="list-style-type: none"> This is such a tough situation. <i>I think anyone would be scared.</i> Could you share more with me?
I need some hope.	<ul style="list-style-type: none"> Tell me about the things you are hoping for? <i>I want to understand more.</i>
You people are incompetent!	<ul style="list-style-type: none"> I can see you are not happy with things. <i>I am willing to do what is in my power to improve things for you.</i> What could I do that would help?
I want to talk to your boss.	<ul style="list-style-type: none"> I can see you are frustrated. <i>I will ask my boss to come by as soon as they can. Please realize that they are juggling many things right now.</i>
Do I need to say my goodbyes?	<ul style="list-style-type: none"> I'm hoping that's not the case and I worry time could indeed be short. <i>What is most pressing on your mind?</i>

Symptom Management Guidelines

(Adapted from BC Centre for Palliative Care COVID-19 Resources and Information, bc-cpc.ca/cpc)

1. Patients with COVID-19 infections experience many of the same symptoms as other patients: dyspnea, oral secretions, anxiety and pain. Symptom management should be individualized based on clinical status.
 - a. **Dyspnea – dyspnea can present as anxiety – treat the dyspnea!**
 - Non-pharmacologic management for shortness of breath:
 - Positioning, cool room temperatures, removing restrictive clothing
 - Avoid bedside fan for patients with COVID-19. Consider bronchodilator therapy, fluid overload therapies, and heart rate control if >120 BPM. h:
 - Opioids are the mainstay of comfort care in severe dyspnea. When dosed effectively to control dyspnea, they **do not** contribute to a hastened death.
 - Treat and reassess. IV opioids works within 10-15 min, oral opioids within 30-45 min.
 - Goals for treatment: respiratory rate <25, minimal use of accessory muscles, resolution of pursed lip breathing, nasal flaring, and retractions or subjective dyspnea. Patient comfort is the goal.
 - See Table 8 for recommended opioid dosing. If the dose does not work, increase it!
 - b. **Respiratory Secretions/Congestion Near End of Life**
 - Discuss congestion and secretions with family and bedside staff. Pharyngeal secretions are normal at end of life and rarely require treatment. A productive cough may benefit from mucolytics or opioids (Table 7). Limit oropharyngeal suction. Reduce or stop saline infusions.
 - Medications may include:
 - **Glycopyrolate** 0.4 mg SQ/IV q4H PRN
 - If severe and refractory to above medications, consider:
 - **Furosemide** 20 mg SQ/IV q2h PRN with close monitoring of response.
 - c. **Anxiety**
 - Patients with dyspnea have associated fear and anxiety-- opioids are the first line of treatment. The following adjuncts may be helpful in refractory anxiety:
 - **Lorazepam** 0.5 – 1 mg PO/IV q1-4H PRN, consider scheduling Q4H if goals are for comfort-directed care and the patient is requiring frequent PRN dosing.
 - **Midazolam** 1 – 4 mg SQ/IV q30min PRN, consider scheduling Q4H if goals are for comfort-directed care and the patient is requiring frequent PRN dosing. (*for severe anxiety or SOB in ICU*)
 - d. **Delirium**

Clinical Management of COVID-19, v4

- Delirium, either hypoactive or agitation, is common in hospitalized patients and can be distressing. Avoid benzos. Treat underlying causes of delirium if possible.
- **Haloperidol** 0.5 mg PO **OR** 0.5 – 1 mg IV q4H PRN. Consider scheduling the medication Q4H if requiring frequent PRN dosing. Titrate dose in 0.5mg increments.
- **Olanzapine** 2.5 – 5 mg PO qHS and q8 hr PRN. This comes as a regular or oral dissolving tablet and can be titrated.

Table 8. Opioid Dosing to Relieve Dyspnea and Pain in Adults

Intermittent Dosing													
Dosing for Opioid Naïve Patient (patient not on opioid therapy) (For frail, elderly patients, begin at low end of any range)													
Morphine	<ul style="list-style-type: none"> • 15 mg tablet ½ to 1 tab PO q 3 hours prn OR 5 mg SQ/IV q1H PRN shortness of breath (SQ/IV can be given as frequently as q30min PRN) 												
Hydromorphone	<ul style="list-style-type: none"> • 2 mg tablet ½ to 1 tab PO q 3 hours prn OR 0.4-0.8 mg SQ/IV q1H PRN shortness of breath (SQ/IV can be given as frequently as q30min PRN) • If more than 6 PRN doses of opioid in 24 hours: Consider a basal opioid such as MSContin 15 mg PO BID. If patient unable to make needs known, consider SCHEDULED dosing of the immediate release opioid (q4H or 6H for frail elderly) AND continue PRN dose. <p style="text-align: center;">TITRATE UP AS NEEDED for relief of dyspnea and/or pain</p>												
Dosing for Patients ALREADY Taking Opioids													
Applies to any opioid	<ul style="list-style-type: none"> • Continue previous opioid, consider increasing dose by 25% • To manage breakthrough symptoms: Start PRN opioid at 10% of total daily (24 hour) opioid dose. • PRN q1H for PO and q30mins for SQ/IV 												
PCA Infusion Dosing: For alert patients who need IV opioids (unable to take PO or with severe symptoms)													
PCA Infusion Pump Dosing for Opioid Naïve Patient NOT Already Taking Opioids													
	<table border="1"> <thead> <tr> <th>Opioid</th> <th>Bolus Dose</th> <th>Basal Rate (if severe symptoms)</th> </tr> </thead> <tbody> <tr> <td>MORPHINE</td> <td>1.5 mg q10mins</td> <td>1-2 mg/hour</td> </tr> <tr> <td>HYDROMORPHONE</td> <td>0.2 mg q10mins</td> <td>0.1 – 0.3 mg/hour</td> </tr> <tr> <td>FENTANYL</td> <td>20 micrograms q10mins</td> <td>10-25 micrograms/hr</td> </tr> </tbody> </table>	Opioid	Bolus Dose	Basal Rate (if severe symptoms)	MORPHINE	1.5 mg q10mins	1-2 mg/hour	HYDROMORPHONE	0.2 mg q10mins	0.1 – 0.3 mg/hour	FENTANYL	20 micrograms q10mins	10-25 micrograms/hr
Opioid	Bolus Dose	Basal Rate (if severe symptoms)											
MORPHINE	1.5 mg q10mins	1-2 mg/hour											
HYDROMORPHONE	0.2 mg q10mins	0.1 – 0.3 mg/hour											
FENTANYL	20 micrograms q10mins	10-25 micrograms/hr											
Titrate the basal rate and bolus dose to effect. If using more than 1 rescue dose/hour, increase the basal rate for improved symptom control.													
PCA Infusion Pump Strategy for Patient ALREADY Taking Opioids													
<ul style="list-style-type: none"> • For patients on chronic opioid therapy, rotate their long acting medication into the basal rate of your PCA. Titrate to effect. • Bolus doses may be given q10 to 15min PRN; if the patient is NOT able to use the button, add a nurse administered bolus order of 5 mg IV q 2 hour prn for morphine PCAs and 0.8 mg IV q 2 hour prn for hydromorphone PCAs. • Example titration: You start a morphine PCA at 1 mg/hr basal rate with 1 mg q 15 minutes rescue. The patient presses the button every 15 minutes and says he “feels nothing” and continues to be short of breath. Increase the rescue dose to 2 mg and reassess. • Adjust bolus doses to 50-100% of new continuous infusion rate (e.g. Bolus dose of 2-4 mg q15min PRN for new rate of 4mg/h). • New rate can be reassessed for adjustment again in 3-4 hours. 													

e. Constipation

- Use of opioids will cause constipation. If the patient has more than 24 hours to live:
 - Start a stimulant laxative, such as Senna 8.6mg PO daily if they are tolerating PO.
 - PRN enema if unable to take PO and patient uncomfortable from distention
 - Escalate bowel regimen as needed, with a goal of one soft bowel movement at minimum every other day.

Palliative Ventilator De-Escalation

(Adapted from “Palliative Ventilator De-escalation Recommendations for COVID-19 Positive or PUI. Developed by Bartlett, Christi for The University of Kansas Health System)

The following protocol is assumed to take place after appropriate goals of care discussions with family and/or surrogate decision makers. The endotracheal tube will remain in place and the ventilator circuit will remain intact to reduce the risk of COVID-19 exposure.

Pre-Procedure.

1. Prepare family that prognosis can be as short as a few minutes but as long as a few days.
2. Deactivate defibrillators first. A magnet can also be placed over the device if needed to deactivate.
3. Ensure no paralytic medications are on board.
4. Code status should be DNR/ comfort measures only for patients at the end of life.
5. Discontinue tube feeds and remove unnecessary equipment from the room.

Procedure.

1. Turn off alarms and change room monitor to comfort care setting or turn off if family is present.
2. If a continuous opioid infusion is in place, continue THE SAME medication. All opioids contribute to relief of pain/dyspnea.
3. If the patient is already on a continuous opioid infusion, double current drip rate and order bolus doses of 100-200% of drip rate to be given q10min PRN. Use bedside infusion to provide boluses whenever possible.
4. If the patient is opioid naïve and not on a continuous infusion, begin with morphine 5mg IV or hydromorphone 0.5 – 1 mg IV q10min PRN. If possible, bring at least four doses into patient room for ventilator de-escalation.
5. Order midazolam 2-4 mg q10min PRN or lorazepam 2 mg q30min PRN for anxiety/breathlessness. If patient is already on a midazolam continuous infusion, double current rate and give boluses of 100-200% of drip rate available q10min PRN. Use bedside infusion to provide boluses PRN.
6. Pre-medicate with an opioid bolus as above (100% of drip rate) 10 minutes prior to de-escalation.
7. Pre-medicate with 2 – 4 mg of midazolam 10-15 min or 1 – 2 mg of IV lorazepam or prior to de-escalation.
8. Recommend glycopyrrolate 0.4 mg IV q4H PRN for secretions.
9. If patient requires sedative medication (propofol, precedex, etc) for comfort, continue as ventilator is weaned.
10. Stop vasopressors prior to weaning ventilator.
11. Ensure that patient appears comfortable prior to reducing ventilator settings. Titrate to comfort to palliate signs of discomfort: grimacing, agitations, or labored respirations.
12. For agitation/delirium management, consider Haloperidol 0.5 – 1 mg IV q30mins PRN.
13. If patient is obtunded and expected to die abruptly after ventilator is weaned, recommend immediate reduction in ventilator settings to pressure support 5/5 and room air. Bolus opioid and benzodiazepine aggressively as needed to ensure comfort.
14. If the patient is alert, consider a gradual reduction in ventilator settings. Decrease FiO₂ to 40%, PEEP to 10, RR to 16. Recheck patient comfort and re-bolus opioids prn to achieve comfort. Reduce PEEP to 5, FiO₂ to 0.21.
15. Once the ventilator is set at PS 5/5 and FiO₂ of 21%, leave endotracheal tube in place and leave the ventilator circuit intact for the end of life.
16. Continue to re-bolus opioids, benzodiazepines and sedation as needed to ensure comfort.

IMPLICATIONS OF SURGICAL AND INVASIVE PROCEDURES

Priorities for Surgical Resources

Force Protection: Protection of personnel and patients from disease transmission

Mission Capability: Maintaining capability to provide safe and effective surgical care

Mission Support: Support of the healthcare community response to COVID-19 through preservation of critical resources and re-deployment of personnel

Triage and Decision-making

The ability to provide surgical care should be determined by health protection conditions, local and regional healthcare capability and capacity with consideration of logistic constraints.

1. During sustained or widespread community transmission, surgical care should be restricted to reduce risks of transmission between patients and healthcare personnel. To the extent possible, clinical encounters should be accomplished through virtual means and surgical treatment options deferred or delayed.(246)
2. MTFs should establish a review process to triage and prioritize medically necessary and time-sensitive surgical care.(247) This process should include multidisciplinary representation and be led by a senior surgeon, preferably the Department/Service Chief.(248)
 - a. This review should consider medically necessity, time sensitivity, risk and impact of viral transmission to either the patient or medical personnel, suitability of alternative treatment options, resource utilization, impact of delay of treatment, as well as readiness and mission impact.
 - b. Consider using an acuity scale or scoring system to assist in decision making.(249)
3. Preoperative COVID-19 testing is recommended to assist in decision-making for all surgical patients including symptomatic and asymptomatic. In the event of a positive test, the surgical treatment plan can be

Clinical Management of COVID-19, v4

reconsidered to reduce patient risk of morbidity and mortality and to reduce the risk of transmission to medical personnel and the community. Surgical teams and their patients should have access to preoperative testing to ensure adequate information is available to determine the best treatment strategy

- a. Treatment facility preoperative testing strategies should consider local prevalence of disease and the availability and performance of testing capability. Testing is expected to be most beneficial if performed within 48 hours of the surgical procedure. Recommendations for prioritization of testing are as follows:
 - i. All patients with symptoms suggestive of COVID-19.
 - ii. High-risk procedures such as head & neck, thoracic, and upper gastrointestinal surgery.
 - iii. Surgical procedures/patients with anticipated requirement for intensive care and/or prolonged hospitalization.
 - iv. Surgical procedures requiring inpatient postoperative care.
 - v. Outpatient procedures on patients whose age or comorbid conditions suggests a high-risk of morbidity or mortality from COVID-19.
 - vi. Routine outpatient procedures.
 - b. MTF policies and procedures regarding preoperative testing must balance the desire to support safe, high-quality surgical care with efficient operations. Preoperative testing policies should not adversely impact capability to provide emergent and urgent surgical care.
4. In areas with low incidence or sustained reduction in the rate of new COVID-19 cases, expansion of surgical services should be considered.(250) Surgical services must adapt traditional and contingency operations into a new normal of patient care in the setting of ongoing COVID-19 risk. The surgical review and triage process should continue to prioritize surgical care as outlined in 2a above but can apply a progressively lower threshold to proceed with surgical care and utilization of healthcare resources.
5. Prior to resumption of elective surgery, the following should be established:
- a. Local Objective Triggers: Transition of an MTF to the next phase of medical activities should be guided by the local Health Protection Condition (HPCON) level or guidance from local or state governments or from the Uniformed Military Department or installation commander, in addition to DHA recommendations based on coordination in key healthcare-sustaining areas discussed below. Triggers at the local level are:
 - i. Symptoms: Downward trajectory of influenza-like illnesses (ILI) reported within a 14-day period; and a downward trajectory of COVID-like cases reported within a 4-day period.
 - ii. Cases: Downward trajectory of documented cases within a 14-day period or a downward trajectory of positive tests as a percent of total tests within a 14-day period (flat or increasing volume of tests).
 - iii. Hospitals: Treat all patients without crisis care; and robust testing program in place for at-risk HCP.
 - b. Sufficient resources are available across phases of care, including PPE, healthy workforce, facilities, supplies, testing capacity, and post-acute care.
 - i. Performance of elective surgery must not negatively impact capability to provide medically necessary and time sensitive surgical care.
 - ii. Surge capacity should be preserved.
 - c. Policies and process are established for perioperative screening and testing of surgical patients.
 - d. Evidence-based infection prevention policies and procedures are established to ensure a safe environment in which elective surgery can occur. (i.e., access control, workflow and distancing)
 - e. Non-COVID Care areas should be established to reduce risk of COVID-19 exposure and transmission; preferably these areas should be separate from other facilities to the extent possible (i.e., separate building, or designated rooms or floor with a separate entrance and minimal crossover with COVID-19 areas).
 - f. Establish policies, process and content for patient education on COVID related care risks and the risk mitigation strategies employed to ensure their safety.(251)
 - g. CONUS: The guidelines for “Opening Up American Again” based on Centers for Medicare & Medicaid Services recommendations offer the following 3 phases:(252)

Clinical Management of COVID-19, v4

- i. Phase 1: Elective surgeries and procedures can resume, as clinically appropriate, on an outpatient basis
- ii. Phase 2: Elective surgeries and procedures can resume, as clinically appropriate, on in-patient bases
- iii. Phase 3: Patient care as indicated. Visits to hospitals can resume. Those who do interact with patients must adhere to strict hygiene protocols.

Phases of Surgical Care Recommendations

Preoperative Care.

1. Virtual and telehealth should be utilized to accomplish preoperative administrative tasks, education, and assessments that do not require face to face interaction.
2. Post-operative care needs should be assessed and resource availability confirmed.
3. Patients planned for surgical care should be screened for symptoms or exposure history. Those patients that screen positive should undergo testing and their surgical treatment plan should be reconsidered.
4. Consider use of a Facility Readiness Checklist and/or Patient Information Sheet.
5. When available, preoperative COVID testing should be performed to identify asymptomatic patients whose surgical care plan can be altered in the event of a positive test result. Timing of the test should balance considerations regarding the availability and turnaround time of test against the risk of patient exposure and infection in the interval between testing and the scheduled procedure.

Immediate Preoperative Care.

1. Recommend establishing Intubation/Extubation Airway Teams consisting of providers with a high degree of comfort with PPE and airway skills. Teams should bring their own PPE, medications, and airway equipment to avoid delays while limited or unfamiliar PPE is made available. During the pandemic, any emergency airway should be treated as potentially COVID-19 positive and full PPE worn.
2. For purposes of perioperative care, patients should be treated as presumed COVID-19 positive if they have symptoms/exposure history that warrants testing. PUIs at MTFs without an urgent indication for surgery preferably are tested for COVID-19 before any operative intervention (provided testing availability).
3. Optimally, an OR or pod of ORs should be predesignated with a distinct anteroom to maintain separation from non COVID-19 patients. Negative pressure is not recommended for operating rooms. Consider reducing positive pressure and using a HEPA filtration system. Consult with facilities to ensure air handling is routed through the HEPA filter (i.e., air scrubber). An air scrubber is a portable filtration system that removes particles, gasses, and/or chemicals from the air within a given area. These machines draw air in from the surrounding environment and pass it through a series of filters to remove contaminants.
4. All patient interaction with COVID-19 positive or PUI patients will be performed with airborne and contact precautions, including eye protection:
 - a. N95 mask with surgical mask over the N95 mask, consider PAPR for aerosol generating procedures.
 - b. Eye protection consisting of goggles, full face shield/mask worn over N95, or plastic disposable wrap-around glasses. Eyeglasses alone are not adequate.
 - c. Gown, double gloves, hair cover, shoe covers
 - d. Remove PPE except N95 mask before exiting the room. Surgical scrubs should be changed after each case.
5. The anesthesia service provider should attempt to remove all necessary medications and equipment from the carts prior to bringing the patient into the room. Avoiding contamination of the carts/machine should be prioritized over wasting consumable supplies.
6. Anesthesia service providers should not expect routine breaks during the case. Consider leaving cell phones, smart watches, and other personal devices out of the OR. Ensure there's a way to communicate/call for assistance organic to the OR. Recommend additional support staff immediately available outside the OR to assist with providing requested medications and supplies to the operating room team.
7. Patients on the ward should be transported directly to the OR by the anesthesia service team. If assistance is needed with transport, every attempt should be made to enroll a member from the care team (nurse, surgeon, and technician) to minimize staff exposure.

Clinical Management of COVID-19, v4
Intraoperative Care. (253, 254)

1. Surgeons and non-essential staff should not be present in the OR for either intubation or extubation unless necessary for patient safety. Exposure risks after these airway procedures is affected by risk mitigation strategies and engineering controls (airflow and filtration); therefore OR workflow and staff entry after airway manipulation should be adjusted based on a thorough understanding of these factors.
2. Only essential staff should be present in the OR during surgery. Enhanced droplet PPE protection should be worn for all aerosol generating procedures.
3. Airway procedures should be performed in accordance with Anesthesia Patient Safety Foundation (APSF) guidelines.(255)
4. Place a HME/HEPA filter between the Y-piece of the breathing circuit and the patient's mask, endotracheal tube or laryngeal mask airway. The gas sampling line must exit the circuit proximal (closer to the machine) than the filter. The ASA/APSF recommends adding a second HME/HEPA filter on the expiratory limb before entering the anesthesia machine.
5. For sedation cases, a procedural/OR mask should be placed on the patient over the oxygen source. If a gas sampling line is used to monitor end tidal CO₂, ensure a filter is used prior to gases entering the machine. The filter found in most epidural kits may be placed in-line and provide adequate machine protection. For sedation procedures that instrument the esophagus and generate high volume aerosolized secretions, intubation of the airway may be the best way to limit room exposure. Alternately, a Procedural Oxygen Mask may limit room exposure where intubation is contraindicated.
6. For pediatric patients or patients in whom the additional dead space or weight of the filter may be problematic, the HEPA filter can be placed on the expiratory end of the corrugated breathing circuit before expired gas enters the anesthesia machine. Again, ensure the gas sampling line is protected from contaminating the anesthesia machine.
7. Use disposable covers whenever possible (e.g., plastic sheets for surfaces, long ultrasound probe sheath covers) to reduce droplet and contact contamination of equipment and other environmental surfaces.

Postoperative Care.

1. Non-ICU patients should recover in a PACU negative pressure room. If a suitable recovery room isn't available, the OR may substitute until ready for Phase II of recovery from anesthesia.
2. Remove all PPE (except N95 mask) before exiting the OR. Avoid touching hair or face & perform hand hygiene.
3. Surgical scrubs should be changed immediately at the conclusion of each case.
4. Cloth surgical caps should not be worn in PUI cases.
5. The room should be cleaned in accordance with the designated processes for terminally cleaning rooms.
6. Consider air exchange rates for the treatment area and ensure an adequate interval of time between the completion of a procedure and return of staff or initiation of further patient care in that treatment area.
7. When transporting a ventilated patient, ensure a HEPA filter is placed between the ETT and the bag valve mask. Connect the bag valve mask to the ETT prior to opening the door in the negative pressure room. Ensure the door is closed when returning the patient before switching to the ventilator. The same filter may also be used on the exhalation loop of the anesthesia machine.
8. When transporting patients between the OR, a "clean" person who does not contact the patient should accompany the team to safely interact with the environment (e.g. open doors or elevators).

Post-discharge Care.

1. Post-discharge care needs should be assessed and resource availability confirmed.
2. Discharge care plans should consider the risks of exposure from extended healthcare stay (nursing home or other inpatient care facility) and face to face follow-up.
3. Virtual and telehealth should be utilized to accomplish postoperative assessments that do not require face to face interaction.

Special Considerations

Aerosol-Generating Procedures (AGP).

Viral concentration in the aerodigestive tract and respiratory system and aerosol generation during surgical care

Guideline Only/Not a Substitute for Clinical Judgment

Clinical Management of COVID-19, v4

present additional risks to surgical personnel.

1. The performance of high-risk procedures should be limited and risk mitigated through refinement of technique and/or utilization of adjunctive technology and protections. High risk activities include:
 - a. Endotracheal intubation
 - b. Oral surgery
 - c. Tracheostomy and endotracheal tube manipulation/care
 - d. Upper aerodigestive endoscopy (including nasal endoscopy, laryngoscopy, bronchoscopy, and esophago-gastro-duodenoscopy)
 - e. Surgery involving the airway/upper aerodigestive tract, lower airways, or the potential to enter into the upper aerodigestive tract or lower airway.
2. Aerosol generation during surgical procedures can also be limited through the following:
 - a. Electrocautery should be set to the lowest effective setting and a smoke evacuator used if available.
 - b. Chest tubes and surgical drains are all potential sources of aerosolized droplets, and enhanced precautions should be taken during placement, manipulation, or removal.
3. Laparoscopy: Aerosol generation during laparoscopy can be minimized through scrupulous management of access sites, pneumoperitoneum, and through ultrafiltration of aerosolized particles in released CO₂.(256)
 - a. CO₂ insufflation should be set to the lowest effective pressure, and a filtration device should be used for CO₂ release if available.
 - b. Release all pneumoperitoneum via filtration device (if available) or contained suction device prior to specimen removal, port removal, or converting to open surgery.
 - c. Avoid venting insufflation from the ports during surgery.

Endoscopy

Aerosol generation during endoscopy may be difficult to control, therefore performance of these procedures should be carefully considered and engineering controls and PPE optimized to reduce the risk of personnel exposure.

1. In COVID-19 positive patients or PUIs:
 - a. Endoscopy should be performed only for emergent or urgent indications (i.e., cholangitis or gastrointestinal bleeding refractory to medical management).
 - b. Procedures should be performed in negative pressure rooms using PPE as described above in Clinical Management of COVID-19 using endotracheal intubation or a procedural oxygen mask for upper endoscopies, as described above in Intraoperative Care.
 - c. Due to the presence of SARS-CoV-2 RNA in the stool, colonoscopies should be treated as aerosol generating procedures and a procedural/OR mask should be placed on the patient over the oxygen source.
2. After endoscopic procedures in COVID-19 positive patients or PUIs, sufficient time for enough air changes to remove potentially infectious particles should occur before terminal cleaning of the room. The time required for airborne contaminant removal depends on the number of air changes per hour in the room.(257)
3. Endoscopes used in COVID-19 positive patients or PUIs may be reprocessed following standard guidelines for manual cleaning followed by high level disinfection.(258)
4. Consider standard PPE, engineering controls, and room turnover only when the following criteria are met:(258, 259)
 - a. Low incidence or sustained reduction in the number of new COVID-19 cases
 - b. Patients at low risk for COVID-19 (i.e., no concerning symptoms or recent COVID-19 exposure)
 - c. Negative pre-procedure COVID-19 testing (see Surgical Triage and Decision-making, above)

Trauma and Emergency Care. (260)

1. All trauma/injured patients should be presumed positive/PUI in the downrange setting until they can be ruled out (by testing or risk factor assessment). All patients undergoing evaluation and resuscitation for traumatic

Clinical Management of COVID-19, v4

injury require screening and risk-factor assessment to determine the optimal treatment and isolation strategies and potential value of timing of COVID testing.

- a. Trauma team members should all wear appropriate PPE, including airway and eye protection.
 - b. Unnecessary individuals in the trauma bay should be minimized.
 - c. Individuals should remove all PPE (except N95 mask) prior to exiting the resuscitation area.
 - Any clothing worn in the resuscitation bay/ATLS area should be removed after PUI patient contact.
 - Commanders should modify uniform requirements as necessary to allow for multiple rapid clothing changes to avoid cross contamination.
 - d. All equipment in the resuscitation bay and ATLS area (i.e. x-ray, ultrasound, instrument packs, etc.) must be terminally cleaned after every PUI encounter.
 - e. Non-intubated patients should have a surgical mask applied during transport between the resuscitation bay and CT scanner and during any transit within the facility. Patients requiring oxygen should have a non-rebreather mask applied instead of a simple face mask.
 - f. All PUIs either requiring admission or transferred to the emergency department should be kept in isolation rooms (if available) until ruled out or ready for discharge (to quarantine facilities).
2. Staffing risk reduction
 - a. Aerosol generating procedures should have only necessary staff members present in the room (i.e. intubation, chest tube placement, etc.), and all staff must wear enhanced droplet precaution PPE. Intubation should be followed by immediate connection to ventilator to avoid bagging and ETCO₂ monitoring available.
 - b. Each facility should consider options to minimize staff members entering the resuscitation area. This could include the use of runners or pass-through windows for deliveries from pharmacy, lab, etc.
 - c. All visitors should be restricted during the initial phase of resuscitation, and based on risk, may be restricted throughout the entire hospitalization at the discretion of the Commander.
 3. Consultations and therapies should be performed as needed and not delayed solely because a casualty is pending COVID-19 evaluation. This includes specialty and subspecialty consultations, routine nursing care (i.e. pressure injury reduction, oral care, etc.), radiology, lab analyses, and physical/occupational/speech therapy.

Key References:

1. Clinical Issues and Guidance From the American College of Surgeons 2020 [Available: <https://www.facs.org/covid-19/clinical-guidance>].
2. The ASA and APSF Joint Statement on Perioperative Testing for the COVID-19 Virus 2020 [Available: <https://www.asahq.org/about-asa/newsroom/news-releases/2020/04/asa-and-apsf-joint-statement-on-perioperative-testing-for-the-covid-19-virus>].
3. Perioperative Considerations for the 2019 Novel Coronavirus (COVID-19) 2020 [Available: <https://www.apsf.org/news-updates/perioperative-considerations-for-the-2019-novel-coronavirus-covid-19/>].
4. JTS Management of COVID-19 in Austere Operational Environments v2.0 Clinical Practice Guideline 2020 [Available: https://jts.amedd.army.mil/index.cfm/PI_CPGs/cpgs].

OPERATIONAL CONSIDERATIONS FOR COVID-19: PLANNING AND PREPARATION

Providing safe and effective care in the deployed setting during an infectious disease pandemic is particularly challenging given limited resources, close living conditions, and delays in test results and supply arrival. The DOD GCP PI & ID 3551-13 provide a wealth of information, guidelines, and mitigation strategies for a pandemic, but are not tailored to the specific nuances of COVID-19. This section focuses on the unique aspects of dealing with the COVID-19 pandemic in the deployed environment. Collaboration between base commanders and medical teams is an essential component of pandemic response. Additionally, coordination with TRANSCOM is essential as some Geographic Combatant Commanders (GCC) have published orders indicating all COVID positive service members will be transported out of theater while others have maintained treatment in place unless the patient exceeds the capability of the treating facility.

Division of Labor for Quarantine and Isolation.

1. **Quarantine:** This is a *medically-supported command function* to separate high risk individuals from the general

Clinical Management of COVID-19, v4

population following a potential exposure. Commanders are responsible for establishing and maintaining quarantine facilities within their area of responsibility (AOR), and each unit is responsible to identify at-risk personnel based on best medical guidance. Quarantine has also been advocated for units mobilizing to a location where there are no reported cases of COVID-19; these mobilizing units should do a formal quarantine either prior to arrival (then travel only with those who have completed a quarantine) or on arrival to location prior to interacting with anyone on the base.

2. **Isolation:** This is a *command-supported medical function* to care for those with infection. These patients are identified by symptoms (i.e. fever, cough, dyspnea, diarrhea, etc.) following an exposure (typically within the past 14 days), and may be identified de novo or from quarantine. The duration of isolation can vary based upon the duration of symptoms or the results of testing (see below). Because service members are not deployed with a family, even mildly symptomatic patients, who would typically be returned to the care of their family in the garrison setting, become the responsibility of the medical team.

Physical Requirements and Logistics of Quarantine and Isolation.

1. **Quarantine:** Quarters must be provided for persons suspected of having exposure to COVID-19 in an effort to prevent spread of disease to other service members (SM) and civilians on base. These quarters must be separate from the general population and must have their own dedicated toilet and shower facilities. Meals must be provided to quarantined individuals, and they must be checked regularly (i.e. via telephone or in person) to ensure they remain asymptomatic. If symptoms develop, medical personnel should be notified to arrange evaluation and potential transfer into medical isolation. Quarantined individuals should remain in their designated quarters; however, quarantined individuals should be allowed to go outside and exercise in wide open areas to promote mental and physical wellbeing. Personnel should be designated to do laundry for quarantined individuals. Dirty laundry should be placed in a sealed disposable plastic bag by the quarantined member and then handled with gloves by laundry personnel. Laundry should be placed in the washing machine without handling the clothes, and the bag discarded in an appropriate receptacle. Persons in quarantine should remain in quarantine for the allotted 14 days unless it is determined that the person they were suspected of being exposed to is determined not to be infected with COVID-19. The 14 day quarantine resets if any member of the quarantine group develops symptoms or has a positive PCR test result. To avoid excessively prolonged quarantines, every effort should be made to keep quarantined individuals in the smallest possible groups; individual quarters are the ideal quarantine environment. Any personnel interacting with or evaluating quarantined individuals must wear appropriate PPE.
2. **Isolation:** Patients who are symptomatic or test positive should become the primary responsibility of the medical team in isolation. Medical teams will need to plan for patient monitoring, treatment, housing, meal, and hygiene facilities. Based on the demand and the size of the medical treatment team, commanders may need to consider assigning additional non-medical personnel to assist with these tasks. Isolated patients should be classified by symptoms as asymptomatic, mild, moderate, or severe, which will determine the required level of care. Any personnel interacting or evaluating patients in isolation must wear appropriate PPE.
 - a. **Asymptomatic/Mild Symptoms:** In CONUS locations these patients may be sent home for self-care and outpatient follow-up. In the deployed setting family support is absent and self-isolation is not feasible, so medical teams should coordinate with command to establish appropriate isolation housing with routine medical oversight. Symptom progression should result in prompt medical reassessment. There must be a clear and universally-accessible communication plan to notify the medical team of any change in patient condition. This communication plan may need to include providing reliable WiFi to the living area for the isolated patients to use their cell phone or may need to be medical unit supplied radios or phones.
 - b. **Moderate Symptoms:** These patients require hospital ward admission. These facilities may be located within the MTF or established separately near the MTF; if available, negative pressure facilities should be reserved for aerosol producing procedures. A COVID-19 positive patient should not share a room with a non-COVID-19 patient.
 - c. **Severe Symptoms:** These patients require ICU admission for hemodynamic monitoring/treatment and management of severe respiratory symptoms. ICU care should be performed where the greatest medical capability exists, but these patient should not be placed in the same facility used for other non-COVID-19

Clinical Management of COVID-19, v4

patients (such as trauma patients). Negative pressure facilities should be used (if available) during aerosol generating procedures. If negative pressure facilities are not available then a well-ventilated tent or building can be used if it has an air handling separate from all other inpatient and clinic areas. Oxygen generating capability will need to be established along with continuous patient monitoring and nursing care. This level of care can be resource intensive and medical teams will need to work with TRANSCOM on patient transfer if they do not have adequate resources.

- d. **Discharge:** Patient placed in isolation should be classified as patients under investigation (PUI) while awaiting their test results. They will need to remain in isolation for 10 days and be at least 72 hours without fever if positive for COVID-19. Alternatively, they can be released from isolation if they have two negative COVID-19 tests after their positive test. If their initial COVID-19 test is negative they should remain in isolation for at least 10 days unless an alternative diagnosis can be rendered for their symptoms (i.e. community acquired pneumonia, lymphoma, rhabdomyolysis).

Unique Limitations in the Austere Environment

1. **Ventilator:** COVID-19 may result in ARDS which can be challenging to manage even with the best facilities and equipment; in the deployed setting, providers may only have transport ventilators with limited capabilities. It is important to recognize this limitation and prepare accordingly. Medical teams should prepare protocols for prone positioning and pharmacologic paralysis, which have a mortality benefit if used early in moderate to severe ARDS.
2. **Medications:** At present, there is no specific treatment for COVID-19. Deployed providers may not have access to compassionate use or trial medications, and should be familiar with the supportive care measures described elsewhere in this document. Additionally, the Society of Critical Care Medicine, ARDSNet, and other professional societies provide continuously updated guidelines on their websites. Providers should work closely with pharmacy and logistics leadership to ensure adequate stocks of all commonly required medications, including antimicrobials, sedation, and paralytics.
3. **PPE:** Supply chain challenges have led to PPE shortages worldwide. Fortunately most units are deployed with CBRNE equipment which can be used for staff protection. Staff must be proficient at proper donning, doffing, and cleaning techniques.
4. **Hygiene:** The austere environment lends itself to rapid spread of infectious disease. Command staff should emphasize the importance of handwashing/sanitizing, cleaning quarters, and appropriate social distancing.
5. **Testing:** Epidemiologic data is critical for command decision making, therefore commanders may need to divert resources to ensure rapid case identification and intervention. MTFs should not use non-DOD laboratories for testing unless approved by their AOR HQ.
6. **Transportation:** Units must coordinate with PMC to ensure safe and efficient movement of patients and/or testing samples around theater. Patients can be treated in place unless their clinical condition necessitates a higher level of care or if command directs movement of COVID-19 positive patients. Unnecessary patient movement should be avoided to minimize personnel and resource exposure and transmission risk. TRANSCOM has developed 3 categories for patients based on their oxygen requirement to facilitate planning for safe medical evacuation of patients.
 - a. Category A: intubated or O₂ sats <85% on RA and <92% on 5 L/m O₂
 - b. Category B: O₂ sats <90% on RA or sats >92% on 4 L/m O₂
 - c. Category C: O₂ sats >92% on RA
7. **Housekeeping and Cleaning Services:** Cleaning protocols must be established to ensure adequate sanitization occurs in quarantine, isolation, and medical facilities, as well as workspaces and quarters of those moved to quarantine/isolation status. PPE should be worn by cleaning personnel and disposed of in a manner that avoids the potential for cross-contamination.
8. **Mortuary Affairs and Casualty Liaison Teams:** While the COVID-19 mortality rate is expected to be low in the deployed military population, Mortuary Affairs teams should be prepared for increased demands and requirements. Casualty Liaison teams should be ready to work with commanders, medical teams and families on accurately reporting patient status.

BEHAVIORAL HEALTH AND WELLNESS IN COVID-19 CLINICAL MANAGEMENT

Delirium

1. Delirium is anticipated in up to 82% of ICU intubated patients, and the expectation is ICU patients with COVID-19 will experience delirium at a similar rate.(261) Traditional methods of diagnosis, non-pharmacologic management are difficult given isolation precautions.(262) In addition to the CAM-ICU, the Stanford Proxy Test for Delirium (S-PTD) is a validated tool that relies on nursing report of their interactions with patients over their full shift to confirm diagnosis of delirium and would be useful in this setting.(261, 263) No studies exist validating pharmacological management of delirium in COVID-19 positive patients but the following may be considered:
 - a. Melatonin 10-15mg enteric at night for anti-inflammatory effects and regulation of sleep-wake cycle
 - b. Suvorexant 5-20mg enteric at night for sleep-wake cycle regulation
 - c. Alpha-2 agonists to down titrate cytokine and moderate adrenergic storm
 - Dexmedetomidine IV 0.1-2.4 mcg/kg/hr to manage acute agitation and cycling
 - Guanfacine 0.5mg BID – 1mg TID enteric to taper off sedative drips
 - d. Antipsychotics to downregulate excess dopamine inherent to delirium (Haloperidol IV 0.5mg-30mg per 24 hours) **must monitor QTc prolongation*
 - e. Valproic Acid in hyperactive and/or mixed delirium due to potential anti-inflammatory and anti-oxidant effects, and might decrease transcription of interleukin-6 (enteric or IV, 250-500mg BID and titrate to 500mg qAM, 500mg q afternoon, and 1000mg qHS) **monitor LFTs, platelets, ammonia levels* (261)
2. Consider avoiding/ minimizing use of benzodiazepines, opioids, medications with strong anticholinergic properties as they can be deliriogenic, though there are clinical circumstances where these are appropriate.

Psychopharmacology

1. COVID-19 can invade the CNS and may be causing development of psychiatric illness. Full review of all medications as well as substance use history is necessary to delineate primary psychiatric symptoms versus medication side effects or symptoms directly attributable to COVID-19 infection. Case reports of new onset psychosis, presenting with severe anxiety, agitation, paranoia, disorganized thinking and auditory hallucinations, was associated with otherwise asymptomatic COVID-19 positive patients. These cases were responsive to typical treatment with second generation antipsychotics.(264)
2. Due to the multi-organ system effects of COVID-19, consideration for use and need for monitoring of psychotropics must be tailored to the patient's specific situation. This list is not meant to be wholly inclusive – but use caution when the following symptoms are of clinical concern:
 - a. Leukopenia, neutropenia, agranulocytosis: Carbamazepine, clozapine, and all first and second generation antipsychotics **Clinicians who have patients on clozapine should consider cutting the dose by half if the patient develops fever and/or other signs of infection*
 - b. Platelet dysfunction and increased bleeding risk: Medications that inhibit serotonin reuptake (SSRIs, SNRIs, TCAs) and valproic acid
 - c. QTc prolongation and concern for exacerbation with some COVID-19 treatment options: some antipsychotics, tricyclic antidepressants, citalopram
 - d. Drug-induced liver injury: chlorpromazine, carbamazepine, valproate, duloxetine, and nefazodone
 - e. Impaired renal excretion: Lithium, gabapentin, topiramate, pregabalin, paliperidone, duloxetine
 - f. Lowered seizure threshold: most antipsychotics, bupropion, tricyclic antidepressants (265)
3. There are multiple neuropsychiatric side effects associated with current medications used for treatment of COVID-19, to include psychosis, depression, sleep disruption, and anxiety/agitation. It is recommended these symptoms be treated as clinically appropriate, with cautious monitoring if there are concerns for additional complications (i.e., benzodiazepine use for severe anxiety symptoms). Specifics of medications:
 - a. Remdesivir: None noted
 - b. Chloroquine and Hydroxychloroquine: psychosis, delirium, agitation, suicidality, personality changes, depression, sleep disturbance
 - c. Tocilizumab: Exacerbation of depression, anxiety, pain, and sleep disruption

Clinical Management of COVID-19, v4

- d. Favipiravir: No published information
- e. Lopinavir/Ritonavir: Abnormal dreams, agitation, anxiety, confusion and emotional lability **Ritonavir has been shown to lower concentration of some psychotropics due to presumed CYP induction (bupropion, methadone, lamotrigine, olanzapine)*
- f. Azithromycin: Psychotic depression, catatonia, delirium, aggression, anxiety, dizziness, headache, vertigo, somnolence
- g. Corticosteroids: depression, mania, agitation, mood lability, anxiety, insomnia, catatonia, depersonalization, delirium, psychosis
- h. Interferon-Alpha: boxed warning for “life threatening or fatal neuropsychiatric disorders” – fatigue, mood disorders, suicidality, anxiety disorders, irritability, lability, apathy, sleep disturbance, cognitive deficits (265)

For Defense Health Agency COVID-19 Related Behavioral Health (BH) Resources:

<https://info.health.mil/army/bhsl/Covid19/Forms/AllItems.aspx> (DoD CAC Enabled only)

Pandemic conditions require medical staff to be sensitive and responsive to patient, family, provider, and leader needs. Common pandemic responses include a predictable range of distress reactions (e.g. insomnia, fear, grief), health risk behaviors (e.g. increased use of alcohol/other substances, work/life imbalance), and may also result in BH disorders (e.g. PTSD, depression, and anxiety). In response to multiple stressors, associated with quarantine or in support of critical care operations, common responses may also include anger, irritability, detachment, avoidance, impaired function, and burnout. Addressing stress responses early can mitigate enduring impacts.

General Considerations for Frontline Workers, First Responders, and Support Staff

1. Prioritize basic needs. Proper sleep, nutrition and hydration, regular exercise, regular breaks, and appreciation/gratitude can sustain performance and enhance decision-making.
2. Social distancing, infection control, and isolation present a significant barrier to our usual approach to care, requiring innovative approaches.
3. Communication – words matter now more than ever. Clear and consistent messaging from leadership, between team members, and to patients and family is vital during this crisis.
4. Anticipate fears of returning to work and provide thoughtful, transparent information.
5. Resources for leaders in support of Healthcare Workers can be found at: <https://www.cstsonline.org/covid-19/supporting-healthcare-workers>

General BH Care for Patients with known or suspected COVID-19

1. In accordance with HPCON, use telehealth and virtualization tools as much as possible for BH assessments and ongoing care of isolated patients. Promptly identify all COVID-19 patients with known mental illness and consult BH to assist with ongoing care.
2. Recognize isolation as a barrier to communication. Keeping patients informed as to what is happening, what is likely to happen, and next steps in their care may provide a sense of control in the midst of a stressful and confusing situation. Expand virtual approaches to care and provide regular updates to patients and families.
3. Anticipate patient concerns and misconceptions. Concerns that may be present include fears related to transmission to family members, fears related to intensive care or ventilator availability, duration and impact of isolation, or external stressors such as impact on job, housing, and finances.
4. Healthcare systems should establish easily accessible pathways for BH referrals for family members of patients admitted for COVID-19.
5. Attend to negative impacts of isolation by facilitating virtual connection with providers, family, and loved ones as much as possible. This could include providing patients with dedicated mobile devices/tablets.
6. Resources to help in caring for Patients and Families can be found at: <https://www.cstsonline.org/covid-19/caring-for-patients-and-families>

For Medical Staff

1. Self-care is important for providers, patients, and families.
2. Connect to a sense of unified purpose; foster hope, fortitude, and tolerance in self and others.

Clinical Management of COVID-19, v4

3. Amplify positive stories and stories about competent efforts by self and colleagues. Encourage perceptions of competence among staff, especially junior and/or less experienced colleagues.
4. Recognize and attend to signs of stress reactions or burnout in self and others (e.g. out of character sadness, frustration, irritability, isolation/disconnectedness, substance use, and lack of self-care). Usually these can be addressed with simple measures, including normalization, peer-support, and rest with expectation of rapid return to full functioning.
5. Focus on what can be controlled – checklists, routines, self-care; and accept what cannot be controlled.
6. Promote a climate where it is acceptable for team members to talk about difficult events (e.g., death, triage, errors), as avoidance and fear of such thoughts are associated with greater long-term mental health problems.
7. Establish a routine of regular team meetings as an opportunity to pass relevant information, but also as an opportunity to check in with each other and rotate duties as needed. Maintain a climate where it is okay to not be okay and offer peer support when needed.
8. Resources for Healthcare Worker Self Care can be found at: <https://www.cstsonline.org/covid-19/healthcare-worker-self-care>

For BH Providers

1. Provide proactive support to frontline workers where possible, and at times of peak stress, ideally, in the form of BH outreach teams with established relationships to frontline and medical staff points of contact. Consider BH team outreach routinely (e.g. during daily rounds, at shift changes).
2. Be careful not to overlook other at risk groups such as janitorial staff, transport, food service, and others who make the medical system run, and may also be at risk of exposure and are likely to experience distress.
3. Behavioral health care teams can provide both non-clinical support to frontline staff as well as be available to facilitate referral for additional BH care when needed.
4. Tailor resources and support as much as is feasible – and plan on changing/adapting resources with the unfolding realities of the medical mission. Flexibility is important.
5. Supportive care of healthcare workers is different from usual clinical care, and includes:
 - a. Check in with the physicians, nurses, technicians, and support staff, and get to know their mission and challenges in a non-intrusive manner.
 - b. Link with support services, such as Red Cross, providing food and beverages.
 - c. Provide information on normal stress reactions and adaptive responses.
 - d. Promote positive peer support and facilitate connections.
 - e. Make connections during a calm time. Do not interrupt urgent patient care or sign-out.
 - f. Offer combinations of simple supportive non-clinical strategies, as well as clinical triage when appropriate (e.g. find a quiet space to talk when things are chaotic).
 - g. Ensure individuals have access to safe spaces and emotional/spiritual support.
6. Unique issues to consider when supporting frontline workers:
 - a. Be aware of the potential for distress related to ethical issues in providers making difficult and potentially life or death triage and management decisions.
 - b. Be aware of potential concerns of individual front line workers, including single parents, dual healthcare worker families, families with serious medical issues, workers living separate from their families, and individuals facing the community stigma of being “infected.”
7. Resources for Patients can be found at: <https://www.cstsonline.org/covid-19/mental-health-support>

For additional COVID-19 Related Behavioral Health (BH) Resources:

<https://www.cstsonline.org/>

<https://asprtracie.hhs.gov/COVID-19>

REHABILITATION CONCERNS FOR PERSONS WITH COVID-19

Rehabilitation of COVID-19 Patients and PUIs

Overview.

Clinical Management of COVID-19, v4

1. This document is intended to provide guidance and planning considerations for the provision of acute and critical care rehabilitation for hospitalized patients by practicing acute care Physical Therapy (PT) and Occupational Therapy (OT) providers and augmented staff dedicated to support the COVID-19 response.
2. The goal of acute care inpatient rehabilitation is to improve mobility in order to reduce mortality, decrease hospital length of stay (LOS), decrease ICU and ventilator days, streamline patient throughput, and decrease the burden of acute rehabilitation after discharge.
3. Early rehabilitation involvement in the facility's COVID-19 planning team is recommended to anticipate rehabilitation needs.
4. Rehabilitation personnel should be dedicated to either COVID-19 patients or non-COVID-19 patients to minimize potential exposure.
5. Pool staff resources as able and maximize distancing. Maintaining appropriate work/rest cycles.
6. Screening tools should be used to quantitatively determine a patient's need for therapy intervention.

ICU and Critical Care Staffing ratio recommendations when respiratory rehabilitations is a primary intervention

1. ICU recommendations: 4 therapy providers for the first 22 ICU beds. One FTE for each 4-bed increase.
2. Acute Care Recommendations: therapy provider FTE for the first 11 beds and a potential increase of 2 per additional 11 beds.
3. Subacute and Acute Inpatient Rehabilitation Unit (if present): 2.5 FTEs per 11 beds.
4. Staffing ratios may be lower when rehabilitation interventions are the primary focus rather than on respiratory rehabilitation

Personal Protective Equipment

1. Prior to working with patients with COVID-19, therapy staff should have comprehensive training on the use of PPE.

Treatment Guidelines

1. Positioning: Rehabilitation staff may be involved in prone positioning with COVID-19 patients due to their expertise in safely and optimally performing this task.-
2. Rehabilitation should progress to active movement as soon as possible.
3. Provide patient specific exercise programs and handouts.
4. Partner with nursing for patient active participation in care and exercise.
5. Interactions with COVID-19 patients will be limited to a contained environment where airborne precautions can be maintained.
6. Therapy staff must have advanced understanding of medical implications of COVID-19
7. Emphasize treatment of non-COVID-19 patients for quick discharge to decrease risk of exposure.
8. Attend to the well-being of the whole patient by promoting orientation and communication with patient during therapy sessions-

Discharge Planning

1. Goal should be safe patient discharge to home from the acute hospital setting whenever possible.
2. Therapists should participate in multidisciplinary rounding/discharge planning to ensure necessary patient supports are in place for discharge.
3. Electronic communication with spouses and other care providers should be completed to promote patient and family confidence in the discharge plan.
4. If discharged directly to home, recommend telehealth follow-up with the patient within 3-5 days post-discharge.

More information on Rehabilitation Guidelines can be found in the DHA Patient and Caregiver COVID-19 Rehabilitation Guide as well as the following sites: Allied Health Folder in <https://www.milsuite.mil/book/groups/covid-19-clinical-operations-group> or <https://www.milsuite.mil/book/groups/acute-and-critical-care-physical-therapy>

TELEMEDICINE SUPPORT DURING THE COVID-19 PANDEMIC

1. Telemedicine, also referred to as virtual health (VH), encompasses a set of tools that leverage information and communication technologies to most commonly extend medical care across geographic distances and boundaries. These same tools have a significant and unique potential to support care delivery during an

Clinical Management of COVID-19, v4

infectious pandemic in order to decrease healthcare worker exposure to contagion (i.e. “clinical distancing”), reduce the usage of consumable PPE, while also enabling continued medical care delivery for non-infected patients while in their home. Accordingly, the CDC now recommends the liberal use of telehealth during the COVID19 Pandemic (<https://www.cdc.gov/coronavirus/2019-ncov/healthcare-facilities/guidance-hcf.html>).

2. Telemedicine can be delivered through two primary manners
 - a. Direct-to-patient VH. Services delivered in this manner require credentialing and privileging IAW DHA PM 6025.13 using the centralized privileging by proxy for telemedicine (TPbP) through the Virtual Medical Center. A provider or a patient can be in the home for a telemedicine visit. Direct-to-Patient VH is most appropriate when a provider is directly evaluating a patient, and requires documentation of the encounter in the electronic health record (EHR).
 - b. Tele-Consultation. Services delivered in this manner may occur without separate privileging at the patient’s location, and typically are performed from healthcare professional to healthcare professional (i.e. trained clinician to trained clinician like medic to remote physician or nurse to physician or physician to physician).
3. Telemedicine technology:
 - a. Phone calls can be used for a majority of patient encounters during the COVID Pandemic. The need for clinical video versus telephone and/or secure messaging will be based upon the provider’s individual judgement, and will take into consideration the specific patient complaint evaluated.
 - b. Clinicians engaging in telemedicine (especially forums that utilize video with the patient) must appreciate the burden it places upon valuable network resources. The solution that achieves clinical needs and uses the *minimal* network resources should be utilized when possible.
4. All care provided through telemedicine should be documented in the appropriate EHR. If the provider is delivering care from outside of the MTF, the DHA Application Virtualization Hosting Environment (AVHE) can be utilized to access the EHR.
 - a. AVHE can be accessed from a computer with a CAC-card reader through the following URL: <https://avhe.health.mil>.
 - b. Make sure to select your email certificate
5. There are several use-cases for telemedicine during the COVID-19 Pandemic. Each require planning and practice to be successful.
6. Use cases for which currently available MHS approved solutions exist include:
 - a. Screening and Initial Evaluation (e.g. Virtual Clinics)
 - i. Phone calls can be used for a majority of patient encounters during the CoVID Pandemic. The need for clinical video versus telephone and/or secure messaging will be based upon the provider’s individual judgement, and will take into consideration the specific patient complaint that is being evaluated.
 - ii. Web-portal based screening tools suggest need for patients to engage with their healthcare system (reduces overall burden on the system if patients are screened as low risk). Some examples of online tools are listed below, although none are created or owned by the DOD:
 - a) <https://c19check.com/start>. Site hosted by Emory University Medical Center, which provides likelihood of CoVID infection based on answering series of online questions.
 - b) <https://penn-chime.phl.io/> Site hosted by Penn State Medical Center, Predictive Healthcare Team, which provides patient volume projections during the pandemic.
 - iii. Asynchronous solutions including web-portal based messaging (e.g. Federal Secure Messaging and MHS GENESIS patient portal) and e-mail allow engagement with the healthcare system with minimal network resource use.
 - iv. Where available, portable telemedicine units can be employed by triage and Emergency Department personnel to evaluate patients to reduce clinician exposure to potentially sick patients; Telehealth in a Bag (THIAB), Transportable Exam Station (TES), and Video Teleconferencing (VTC) Carts with/without virtual exam equipment.

Clinical Management of COVID-19, v4

- v. These systems can connect a patient (within an isolation setting) to a provider (within a “clean” setting) by use of either portable data networks (PDN’s), WiFi routers, cellular service, or hospital WiFi networks.
- vi. Synchronous video to the patient’s location can be accomplished through several mechanisms. The preferred and supported solutions are Adobe Connect and Cisco Meeting Server (more below).
- b. Inpatient Wards (non-ICU)
 - i. Where available, portable telemedicine units can be employed by triage and Emergency Department personnel to evaluate patients; Telehealth in a Bag (THIAB), Transportable Exam Station (TES), and Video Teleconferencing (VTC) Carts.
 - ii. These systems can connect a patient (within an isolation setting) to a provider (within a “clean” setting) by use of either portable data networks (PDN’s), WiFi routers, cellular service, or hospital WiFi networks.
- c. Tele-Critical Care
 - i. Sites that are currently enrolled in the Joint Tele-Critical Care Network, should use this existing resource to support care of critically ill patients with or without suspected / confirmed COVID-19.
 - ii. Sites that are not currently enrolled in JTCCN, should attempt triage and management of patients as outlined in this document and per usual standards of care. For hospitals that typically do not care for critically ill patients, this may involve transfer of the patient to a local civilian hospital.
 - iii. MTFs that are not enrolled in the JTCCN that (1) do not have sufficient critical care expertise, and (2) cannot transfer critically ill patients, may be forced to care for these patients. In this situation, tele-consultation is available to support clinicians.
- d. Tele-consultation (outside of JTCCN):
 - i. Advanced Virtual Support for Operational Forces (ADVISOR) Program. 1-833-ADVSRLN (238-7756) or DSN 312-429-9089
 - a) The ADVISOR program was originally designed for operational VH support.
 - b) Due to COVID-19 garrison support has been expanded to include:
 - Critical Care (Non-JTCCN MTFs)
 - Infectious Disease
 - Pediatric Infectious Disease
 - Future expansion: Behavioral Health and Palliative Care
 - c) Phone calls will be routed by live ADVISOR Care coordinator(s) 24/7/365.
 - d) The caller needs to identify that they are requesting support for critically ill patients located in a MTF.
 - e) The care coordinator routes the call to a geographically located MTF with the available specialty.
 - f) ADVISOR is only available for MHS providers.
 - g) Information on the program can be found at:
<https://info.health.mil/army/VMC/Pages/Home.aspx>
 - h) Additional questions or information on ADVISOR can be obtained by emailing dod.advisor_office@mail.mil or scanning the QR code (shown to the right):






Figure 11. Telecritical Care Pandemic/Natural Disaster Decision Pathway

- e. Virtual Health to Patient Location (e.g. home)
 - i. The CDC recommends providing outpatient care where/when possible through telemedicine to minimize infectious exposure in MTFs for at risk patients and staff.
 - ii. Virtual health to patient location can be established through several mechanisms.
 - a) Secure Messaging (e.g. Federal Secure Messaging, MHS GENESIS Patient Portal).
 - b) Establishing a clinic cell phone with texting services and publishing the number
 - c) Using phone calls to discuss patient problems/symptoms as indicated.
 - d) Conducting Synchronous Video Visits can be performed through either Adobe Connect or Cisco Meeting Server (preferred solutions), or through several non-public facing communication platforms.
 - Adobe Connect accounts can be requested from the VMC Front Office at: <https://info.health.mil/army/VMC/Pages/Home.aspx>
 - Online VH training should be completed prior to Adobe Connect account creation, but there are exceptions during the pandemic to get accounts deployed rapidly. The DHA Virtual Health Provider Training (US444) can be found on the JKO training website: <https://jkodirect.jten.mil>.
 - Additional guidance will be forthcoming IRT the Cisco Meeting Server capability. The capability being established by DHA J6 will have several interconnected servers spread across the enterprise.
 - The following non-public facing communications tools are authorized for provider-patient medical interactions, however these technologies are not supported by the DHA or DOD.
 - Apple FaceTime
 - Google Duo
 - Microsoft Skype
 - f. OCONUS MTFs may utilize existing asynchronous virtual health platforms (PATH for INDOPACOM, HELP for EUCOM, AFRICOM, and CENTCOM) to obtain teleconsultation subspecialty consultation.
7. Documentation, Billing, and Coding (See **Appendix O**)
- a. When direct-to-patient telemedicine is performed, encounters should be documented in the appropriate electronic medical record (AHTLA or Genesis for outpatients, Essentris or Genesis for inpatients).
 - b. If the Military Medical Treatment Facility (MTF) is open and conducting normal clinical operations, no change in coding is necessary.

Clinical Management of COVID-19, v4

- c. Up to date virtual health coding references can be found at:
<https://info.health.mil/army/VMC/Pages/COVID-19/CodingGuidance.aspx>
 - d. If the MTF is open, but is restricting access for patients who can be treated virtually, the processes are as follows:
 - i. By telephone only:
 - a) Document as normal for the appropriate encounter type (not in t-con module) to include history, any counseling, assessment and plan, and disposition. Include time spent during the encounter, if required, by service performed.
 - b) Assign the diagnoses, as appropriate.
 - c) Assign G2012 in the procedure (Healthcare Common Procedure Coding System [HCPCS]) code section.
 - d) Assign E/M 99499 or leave blank.
 - ii. By synchronous visual and audio telecommunications:
 - a) Document as normal for the appropriate encounter type to include history, exam if done, any counseling, assessment and plan, and disposition. Include time spent during encounter if required by service performed.
 - b) Assign the diagnoses, as appropriate.
 - c) Assign any procedures performed and documented (e.g., psychotherapy, PHQ-9, etc.)
 - d) Assign appropriate Evaluation and Management (E/M) service, if performed; otherwise assign 99499 or leave blank.
 - e) Apply virtual encounter modifier to encounter (GT=MTF to MTF or 95=provider to patient location other than an MTF).
8. Other Considerations:
- a. Always be conscious of the need to maintain patient privacy and data security and clearly delineate risks to the patient or healthcare professionals using the system.
 - b. Do NOT use photos, video, geospatial positions when you are in an operationally sensitive area: ALWAYS CONSIDER OPSEC!
 - c. Before pursuing a new application of telehealth, CLEARLY DEFINE YOUR USE CASE, then consider technology resources (hardware, software, and network combinations) that can be used for your use case. Most importantly, consider HOW you will use the technology and practice this workflow before implementing it broadly at your location. Consider the following:
 - i. Who will use your solution?
 - ii. Why would they use your solution?
 - iii. When would they use this solution?
 - iv. Where will they use the solution (in a patient room, at a nursing station, from a home/office, to a home/office, etc.)?
 - v. What combination of hardware, software, and network will be used?
 - vi. How will they use it (training, how-to guides, etc.)
 1. How will they document care?
 2. How will you maintain patient regulation (admission/discharge/transfer)?
 3. How will you maintain team-based care as necessary?
 - d. PRACTICE your solution on a small scale before deploying more broadly.
 - e. Establish routine communication with leadership regarding current capabilities and your telehealth solution's potential to off-load aspects of bedside care to telemedicine support. Use telemedicine to triage bedside clinician time and activities. Necessary to do this is good communication and trust between the bedside clinical team and the remote clinical team. One way to facilitate this is to rotate teams from bedside duties to telemedicine duties or to shift infected caregivers toward telemedicine and recovered caregivers towards the bedside. Importantly, asking/having all clinicians participate in telemedicine increases their awareness and understanding of telemedicine capabilities and limitations.
9. Questions regarding MTF and Market telemedicine capabilities should be directed to MTF and Market

Clinical Management of COVID-19, v4

virtual health leads. Questions that cannot be answered by the MTF/Market VH lead, or questions pertains to an enterprise VH service, should be directed to the regional VMC hub site.

- a. CONUS: VMC-C located in San Antonio (1-844-VMEDCEN)
- b. INDOPACOM: VMC-IP located in San Diego, CA
- c. EUROPE: VMC-E located in Landstuhl, Germany

EMERGENCY MEDICAL SERVICES (EMS) AND GROUND TRANSPORT OF PERSONS WITH COVID-19

911 Public Safety Answering Points (PSAPs) and Dispatch Screening for COVID-19

1. Persons assigned to EMS and first responder dispatch function should complete key question interrogation and dispatch resources accordingly. Dispatchers should reference the EMS COVID-19 questionnaire when obtaining information from 911 callers (Table 9). EMS systems may become strained due to an influx of 911 calls regarding known or suspected COVID-19 transmission or infection. In areas where EMS resources are overwhelmed by 911 call volumes, the following should be considered:
 - a. EMS and/or Fire Dispatch should triage 911 calls and prioritize responses accordingly (e.g. if a patient calls reporting signs and symptoms consistent with COVID-19, but denies respiratory distress and other complaints suggestive of a life-threatening condition (i.e. chest pain, etc.), ambulance services should be prioritized to an alternative, higher-acuity call.
 - b. If EMS arrives on scene and determines that a patient does not have a life-threatening condition relating to the potential exposure to, or signs and symptoms of, COVID-19, EMS crews should contact On-line Medical Control to discuss non-transport and/or alternative transport destinations. If non-transport is approved, EMS Dispatch should direct the EMS crew to a higher-acuity 911 call. Refusal of Transport /Treat and Release should be coordinated with local On-line Medical Control
 - c. Callers using the 911 system for questions or concerns regarding COVID-19 testing (e.g. sites, locations, and decisions regarding testing criteria) should be diverted to established local, county, or state COVID-19 call centers. Installations and facilities should consult with their local EMS Medical Directors regarding protocols and policies pertaining to call diversion for information-only requests from 911 callers.

Additional guidance for PSAPs can be found on the CDC site at: <https://www.cdc.gov/coronavirus/2019-ncov/hcp/guidance-for-ems.html>

Pre-Arrival Screening or Initial Patient Assessment of Suspected COVID-19 Patients. *(For utilization by EMS/Fire Department Dispatch OR Responding Crews)*

1. Perform initial assessment from at least six feet away if patient presentation allows. If the patient reports symptoms consistent with a respiratory illness, EMS personnel should don appropriate PPE. With potential widespread COVID-19, all patients should wear a surgical-type mask (best) [or alternatives as available, e.g., cloth (better)].
2. If the patient’s condition allows, to minimize the risk of exposure, one individual should approach the patient, place a surgical-type mask on him/her, and complete the COVID-19 screening questionnaire/ initial assessment. Additional EMS/Fire personnel should be contacted for support only as required.
3. If EMS personnel are first on-scene, and it is determined that the patient has symptoms of a respiratory illness (Box 1) and risk factors for COVID-19 (Box 2), Dispatch should be contacted to minimize response by additional units (Fire and Law Enforcement) to reduce the risk of exposure unless those resources are required for other aspects of the call.

Table 9. Emergency Medical System/First Responder Pre-Arrival Screening for COVID-19

Does the patient have:		
BOX 1	AND	BOX 2
<ul style="list-style-type: none"> • Fever (or are they hot to the touch) • Cough • Shortness of Breathing or Difficulty Breathing • Other flu-like symptoms (sore throat, runny nose, 		<ul style="list-style-type: none"> • Are they currently under investigation or isolation for COVID-19 by public health or other medical professionals? • Have they been in close contact with an individual who is known to be sick with, or under public health/medical

body aches, chills, nausea, vomiting, diarrhea)		professional investigation/isolation for COVID-19?
---	--	--

If the patient meets at least one criteria item from Box 1 and Box 2, see below:
<ul style="list-style-type: none"> • Instruct the individual to isolate him/herself from close contact with others until EMS arrives. • Notify First Responders (to include Fire and Law Enforcement) that the patient meets pre- arrival screening criteria for COVID-19. Advise donning of appropriate PPE prior to patient contact. • Follow local agency policies to limit multi-unit responses. • Transport Agencies will contact the receiving facility as soon as possible, preferably prior to transport (See EMS TRANSPORT OF PERSONS UNDER INVESTIGATION OR PATIENTS WITH CONFIRMED COVID-19).

Table adapted from the Southwest Texas Regional Advisory Council (STRAC); EMS Pre-Arrival Screening for Coronavirus 2019-nCoV - V1.2, issued 02/28/2020.

EMS Non-Transport/Treat on Scene

1. Purpose: Identify patients that do not require EMS transport to a hospital or alternate facility during the COVID-19 pandemic, in order to accomplish the following: 1) Minimize disease transmission to the community and health care system; 2) Protect first responders and health care providers and; 3) Preserve the health care system functionality by not overwhelming emergency resources.
2. Transport decision and final destination versus non-transport with self-care should be considered by EMS Medical Directors, partnering with MTF leadership, to develop local policies. The following are provided as recommendations:
 - a. Careful consideration for EMS Non-Transport should be given for pediatric patients, pregnant females, or patients who are immunocompromised. Discussion with Online Medical Control is advised.
 - b. The below assessment tool is to inform the necessity to transport an adult patient when the patient reports symptoms related to COVID-19.
 - c. If a patient is not transported, he/she should be directed to contact 911 if he/she develops significant shortness of breath, or chest pain. Recommendations for non-emergent care follow up per local resources should be provided. First use local nurse advice line or primary care telemedicine if there is the inability to tolerate oral intake even at very small amounts of 5-10 mL (1-2 tsp every 5 min). Inability to schedule follow-up with an appropriate health care provider/facility is not a 911 call unless emergent symptoms above are present instead a non-emergent resource line should be provided.
 - d. The patient must be in agreement with non-transport and the time taken to explain other resources that are more appropriate to get patients buy-in and understanding.

Table 10. Emergency Management System Patient Considerations for Non-Transport in COVID-19

<u>PATIENT CONSIDERATION FOR NON-TRANSPORT:</u>
INITIAL ASSESSMENT WITH VITAL SIGNS
<i>(initial encounter should ideally be by a single provider in appropriate PPE from a distance of 6 feet)</i>
<ul style="list-style-type: none"> • Temp < 39.4°C (103°F) • GCS 15, Alert & Oriented • Respiratory Rate 10-30 • SpO2 > 90% (with basic ADLs) • HR < 100 bpm • Well appearing, speaks in full sentences, ambulatory • Viral sx: cough, sore throat, body aches, nasal/chest congestion
PATIENT MEDICAL HISTORY & PRESENTATION
<ul style="list-style-type: none"> • Age < 50 years • Non-diabetic • Non-Immunocompromised • No known respiratory disease • No known cardiac disease
LIVING ARRANGEMENTS
<ul style="list-style-type: none"> • Has appropriate support system at home • Patient has means for follow-up
IF THE PATIENT IS IN A PUBLIC LOCATION
<ul style="list-style-type: none"> • Place a surgical mask on the patient. • Discourage the use of public transportation. • Instruct the patient to directly transport themselves home while minimizing exposure to others/the community.

Pre-hospital personal should continue to reference current CDC guidance regarding PPE and Transport of PUIs or Patients with Confirmed COVID-19: <https://www.cdc.gov/coronavirus/2019-ncov/hcp/guidance-for-ems.html>

EMS Transport in Resource-Limited Environments.

Clinical Management of COVID-19, v4

1. During the pandemic, MTFs and civilian EMS services may become inundated with critically ill patients, exceeding MTF treatment and transport capabilities. It is strongly recommended that EMS Medical Directors partner with MTF leadership to discuss disaster response contingency plans relating to inter-facility transports. Nationally Registered Paramedics (NRPs), with approval and guidance from local EMS Medical Directors, are authorized to transport critically ill patients via ambulance. The following are ambulance staffing recommendations to be utilized according to staffing capabilities and patient acuity:

GOOD	
<i>If the patient:</i>	<i>Crew (in addition to the EMT/NRP driver):</i>
Is not ventilated and has no more than two intravenous (IV) or intraosseous (IO) pump infused medications	Paramedic
Is not ventilated and has ≥3 IV/IO pump infused meds	Paramedic AND Critical Care Registered Nurse (CCRN) OR Certified Emergency Nurse (CEN)
Is ventilated and has ≤2 IV/IO pump infused meds	Paramedic x 2 OR Paramedic AND Respiratory Therapist (RT)
Is ventilated and has ≥3 IV/IO pump infused meds	Paramedic x 2 AND CCRN OR CEN OR Paramedic, RT, AND CCRN OR CEN
Is ventilated and has three or more IV/IO pump infused medications	If NRPs are unavailable, consider utilizing MTF CCAT Teams OR hybrid transport teams consisting of a CCRN, Critical Care Technician and a RT. All patient transports should have 2 EMTs on board to assist with ambulance operations.

BETTER		
<i>If the patient:</i>	<i>Crew (in addition to the EMT/NRP driver):</i>	<i>References:</i>
Is ventilated with IV/IO infusion medication, but no central lines or arterial lines	NRP trained in ventilator management ABG should be obtained within 30 minutes of transport. If time allows, patient should be placed on transport ventilator for at least 15 minutes prior to transport.	ALS standards Commission on Accreditation of Medical Transport Systems (CAMTS) 11 th Edition https://www.camts.org/standards/ https://its.amedd.army.mil/assets/docs/cpgs/Prehospital_En_Route_CPGs/Standard_Medical_Operating_Guidelines_(SMOG)_for_Critical_Care_Flight_Paramedics_2020.pdf Procedure A-XII
Is ventilated with central line, or arterial line, or chest tube	At least 2 providers trained at the NRP level or above (physician (MD/DO), physician’s assistant (PA), nurse practitioner (NP), or registered nurse (RN)) Primary care provider requirement: > 3 years ED, ICU, or critical care experience.	Emergency Critical Care standards CAMTS 11 th Edition https://www.camts.org/standards/
Above criteria AND complex ventilator settings OR > 4 IV/IO infusions	Above requirements AND 1 crew member must be an RN with Certified Flight RN, Critical Care RN, or Certified Transport Registered Nurse within 2 years of hire, or equivalent national certification. At least 1 critical care transport provider shall be licensed as a MD/DO, PA, APRN, or RN with documented competency and experience in the provision of critical care in a tertiary critical care unit, commensurate with the type and acuity of patient requiring transport.	Intensive Care Standards CAMTS 11 th Edition https://www.camts.org/standards/ Para 1.2.3 Critical Care Transport Team Association of Critical Care Transport-Critical Care Transport Standards-Version 1.0 ©2016 (AACT is a professional organization recommendation but not a certifying organization.)

BEST	
<i>If the patient:</i>	<i>Crew (in addition to the driver):</i>
requires critical care	Military or civilian trained and equipped critical care transport crew (Ground, Rotary, or Fixed Wing)

2. Additional considerations for interfacility transport include:
 - a. On-line Medical Control. On-line Medical Control must be available to transport critically ill patients.
 - b. Training. Personnel involved in interfacility transports should be trained on ambulances, facility transport ventilators, infusion pumps and all required equipment. Additionally, NRPs with critical care training:

Clinical Management of COVID-19, v4

Critical Care Paramedic Program (CCEMT-P), Certified Critical Care Paramedics (C-CCPs), Certified Flight Paramedics (FP-Cs), or individuals with previous critical care experience should be tasked as primary transport personnel given their increased education/experience.

- c. Ventilators. NRP and RNs should be deemed proficient in ventilator operation and management by the local EMS Medical Director prior to performing patient transport. Ventilated patients should be transported with physician documented orders which detail ventilator settings. All patients will be monitored with wave-form capnography. If a BVM is utilized for transport, or if use of the BVM becomes necessary during transport, a positive-end expiratory pressure (PEEP) valve must be applied and dialed to the ventilator PEEP setting. Ventilators and BVMs should be equipped with HEPA filters.
 - d. IV/IO Infusions. Many pre-hospital NRP infusions are currently delivered without the use of an infusion pump (epinephrine, norepinephrine, dopamine, amiodarone, and magnesium sulfate), however any infusion for an interfacility transfer should be on an infusion pump. Medications not detailed in the formulary outlined by EMS protocols are authorized with a written physician order. Orders should specify the name of the medication, the drug concentration, and the infusion rate. Infusions must be initiated by the sending facility. Infusions will be maintained at the physician-prescribed dosing regimen. Alterations to dosing regimens require authorization from a physician, preferably, On-line Medical Control. Rapid deterioration in patient clinical status negates the requirement for physician authorization (e.g. vasopressor titration).
 - e. Prior to placing a transport request, MTF in-patient units should communicate with local EMS Medical Directors or attending Emergency Department physicians to determine transport capabilities. If possible, patient documentation (to include compact discs containing images) should be prepared prior to transport crew arrival.
3. If trained healthcare personnel are severely limited, local Medical Directors should partner with MTF and Logistics leadership to discuss the use of licensed drivers/ government owned vehicles to transport of low acuity patients.

EMS Personnel Precautions for Procedures.

1. See CDC guidance for aerosol generating procedures: <https://www.cdc.gov/coronavirus/2019-ncov/hcp/guidance-for-ems.html>.
2. If patient presentation allows, EMS personnel providing care to a patient suspected of having COVID-19 should contact Medical Control and/or follow local protocols before initiating an aerosol-generating procedure.
3. Nebulized medications for known or suspected COVID-19 patients should be limited given the risk of virus transmission. It is recommended that local Medical Directors work with MTF leadership to obtain single-use albuterol metered-dose inhalers with spacers for prehospital use. If an aerosol-generating procedure is required/recommended, the doors to the patient compartment of the ambulance should remain open to allow ventilation of the area during these procedures. If the ambulance is equipped with an HVAC system it should remain on during patient transport.
4. If used, BVMs, SGAs, and ET tubes should have a HEPA/viral filter attached. If the EMS agency has access to ventilators, units should contact the specific ventilator manufacturer for additional guidelines and to obtain part numbers for compatible HEPA/viral filters.

Mechanical CPR.

1. The American College of Cardiology recommends use of mechanical CPR devices on PUIs or patients with confirmed COVID-19 who arrest (<https://www.acc.org/latest-in-cardiology/articles/2020/03/19/13/14/cvd-in-the-setting-of-covid-19-coronavirus-considerations-to-prepare-patients-providers-health-systems>). See updated AHA ACLS and PALS Cardiac Arrest Algorithms for best practice for COVID-19 patients (April 2020). (<https://professional.heart.org>)
2. Local Medical Directors & EMS/Fire Leadership are responsible for ensuring personnel education of device indications/contraindications, application, and cleaning of mechanical CPR devices. Initial and continuing education should be documented in training records.
3. Devices should be cleaned according to CDC recommendations for known or suspected COVID-19 patients.

4. Contact the device manufacturer for additional recommendations.

Cleaning EMS Transport Vehicles After Transporting a PUI or Patient with Confirmed COVID-19: See current CDC guidelines: <https://www.cdc.gov/coronavirus/2019-ncov/hcp/guidance-for-ems.html>

Follow-up for EMS Personnel after Caring for a PUI or Patient with Confirmed COVID-19.

1. Local public health and infectious disease authorities should be notified regarding PUIs or confirmed COVID-19 patients so that appropriate follow-up monitoring can occur.
2. EMS personnel who have been exposed to a patient with suspected or confirmed COVID-19 should notify their chain of command to ensure appropriate follow-up.
3. EMS agencies should develop local policies for assessing exposure risk and the management of EMS personnel potentially exposed to COVID-19. Decisions for monitoring and quarantine should be made in consultation with public health and infectious disease authorities.
4. EMS personnel should be alert for fever or respiratory symptoms (e.g. cough, shortness of breath, sore throat). If symptoms develop, it is recommended that they self-isolate and notify their primary care provider and/or public health authority to arrange for evaluation.

EN ROUTE CRITICAL CARE CONSIDERATIONS FOR PERSONS WITH COVID-19

1. The DoD has issued a COVID-19 specific Force Health Protection Guidance (Supplement 5) to DoDI 6000.11 “Patient Movement.” Attachment 1 of this supplement provides guidance for the air movement of COVID-19 patients and COVID-19 exposed persons. “Treatment in Place” remains the primary guidance. The document ([https://www.whs.mil/Portals/75/Coronavirus/FHP%20Guidance%20\(Supplement%205\)%20-%20DoD%20Guidance%20for%20Mvmt.%20and%20Med.%20Treatment.pdf?ver=2020-04-08-142257-760](https://www.whs.mil/Portals/75/Coronavirus/FHP%20Guidance%20(Supplement%205)%20-%20DoD%20Guidance%20for%20Mvmt.%20and%20Med.%20Treatment.pdf?ver=2020-04-08-142257-760)) discusses procedures for obtaining an exception to policy, preferred means of transport, airframe selection, biocontainment, cabin airflow, patient placement, and infection control. It also discusses specific safety precaution considerations regarding mechanical ventilators, clinical specimen management, waste disposal, aircraft cleaning, logistical planning, and in-flight emergencies. Repeating the document is not an intent of this guidance. Medical planners and clinicians are strongly encouraged to review FHP 5 when considering transport for patients with COVID-19.
2. The following suggested treatment plans exist for Critical Care Air Transport (CCAT) teams transporting critically ill COVID-19 patients under an ETP. This section highlights considerations for flight. CCAT team members should review the relevant critical care topics elsewhere in this PMG.
3. **Biocontainment:** Civil Aviation Assets (e.g., Phoenix Air Group) should be the primary means of patient movement if capable. For USAF CCAT or aeromedical evacuation (AE) teams tasked to transport patients on USAF aircraft, the best practice is to use a biocontainment module like the DoD’s transport isolation system (TIS), followed by transport in open aircraft. AMC has issued AMC COVID-19 PMP, which discusses best practices for transport in open aircraft and offers guidance on appropriate PPE measures. FHP Supplement 5 discusses these measures as well.
4. **Initial assessment:** The pre-evacuation assessment requires additional time due to the complexity of these patients. Consider continuing to treat in place those not requiring mechanical ventilation or depleting local resources in austere locations. In environments with fewer resource constraints, consider allowing patients to declare themselves on the ground to require mechanical ventilation before transport. Teleconsultation over time may assist in the management of non-ventilated patients and help determine the need for mechanical ventilation before transport.
5. **Neurologic:** Sedation can be challenging in the controlled environment of the ICU and even more complicated in flight. Adjust management to conserve common medications in short supply. Reports indicate a ceiling dose of propofol (30 mcg/kg/min), with little effect of increasing infusions. Consider combinations of acetaminophen (IV/PO), opiates (gtts/IVP/PO), propofol (gtts), atypical antipsychotics (IV/IM/PO), and sub-dissociative ketamine (IV) for a multi-modal approach to patient analgesia/sedation. Use caution with dexmedetomidine due to reports of significant bradycardia. Utilize low dose benzodiazepines (IVP) as a last

Clinical Management of COVID-19, v4

resort due to their association with delirium and prolonged mechanical ventilation. Continue the same or a more aggressive analgesia/sedation strategy for flight if a patient is receiving neuromuscular blockade.

- Pulmonary:** Anecdotal evidence suggests a subset of COVID-19 patients whose hypoxia rapidly corrects with awake proning and supplemental oxygen. Additionally, anecdotal evidence indicates some patients fare poorly with early intubation. Therefore, the traditional CCAT practice of intubation solely for the flight may not be appropriate. CCAT team leads must consider the difficulty in predicting when COVID-19 patients will deteriorate, and anticipate a need for in-flight intubation. Plan patient placement and airflow characteristics on the aircraft to minimize aircraft/crew exposure in case of in-flight intubation. Signs of persistent respiratory distress, complaints of dyspnea, persistent hypoxia with SpO₂ <92%, or a pH of <7.2 despite preflight proning and conventional supplemental oxygen likely indicate a need for intubation. Careful observation during flight and avoiding intubation may be appropriate for patients whose symptoms improve with awake proning and supplemental oxygen. Consider using the low PEEP table for intubated patients with a low driving pressure (P_{plat} – PEEP) and 6-8 ml/kg IBW tidal volume. Conversely, for patients with moderate to severe ARDS and less compliant lungs with a higher driving pressure, consider using the high PEEP table.

Cabin Altitude Restriction (CAR): Consider a CAR when transporting non-intubated patients requiring supplemental oxygen or intubated patients on high PEEP or high FiO₂ in anticipation of potential in-flight patient decompensation. During pre-mission planning, the CCATT lead should discuss a CAR with the TPRMC validating flight surgeon. A lower CAR is associated with a longer duration of the flight. A sea-level CAR can provide an increased safety buffer if the aircraft is capable.

Prone Positioning: Strongly consider lung team consultation before transporting intubated patients in the prone position or patients requiring PEEP >14, FiO₂ >60%. For intubated patients that are to be transported in the prone position, initiate prone positioning preflight with adequate time (i.e., >4 hours or physician discretion) to verify patient stability and adequate ABG. After proning, wean FiO₂ to maintain SpO₂ >92%. Ideally, design a patient load strategy allowing access for bilateral chest tubes placement, particularly when utilizing high PEEP. Prone positioning complicates the treatment of cardiac arrhythmias, cardiac arrest, pneumothorax, and shock. Before proning, consider placing cardioversion pads for dysrhythmia treatment. Leave the patient on the ventilator to avoid additional aerosolized particles during cardiac arrest. Consider CPR in the prone position during cardiac arrest (see AHA 2010 guidelines). Avoid rotation (proning/reversal) of intubated patients during flight. Refer to **Appendix D** for further discussion of prone positioning. It highlights the need for thorough patient handoff preflight. Be aware prone positioning requires frequent repositioning and padding to prevent pressure wounds. Prone-positioned patients may have intermittent scheduled times in a supine position. Cautiously consider patient movement during the supine period, as FiO₂ requirements usually increase upon supination and may continue to increase throughout the supine period.

Refer to **Appendix G** for a demonstration of the transport ventilator setup.

Table 11 shows the required FiO₂ to maintain a constant PaO₂ at different altitudes. It may be useful when assessing stability for flight and the need for cabin altitude restriction.

Table 11. Altitude Physiology Table

Altitude (ft)	Barometric Pressure (mmHg)	FiO ₂ Required to Maintain Constant PaO ₂ ¹								PiO ₂ While Breathing:					Gas Volume Expansion (% at Sea Level)	
		0.21	0.40	0.60	0.80	100	0.21	0.40	0.60	0.80	100					
16000	412	0.41	0.59	0.78	0.98						76	146	219	292	365	184
14000	446	0.37	0.53	0.71	0.89						83	160	239	319	399	170
12000	483	0.34	0.49	0.65	0.81	0.98					91	174	262	349	436	157
10000	523	0.31	0.45	0.60	0.75	0.90					100	190	286	381	476	145
8000	564	0.29	0.41	0.55	0.69	0.83	0.96				108	207	310	414	517	135
6000	609	0.27	0.38	0.51	0.63	0.76	0.89				118	226	337	450	562	125
4000	656	0.24	0.35	0.47	0.58	0.70	0.82	0.94			127	244	365	487	609	116
2000	707	0.23	0.32	0.43	0.54	0.65	0.76	0.86	0.97		138	264	396	528	660	107
Sea Level	760	0.21	0.30	0.40	0.50	0.60	0.70	0.80	0.90	1.0	149	285	428	570	713	100

Clinical Management of COVID-19, v4

7. **Cardiovascular:** Optimize electrolytes (e.g., Ca, Mg, and K) preflight due to the incidence of tachydysrhythmias. Consider requesting electrolyte supplementation preflight due to the allowance standard limitations. Review the EKG. Consider holding QTc prolonging medications (e.g., chloroquine derivatives, antipsychotics, etc.) if the QTc >500 ms. Due to the incidence of cardiomyopathy, obtain an echo preflight to inform treatment if in-flight shock develops. For intubated patients, place a CVC preflight, in case a vasopressor requirement develops during the flight.
8. **Renal:** AKI is common in COVID-19 patients. Renally adjust medication dosage and convert renally metabolized medications as appropriate (e.g., morphine -> dilaudid, or lovenox -> heparin).
9. **Gastrointestinal:** Continue stress ulcer prophylaxis. Continue post-pyloric enteric feeds, as suggested in **Appendix J**. OGT should be placed preflight and on intermittent suction.
10. **Fluids:** Euvolemia is the goal. If hypovolemia is suspected, consider low volume (250–500ml) boluses of balanced crystalloid solutions. Anticipate K and Mg replacement need if patient diuresis is ongoing. Recall potassium is not in the CCATT allowance standard.
11. **Heme:** Ensure administration of DVT prophylaxis. Anecdotal evidence suggests some COVID-19 patients are affected by pulmonary microvascular thrombosis. Guidance for treatment is not within the realm of this section of the PMG. Recommend discussion with sending/receiving critical care specialists to determine the dose of prophylactic anticoagulation.
Non-COVID-19 patient transports may continue within the PM system. Utilize standard transmission-based precautions in accordance with AFI 48-307. Movements should be requested when it is essential to provide appropriate care while minimizing opportunities for transmission of pathogens within and between theaters and countries.

PUBLIC HEALTH CONSIDERATIONS AND RESPONSE

1. **Public Health Emergency Management (PHEM)**
 - a. Primary reference: DoD Instruction (DoDI) 6200.03 (Public Health Emergency Management (PHEM) within the DoD); March 28, 2019.
 - b. The Public Health Emergency Officer (PHEO). PHEOs provide military commanders with guidance and recommendations on preparing for, declaring, responding to, mitigating, and recovering from public health emergencies. PHEO responsibilities fall into 10 major categories, including: advising the military commander regarding the declaration of a public health emergency and the implementation of emergency health powers, assisting in public affairs risk communications, including dissemination of health protection measures detailed in the Health Protection Condition (HPCON) framework in coordination with the Public Affairs Officer, coordinating with other DoD Components, civilian state, legal, tribal, and territories (SLTT), other federal agencies, and others.
 - c. Declaring a Public Health Emergency (PHE): Commanders must be prepared to make timely decisions in order to protect lives, property, and infrastructure and enable DoD installations and/or military commands to sustain mission-critical operations and essential services. Declaration of a PHE allows the installation commander access to the medical emergency powers described in DoDI 6200.03, including restriction of movement (ROM), directing examinations and testing, and controlling or restricting the distribution of commodities, and others. The process by which the Commander makes decision to declare a PHE is summarized in the DoDI.
 - d. Health Protection Condition (HPCON) levels are used during a health emergency to communicate what health protection measures are currently being used to prevent the spread of disease in the population. The decision to adjust HPCON posture is not based on strictly objective criteria - rather, it is based on a constellation of factors. These factors are similar to deciding whether to declare a Public Health Emergency. The decision to adjust HPCON levels may be more political than one driven by public health statistics. If the public health recommendations are being mostly followed with few exceptions, and the command continues to support public health interventions as they are needed, then the specific HPCON level is less (from a strictly public health perspective) relevant. For example:
 - i. Evidence of repeated noncompliance with public health guidance among the installation community.

Clinical Management of COVID-19, v4

- ii. Difficulties in getting supervisors and/or employees to follow public health guidelines for example, regarding guidance on returning to work.
 - iii. Any difficulties encountered by public health personnel in conducting duties required to investigate a public health threat, e.g. a case investigation and contact tracing.
 - iv. A sense from commanders that the installation community is panicking, or is on the verge of panic.
 - e. Further information on HPCONs can be found in the DoDI and the Army Public Health Center (APHC), <https://phc.amedd.army.mil/topics/campaigns/covid19/Pages/HPCON.aspx>
 - f. PHEM training courses (which is required by DoDI 6200.03) and POCs can be found at: <https://www.health.mil/Training-Center/Defense-Medical-Readiness-Training-Institute/Public-Health-Emergency-Management-Course>
2. **Non-pharmaceutical interventions (NPIs)** are critical when no vaccine or therapeutic is available to mitigate a public health threat. NPIs directed towards control of COVID-19, for example, were largely based on the CDC's "Community Mitigation Guidelines to Prevent Pandemic Influenza—United States, 2017," at: <https://www.cdc.gov/mmwr/volumes/66/rr/rr6601a1.htm>. These include:
- a. Personal Protective Measures (PPMs) for Everyday Use
 - i. Voluntary home isolation (i.e., staying home when ill or self-isolation)
 - ii. Respiratory etiquette
 - iii. In health care settings, screening for respiratory symptoms immediately upon entry.
 - iv. Hand hygiene
 - b. Personal Protective Measures (PPMs) Reserved for Pandemics. During a pandemic, the PPMs described above should be strengthened and augmented with additional measures:
 - i. Active, rapid identification of persons having symptoms consistent with COVID-19, followed by referral for testing and home isolation.
 - ii. Identification and home quarantine of non-ill household members or other close contacts of persons with COVID-19. See "contact tracing" section below.
 - iii. Use of face masks or cloth face coverings by well persons. **IMPORTANT NOTE:** respirators (e.g. N95, PAPR) are medical supplies and are reserved for use by at-risk medical providers. See information differentiating masks and respirators at the APHC website: <https://phc.amedd.army.mil/topics/campaigns/covid19/Pages/healthcare.aspx>
 - iv. Preemptive or reactive school and work closures/dismissals.
 - v. Elimination or reduction of other mass gatherings.
 - vi. Social/physical distancing measures to no less than 6 feet separation.
 - vii. Environmental surface cleaning measures in all settings.
3. **Contact tracing** (also called contact investigation): When a person gets sick, they are interviewed by public health personnel to make a contact list of other individuals who they might have exposed. The steps include:
- a. Contact identification: Each case of COVID-19 is interviewed to identify contacts (people) and activities starting 2 days before symptoms started.
 - b. Contact notification: All contacts are notified that they may have been exposed to COVID-19.
 - c. Contact follow-up: Regular follow-up may be needed with all contacts to monitor for symptoms and provide additional information about COVID-19.
- PLEASE NOTE! Contact tracing is very time consuming and requires large amount of man power! Therefore, force multiplying protocols were developed to train nonmedical individuals to assist in the process. Additional information on contact tracing, including a toolkit for contact tracing, can be found at the APHC website: <https://phc.amedd.army.mil/topics/campaigns/covid19/Pages/healthcare.aspx>
4. **Risk assessment for potential COVID-19 exposures:**
- a. Travel exposures: Includes travel from a country with [widespread ongoing transmission](#) (currently includes all countries), or travel on cruise ship or river boat. Public health actions include (<https://www.cdc.gov/coronavirus/2019-ncov/php/risk-assessment.html>): (1) Stay home until 14 days after arrival and maintain a distance of at least 6 feet (2 m) from others; (2) Self-monitor for symptoms and check temperature twice a day; (3) Avoid contact with [people at higher risk for severe illness](#); and (4) Follow [CDC guidance](#) if symptoms develop.

- b. Community-based exposures (<https://www.cdc.gov/coronavirus/2019-ncov/php/public-health-recommendations.html>):
 - i. Applies to: Household members, intimate partners, individuals providing care in a household without using recommended [infection control precautions](#), and Individuals who have had close contact (< 6 feet) for a prolonged period of time
 - ii. Exposure to: Person with symptomatic COVID-19 during period from 48 hours before symptoms onset until meets criteria for [discontinuing home isolation](#) (can be a laboratory-confirmed disease or a clinically compatible illness [in a state or territory with widespread community transmission](#))
 - iii. Public health actions: same as under “travel exposures” above
 - c. Health care personnel (HCP) exposures (<https://www.cdc.gov/coronavirus/2019-ncov/hcp/guidance-risk-assesment-hcp.html>).
 - i. HCP exposures in areas with **moderate to substantial transmission**: facilities should consider foregoing formal contact tracing and work restriction for HCP in favor of universally applied screening and source control strategies. Proper adherence to currently recommended infection control practices, including all recommended PPE, should protect HCP having prolonged close contact with patients infected with COVID-19. However, to account for any inconsistencies in use or adherence that could result in unrecognized exposures, HCP should still perform self-monitoring with delegated supervision.
 - ii. HCP in areas with **minimal to no community transmission** may have the ability to apply risk assessment and work restrictions:
 - a. HCP who have had prolonged (≥ 15 minutes) close (< 6 feet) contact with patients with COVID-19 (**beginning 48 hours before onset of symptoms**) and the HCP was: 1) not wearing a respirator or facemask (n.b. not a face covering), 2) not wearing eye protection, or 3) now wearing all recommended PPE (i.e. gown, gloves, eye protection, and respirator) while performing an aerosol-generating procedure) should be excluded from work for 14 days and self-monitor for symptoms.
 - b. HCP with exposures other than those listed above have no work restrictions.
 - iii. HCP with travel or community exposures should consult occupational health.
5. **Guidance for when to discontinue isolation.**
- a. Non-health care settings (<https://www.cdc.gov/coronavirus/2019-ncov/hcp/disposition-in-home-patients.html>)
 - i. *Test-based strategy*. Exclude from work until:
 - a) Resolution of fever without the use of fever-reducing medications **and**
 - b) Improvement in respiratory symptoms (e.g., cough, shortness of breath), **and**
 - c) Negative results of an FDA Emergency Use Authorized molecular assay for COVID-19 from at least two consecutive nasopharyngeal swab specimens collected ≥ 24 hours apart (total of two negative specimens)
 - ii. *Non-test-based strategy*. Exclude from work until:
 - a) At least 3 days (72 hours) have passed *since recovery* defined as resolution of fever without the use of fever-reducing medications **and** improvement in respiratory symptoms (e.g., cough, shortness of breath); **and**
 - b) At least 7 days have passed *since symptoms first appeared*
 - b. Return to work criteria for HCP: The *test-based strategy* or the *non-test-based strategy* (aka *symptom-based strategy*) above may be used with the following differences:
 - i. HCP *with symptoms* may use either strategy.
 - ii. HCP *without symptoms* may use the *test-based strategy* or a *time-based strategy*, in which HCP are excluded from work until 10 days have passed since the date of their first positive COVID-19 diagnostic test. If they develop symptoms, then the *symptom-based* or *test-based strategy* should be used.
- https://www.cdc.gov/coronavirus/2019-ncov/hcp/return-to-work.html?CDC_AA_refVal=https%3A%2F%2Fwww.cdc.gov%2Fcoronavirus%2F2019-ncov%2Fhealthcare-facilities%2Fhcp-return-to-work.html

Clinical Management of COVID-19, v4

6. **Reporting and Surveillance:** All confirmed and probable cases of COVID-19 must be reported to design, inform, and evaluate control and prevention efforts. Cases are reported by military public health personnel to BOTH: 1) military and 2) civilian public health authorities. Military service members and other beneficiaries must be reported through military public health authorities via the Disease Reporting System internet (DRSi) in coordination with the Service-specific public health chain of command. All cases must also be reported to the supporting local or state health department according to state requirements. All DoD medical reporting entities should report cases of COVID-19 to the DRSi using the "COVID-19" and answer all event-related questions in the report. Cases must be classified according to the most recent DoD case definition for COVID-19.

WHOLE OF GOVERNMENT RESPONSE IN COORDINATION OF RESOURCES

On 13 Mar 2020, President Trump declared a nationwide emergency under Sec. 501(b) of the Stafford Act, increasing support to HHS in this role as the lead federal agency for the federal government's response to the COVID-19 pandemic. Under this declaration, FEMA, in coordination with HHS, was empowered to assist state, local, tribal, territorial governments and other eligible entities to access resources made available through the Stafford Act.

HHS has many resources to leverage in the federal response to COVID-19, including the Strategic National Stockpile (SNS). The SNS has ventilators, medications, personal protective equipment and other important equipment and supplies that may be requested for COVID-19 response where state and local resources are overwhelmed or anticipated to be overwhelmed. SNS depots are located around the country by region. There is a Defense Coordinator at regional FEMA offices to coordinate requests to/from civilian and military hospitals and other entities for resources. MTFs can identify anticipated shortages and push a request through their local unit Crisis Action Team to the Regional FEMA Defense Coordinator for items in the SNS. It is recommended that facilities leverage available resources before running out of critical items such as PPE.

HHS link to Resources: <https://www.phe.gov/emergency/Tools/Pages/default.aspx>

HHS Regional Emergency Coordinators Contact List: <https://www.phe.gov/Preparedness/responders/rec/Pages/default.aspx>

State FEMA Office contacts: <https://www.fema.gov/emergency-management-agencies>

OTHER CONSIDERATIONS RELATED TO COVID-19

Facilities.

Medical Heating, Ventilation and Air Conditioning (HVAC) Systems.

1. DHA Facilities Enterprise recommends maintaining building ventilation systems in balance and compliant. Attempts to adjust without professional mechanical engineering support may cause harm and rework later.
2. Medical facilities (hospitals/clinics) or administrative facilities are recommended not to alter the HVAC system operations or filtration in any way due to the outbreak of COVID-19.
3. Building maintenance personnel should not be exposed to COVID-19 unless they are physically in the same room as an infected person or come in contact with surfaces that have not been disinfected (such as air filters). No special COVID-19 PPE is required for maintenance personnel unless they are charged with disinfecting surfaces or working where infected persons may have deposited live virus. In those cases, the maintenance personnel should follow CDC guidelines.
4. Although it is not known exactly how long the virus can survive on a surface outside the human carrier, some reports suggest up to 4 days on some materials.
5. If a maintenance worker becomes infected with COVID-19, it is recommend to clean all surfaces the worker may have been in contact with for the past 7 days. A review of all work orders completed by the infected maintenance staff will aid in discovering where and when the employee contacted other surfaces.
6. DHA Facilities Enterprise does NOT recommend increasing filter media such as changing Minimum Efficiency Reporting Value (MERV) rated filters to High Efficiency Particulate Air (HEPA) if it is being done purely in hope of stopping the spread of COVID-19. MTFs should not add higher rated filters to existing HVAC systems

Clinical Management of COVID-19, v4

without proper engineering management since the HVAC system may become imbalanced which could result in loss of isolation rooms. Care must be taken not to exceed the design performance of the HVAC as it will likely reduce equipment life with little or no positive impact.

7. The use of Ultraviolet (UV) lights in the HVAC system (e.g., AHU or ductwork) is not recommended for COVID mitigation.
8. The use of mobile or fixed air scrubber with integral HEPA or Ultra-Low Particulate Air (ULPA) filter may be used to increase the air changes in a room. Air scrubbers when used to create negative pressure rooms must be cautious in discharging exhaust air to the outside of the building or into the return air system. Coordination with Facilities Management, a professional mechanical engineer, Industrial Hygiene and Infection Control team to ensure virus exposures are minimized and tested prior to room use.
9. There are many new and evolving technologies coming out of industry today as a result of the COVID pandemic that claim to have outstanding results in mitigating COVID-19 viruses. Many of these systems are either experimental or have not been proven in the healthcare setting. DHA FE cannot advocate the use these systems at this time. Should a MTF wants to install a new technology, we recommend a multi-discipline support team with engineers, infection control, and industrial hygiene practitioners to review and validate the product before purchasing to ensure it meets the building's requirements, is maintainable, and can produce the desired mitigation for the MTF.
10. When installing Plexiglas sneeze guards/barriers at reception desk or pharmacy window areas, the MTF should consider which ones should be permanently installed while other may be temporarily installed. Those reception areas with high volume should be more durable in construction while the low volume may be temporary. Also consider the choice of barrier material that is easily cleaned.
11. Due to dental procedures being high aerosolizing, it is recommend to use a room with a door and an air scrubber to create an Airborne Infection Isolation Room (with negative pressure in relation to the corridor) when treating suspected or infected COVID patients. The best option is to utilize an oral surgery suite with a door. If there is no door, it is recommended to create a temporary door with a plastic door curtain or magnetic door curtain. Dental staff should work with their local facilities management and industrial hygiene staff to ensure a negative pressure condition is created for the room before starting treatment.
12. MTFs should follow their Joint Commission required Water Management Plans for reopening their closed facilities to ensure opened facilities are safe to include eye wash stations, cooling towers, hot and cold domestic water systems and water heaters. The CDC has information on reopening closed facilities to include the stagnant water, mold and other issues, available at: https://www.cdc.gov/coronavirus/2019-ncov/php/building-water-system.html?deliveryName=USCDC_248_DM25447.

REFERENCES

1. Wu Z, McGoogan JM. Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72314 Cases From the Chinese Center for Disease Control and Prevention. JAMA. 2020.
2. TeamNCPERE. Vital surveillances: the epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19) – China. China CDC Weekly. 2020;2(8):113-22.
3. Yang X, Yu Y, Xu J, Shu H, Xia J, Liu H, et al. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. Lancet Respir Med. 2020.
4. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. Lancet. 2020;395(10223):497-506.
5. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. Lancet. 2020.
6. Du Y, Tu L, Zhu P, Mu M, Wang R, Yang P, et al. Clinical Features of 85 Fatal Cases of COVID-19 from Wuhan: A Retrospective Observational Study. Am J Respir Crit Care Med. 2020.
7. Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, et al. Clinical Characteristics of Coronavirus Disease 2019 in China. N Engl J Med. 2020.

Clinical Management of COVID-19, v4

8. Li Q, Guan X, Wu P, Wang X, Zhou L, Tong Y, et al. Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus-Infected Pneumonia. *N Engl J Med*. 2020.
9. Lauer SA, Grantz KH, Bi Q, Jones FK, Zheng Q, Meredith HR, et al. The Incubation Period of Coronavirus Disease 2019 (COVID-19) From Publicly Reported Confirmed Cases: Estimation and Application. *Ann Intern Med*. 2020.
10. Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, et al. Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. *JAMA*. 2020.
11. Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet*. 2020;395(10223):507-13.
12. Holshue ML, DeBolt C, Lindquist S, Lofy KH, Wiesman J, Bruce H, et al. First Case of 2019 Novel Coronavirus in the United States. *N Engl J Med*. 2020;382(10):929-36.
13. Pan L, Mu M, Ren HG, Yang P, Sun Y, Wang R, et al. Clinical characteristics of COVID-19 patients with digestive symptoms in Hubei, China: a descriptive, cross-sectional, multicenter study. 2020.
14. American Academy of Otolaryngology-Head and Neck Surgery Coronavirus Disease 2019: Resources 2020 [Available from: <https://www.entnet.org/content/coronavirus-disease-2019-resources>].
15. Cheung KS, Hung IF, Chan PP, Lung KC, Tso E, Liu R, et al. Gastrointestinal Manifestations of SARS-CoV-2 Infection and Virus Load in Fecal Samples from the Hong Kong Cohort and Systematic Review and Meta-analysis. *Gastroenterology*. 2020.
16. Richardson S, Hirsch JS, Narasimhan M, Crawford JM, McGinn T, Davidson KW, et al. Presenting Characteristics, Comorbidities, and Outcomes Among 5700 Patients Hospitalized With COVID-19 in the New York City Area. *JAMA*. 2020.
17. Jacobs M, Muthukumar S, Panneer Selvam A, Engel Craven J, Prasad S. Ultra-sensitive electrical immunoassay biosensors using nanotextured zinc oxide thin films on printed circuit board platforms. *Biosens Bioelectron*. 2014;55:7-13.
18. Galvan Casas C, Catala A, Carretero Hernandez G, Rodriguez-Jimenez P, Fernandez Nieto D, Rodriguez-Villa Lario A, et al. Classification of the cutaneous manifestations of COVID-19: a rapid prospective nationwide consensus study in Spain with 375 cases. *Br J Dermatol*. 2020.
19. Recalcati S. Cutaneous manifestations in COVID-19: a first perspective. *J Eur Acad Dermatol Venereol*. 2020;34(5):e212-e3.
20. Interim Clinical Guidance for Management of Patients with Confirmed Coronavirus Disease (COVID-19): Center for Disease Control (CDC); 2020 [updated March 7, 2020. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/hcp/clinical-guidance-management-patients.html#foot09>].
21. Petrilli CM, Jones SA, Yang J, Rajagopalan H, O'Donnell L, Chernyak Y, et al. Factors associated with hospital admission and critical illness among 5279 people with coronavirus disease 2019 in New York City: prospective cohort study. *BMJ*. 2020;369:m1966.
22. Breslin N, Baptiste C, Miller R, Fuchs K, Goffman D, Gyamfi-Bannerman C, et al. COVID-19 in pregnancy: early lessons. *Am J Obstet Gynecol MFM*. 2020:100111.
23. Chen H, Guo J, Wang C, Luo F, Yu X, Zhang W, et al. Clinical characteristics and intrauterine vertical transmission potential of COVID-19 infection in nine pregnant women: a retrospective review of medical records. *Lancet*. 2020;395(10226):809-15.
24. Di Mascio D, Khalil A, Saccone G, Rizzo G, Buca D, Liberati M, et al. Outcome of Coronavirus spectrum infections (SARS, MERS, COVID 1 -19) during pregnancy: a systematic review and meta-analysis. *Am J Obstet Gynecol MFM*. 2020:100107.
25. Mullins E, Evans D, Viner RM, O'Brien P, Morris E. Coronavirus in pregnancy and delivery: rapid review. *Ultrasound in Obstetrics & Gynecology*. n/a(n/a).
26. Della Gatta AN, Rizzo R, Pilu G, Simonazzi G. COVID19 during pregnancy: a systematic review of reported cases. *Am J Obstet Gynecol*. 2020.
27. Kam KQ, Yung CF, Cui L, Lin Tzer Pin R, Mak TM, Maiwald M, et al. A Well Infant with Coronavirus Disease 2019 (COVID-19) with High Viral Load. *Clin Infect Dis*. 2020.
28. Cai J, Xu J, Lin D, Yang Z, Xu L, Qu Z, et al. A Case Series of children with 2019 novel coronavirus infection: clinical and epidemiological features. *Clin Infect Dis*. 2020.
29. Wei M, Yuan J, Liu Y, Fu T, Yu X, Zhang ZJ. Novel Coronavirus Infection in Hospitalized Infants Under 1 Year of

Clinical Management of COVID-19, v4

Age in China. JAMA. 2020.

30. Cui Y, Tian M, Huang D, Wang X, Huang Y, Fan L, et al. A 55-Day-Old Female Infant infected with COVID 19: presenting with pneumonia, liver injury, and heart damage. J Infect Dis. 2020.
31. Dong Y, Mo X, Hu Y, Qi X, Jiang F, Jiang Z, et al. Epidemiological Characteristics of 2143 Pediatric Patients With 2019 Coronavirus Disease in China. Pediatrics. 2020.
32. Liu W, Zhang Q, Chen J, Xiang R, Song H, Shu S, et al. Detection of Covid-19 in Children in Early January 2020 in Wuhan, China. N Engl J Med. 2020.
33. CDC COVID-19 Response Team. Coronavirus Disease 2019 in Children - United States, February 12-April 2, 2020. MMWR Morb Mortal Wkly Rep. 2020 Apr 10;69(14):422-426. [Available from: <https://www.cdc.gov/mmwr/volumes/69/wr/mm6914e4.htm>].
34. Xiao AT, Tong YX, Zhang S. Profile of RT-PCR for SARS-CoV-2: a preliminary study from 56 COVID-19 patients. Clin Infect Dis. 2020.
35. Xu Z, Shi L, Wang Y, Zhang J, Huang L, Zhang C, et al. Pathological findings of COVID-19 associated with acute respiratory distress syndrome. Lancet Respir Med. 2020.
36. Klok FA, Kruip M, van der Meer NJM, Arbous MS, Gommers D, Kant KM, et al. Incidence of thrombotic complications in critically ill ICU patients with COVID-19. Thromb Res. 2020.
37. Varga Z, Flammer AJ, Steiger P, Haberecker M, Andermatt R, Zinkernagel AS, et al. Endothelial cell infection and endotheliitis in COVID-19. Lancet. 2020.
38. Dichter JR, Kanter RK, Dries D, Luyckx V, Lim ML, Wilgis J, et al. System-level planning, coordination, and communication: care of the critically ill and injured during pandemics and disasters: CHEST consensus statement. Chest. 2014;146(4 Suppl):e87S-e102S.
39. Sprung CL, Zimmerman JL, Christian MD, Joynt GM, Hick JL, Taylor B, et al. Recommendations for intensive care unit and hospital preparations for an influenza epidemic or mass disaster: summary report of the European Society of Intensive Care Medicine's Task Force for intensive care unit triage during an influenza epidemic or mass disaster. Intensive Care Med. 2010;36(3):428-43.
40. Stroud C, Altevogt B, Nadig L. Crisis Standards of Care: Summary of a Workshop Series. Washington DC; 2010.
41. Devereaux AV, Tosh PK, Hick JL, Hanfling D, Geiling J, Reed MJ, et al. Engagement and education: care of the critically ill and injured during pandemics and disasters: CHEST consensus statement. Chest. 2014;146(4 Suppl):e118S-33S.
42. Ratnapalan S, Martimianakis MA, Cohen-Silver JH, Minnes B, Macgregor D, Allen U, et al. Pandemic management in a pediatric hospital. Clin Pediatr (Phila). 2013;52(4):322-8.
43. Tosh PK, Feldman H, Christian MD, Devereaux AV, Kisson N, Dichter JR, et al. Business and continuity of operations: care of the critically ill and injured during pandemics and disasters: CHEST consensus statement. Chest. 2014;146(4 Suppl):e103S-17S.
44. Hota S, Fried E, Burry L, Stewart TE, Christian MD. Preparing your intensive care unit for the second wave of H1N1 and future surges. Crit Care Med. 2010;38(4 Suppl):e110-9.
45. Einav S, Hick JL, Hanfling D, Erstad BL, Toner ES, Branson RD, et al. Surge capacity logistics: care of the critically ill and injured during pandemics and disasters: CHEST consensus statement. Chest. 2014;146(4 Suppl):e17S-43S.
46. Christian MD, Devereaux AV, Dichter JR, Rubinson L, Kisson N, Task Force for Mass Critical C, et al. Introduction and executive summary: care of the critically ill and injured during pandemics and disasters: CHEST consensus statement. Chest. 2014;146(4 Suppl):8S-34S.
47. Lowe JJ, Gibbs SG, Schwedhelm SS, Nguyen J, Smith PW. Nebraska Biocontainment Unit perspective on disposal of Ebola medical waste. Am J Infect Control. 2014;42(12):1256-7.
48. World Health Organization. WHO Checklist for Influenza Pandemic Preparedness Planning. Switzerland; 2005.
49. Wise RA. The creation of emergency health care standards for catastrophic events. Acad Emerg Med. 2006;13(11):1150-2.
50. Martland A, Huffines M, Henry K. Surge Priority Planning COVID-19: Critical Care Staffing and Nursing Considerations 2020 [Available from: <https://www.chestnet.org/Guidelines-and-Resources/Resources/Surge-Priority-Planning-COVID-19-Critical-Care-Staffing-and-Nursing-Considerations>].

Clinical Management of COVID-19, v4

51. Halpern NA, Tan KS. United States Resource Availability for COVID-19: Society of Critical Care Medicine; 2020 [updated March 19, 2020. Version 2:[Available from: <https://www.sccm.org/Blog/March-2020/United-States-Resource-Availability-for-COVID-19>.
52. Wax RS, Christian MD. Practical recommendations for critical care and anesthesiology teams caring for novel coronavirus (2019-nCoV) patients. *Can J Anaesth*. 2020.
53. Tosh PK, Burry L. Essential Institutional Supply Chain Management in the Setting of COVID-19 Pandemic 2020 [Available from: <https://www.chestnet.org/Guidelines-and-Resources/Resources/Essential-Institutional-Supply-Chain-Management-in-the-Setting-of-COVID-19-Pandemic>.
54. Bilben B, Grandal L, Sovik S. National Early Warning Score (NEWS) as an emergency department predictor of disease severity and 90-day survival in the acutely dyspneic patient - a prospective observational study. *Scand J Trauma Resusc Emerg Med*. 2016;24:80.
55. Redfern OC, Smith GB, Prytherch DR, Meredith P, Inada-Kim M, Schmidt PE. A Comparison of the Quick Sequential (Sepsis-Related) Organ Failure Assessment Score and the National Early Warning Score in Non-ICU Patients With/Without Infection. *Crit Care Med*. 2018;46(12):1923-33.
56. Martinez L, Cheng W, Wang X, Ling F, Mu L, Li C, et al. A Risk Classification Model to Predict Mortality Among Laboratory-Confirmed Avian Influenza A H7N9 Patients: A Population-Based Observational Cohort Study. *J Infect Dis*. 2019;220(11):1780-9.
57. Liao X, Wang B, Kang Y. Novel coronavirus infection during the 2019-2020 epidemic: preparing intensive care units-the experience in Sichuan Province, China. *Intensive Care Med*. 2020;46(2):357-60.
58. Sprung CL, Joynt GM, Christian MD, Truog RD, Rello J, Nates JL. Adult ICU Triage During the Coronavirus Disease 2019 Pandemic: Who Will Live and Who Will Die? Recommendations to Improve Survival. *Crit Care Med*. 2020.
59. US FDA Emergency Use Authorization 2020 [Available from: <https://www.fda.gov/emergency-preparedness-and-response/mcm-legal-regulatory-and-policy-framework/emergency-use-authorization#covidinvtrodev>
60. Kim S, Kim D, Lee B. Insufficient Sensitivity of RNA Dependent RNA Polymerase Gene of SARS-CoV-2 Viral Genome as Confirmatory Test using Korean COVID-19 Cases. Preprints 2020.
61. Tran K, Cimon K, Severn M, Pessoa-Silva CL, Conly J. Aerosol generating procedures and risk of transmission of acute respiratory infections to healthcare workers: a systematic review. *PLoS One*. 2012;7(4):e35797.
62. Wang W, Xu Y, Gao R, Lu R, Han K, Wu G, et al. Detection of SARS-CoV-2 in Different Types of Clinical Specimens. *JAMA*. 2020.
63. Hanson KE, Caliendo AM, Arias CA, Englund JA, Lee MJ, Loeb M, et al. Infectious Diseases Society of America Guidelines on the Diagnosis of COVID-19 2020 [Available from: <https://www.idsociety.org/practice-guideline/covid-19-guideline-diagnostics/>.
64. Azzi L, Carcano G, Gianfagna F, Grossi P, Gasperina DD, Genoni A, et al. Saliva is a reliable tool to detect SARS-CoV-2. *J Infect*. 2020.
65. To KK, Tsang OT, Chik-Yan Yip C, Chan KH, Wu TC, Chan JMC, et al. Consistent detection of 2019 novel coronavirus in saliva. *Clin Infect Dis*. 2020.
66. To KK, Tsang OT, Leung WS, Tam AR, Wu TC, Lung DC, et al. Temporal profiles of viral load in posterior oropharyngeal saliva samples and serum antibody responses during infection by SARS-CoV-2: an observational cohort study. *Lancet Infect Dis*. 2020;20(5):565-74.
67. Williams E, Bond K, Zhang B, Putland M, Williamson DA. Saliva as a non-invasive specimen for detection of SARS-CoV-2. *J Clin Microbiol*. 2020.
68. Wyllie AL, Fournier J, Casanovas-Massana A, Campbell M, Tokuyama M, Vijayakumar P, et al. Saliva is more sensitive for SARS-CoV-2 detection in COVID-19 patients than nasopharyngeal swabs. medRxiv. 2020:2020.04.16.20067835.
69. Letko M, Marzi A, Munster V. Functional assessment of cell entry and receptor usage for SARS-CoV-2 and other lineage B betacoronaviruses. *Nat Microbiol*. 2020;5(4):562-9.
70. FDA. Coronavirus (COVID-19) Update: FDA Authorizes First Antigen Test to Help in the Rapid Detection of the Virus that Causes COVID-19 in Patients 2020 [updated May 9, 2020. Available from: <https://www.fda.gov/news-events/press-announcements/coronavirus-covid-19-update-fda-authorizes-first-antigen-test-help-rapid-detection-virus-causes>.
71. Sethuraman N, Jeremiah SS, Ryo A. Interpreting Diagnostic Tests for SARS-CoV-2. *JAMA*. 2020;323(22):2249-

- 51.
72. IDSA COVID-19 Antibody Testing Primer, Updated April 29, 2020 [Available from: <https://www.idsociety.org/globalassets/idsa/public-health/covid-19/idsa-covid-19-antibody-testing-primer.pdf>.
73. WHO. Immunity passports in the context of COVID-19 2020 [updated April 24, 2020. Available from: <https://www.who.int/news-room/commentaries/detail/immunity-passports-in-the-context-of-covid-19>.
74. CDC. Interim Guidelines for COVID-19 Antibody Testing in Clinical and Public Health Settings 2020 [Available from: <https://www.cdc.gov/coronavirus/2019-ncov/lab/resources/antibody-tests-guidelines.html>.
75. CDC. Coronavirus Disease 2019 (COVID-19) Discontinuing Transmission-Based Precautions 2020 [Available from: <https://www.cdc.gov/coronavirus/2019-ncov/hcp/disposition-hospitalized-patients.html>.
76. NIH. Coronavirus Disease 2019 (COVID-19) Treatment Guidelines 2020 [Available from: <https://covid19treatmentguidelines.nih.gov>.
77. WHO. Home care for patients with COVID-19 presenting with mild symptoms and management of their contacts: Interim Guidance, 17 Mar 2020. 2020 [Available from: [https://www.who.int/publications-detail/home-care-for-patients-with-suspected-novel-coronavirus-\(ncov\)-infection-presenting-with-mild-symptoms-and-management-of-contacts](https://www.who.int/publications-detail/home-care-for-patients-with-suspected-novel-coronavirus-(ncov)-infection-presenting-with-mild-symptoms-and-management-of-contacts).
78. CDC. Interim Guidance for Implementing Home Care of People Not Requiring Hospitalization for Coronavirus Disease 2019 (COVID-19) 2020 [Available from: <https://www.cdc.gov/coronavirus/2019-ncov/hcp/guidance-home-care.html>.
79. Xia W, Shao J, Guo Y, Peng X, Li Z, Hu D. Clinical and CT features in pediatric patients with COVID-19 infection: Different points from adults. *Pediatric Pulmonology*.n/a(n/a).
80. Metlay JP, Waterer GW, Long AC, Anzueto A, Brozek J, Crothers K, et al. Diagnosis and Treatment of Adults with Community-acquired Pneumonia. An Official Clinical Practice Guideline of the American Thoracic Society and Infectious Diseases Society of America. *Am J Respir Crit Care Med*. 2019;200(7):e45-e67.
81. Alhazzani W, Møller MH, Arabi YM, Loeb M, Gong MN, Fan E, et al. Surviving Sepsis Campaign: Guidelines on the Management of Critically Ill Adults with Coronavirus Disease 2019 (COVID-19). *Critical Care Medicine*. 2020.
82. Clinical management of severe acute respiratory infection (SARI) when COVID-19 disease is suspected: World Health Organization; 2020 [13 March 2020:[Available from: [https://www.who.int/publications-detail/clinical-management-of-severe-acute-respiratory-infection-when-novel-coronavirus-\(ncov\)-infection-is-suspected](https://www.who.int/publications-detail/clinical-management-of-severe-acute-respiratory-infection-when-novel-coronavirus-(ncov)-infection-is-suspected).
83. He HW, Liu DW. Permissive hypoxemia/conservative oxygenation strategy: Dr. Jekyll or Mr. Hyde? *J Thorac Dis*. 2016;8(5):748-50.
84. Hui DS, Chow BK, Lo T, Tsang OTY, Ko FW, Ng SS, et al. Exhaled air dispersion during high-flow nasal cannula therapy versus CPAP via different masks. *Eur Respir J*. 2019;53(4).
85. Mauri T, Spinelli E, Mariani M, Guzzardella A, Del Prete C, Carlesso E, et al. Nasal High Flow Delivered within the Helmet: A New Noninvasive Respiratory Support. *Am J Respir Crit Care Med*. 2019;199(1):115-7.
86. Patel BK, Wolfe KS, Pohlman AS, Hall JB, Kress JP. Effect of Noninvasive Ventilation Delivered by Helmet vs Face Mask on the Rate of Endotracheal Intubation in Patients With Acute Respiratory Distress Syndrome: A Randomized Clinical Trial. *JAMA*. 2016;315(22):2435-41.
87. Patel BK, Hall JB, Kress JP. Face Mask vs Helmet for Noninvasive Ventilation-Reply. *JAMA*. 2016;316(14):1497.
88. Puah SHea. ATS/APSR Joint Webinar: Global Perspectives of COVID 19. March 30, 2020 [Available from: <https://www.youtube.com/watch?v=fl-Kf3vkCwQ&feature=youtu.be>.
89. Scaravilli V, Grasselli G, Castagna L, Zanella A, Isgro S, Lucchini A, et al. Prone positioning improves oxygenation in spontaneously breathing nonintubated patients with hypoxemic acute respiratory failure: A retrospective study. *J Crit Care*. 2015;30(6):1390-4.
90. Ding L, Wang L, Ma W, He H. Efficacy and safety of early prone positioning combined with HFNC or NIV in moderate to severe ARDS: a multi-center prospective cohort study. *Crit Care*. 2020;24(1):28.
91. Zuo MZ, Huang YG, Ma WH, Xue ZG, Zhang JQ, Gong YH, et al. Expert Recommendations for Tracheal Intubation in Critically ill Patients with Novel Coronavirus Disease 2019. *Chin Med Sci J*. 2020.
92. Mohlenkamp S, Thiele H. Ventilation of COVID-19 patients in intensive care units. *Herz*. 2020.
93. Gattinoni L, Coppola S, Cressoni M, Busana M, Rossi S, Chiumello D. Covid-19 Does Not Lead to a “Typical” Acute Respiratory Distress Syndrome. *American Journal of Respiratory and Critical Care Medicine*.0(ja):null.
94. Brower RG, Matthay MA, Morris A, Schoenfeld D, Thompson BT, Wheeler A. Acute Respiratory Distress

Clinical Management of COVID-19, v4

- Syndrome Network. Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. *N Engl J Med*. 2000;342(18):1301-8.
95. Guerin C, Reignier J, Richard JC. Prone positioning in the acute respiratory distress syndrome. *N Engl J Med*. 2013;369(10):980-1.
96. Pan C, Zhang W, Du B, Qiu HB, Huang YZ. [Prone ventilation for novel coronavirus pneumonia: no time to delay]. *Zhonghua Nei Ke Za Zhi*. 2020;59(0):E007.
97. Albert RK, Hubmayr RD. The prone position eliminates compression of the lungs by the heart. *Am J Respir Crit Care Med*. 2000;161(5):1660-5.
98. Cornejo RA, Diaz JC, Tobar EA, Bruhn AR, Ramos CA, Gonzalez RA, et al. Effects of prone positioning on lung protection in patients with acute respiratory distress syndrome. *Am J Respir Crit Care Med*. 2013;188(4):440-8.
99. Nyren S, Radell P, Lindahl SG, Mure M, Petersson J, Larsson SA, et al. Lung ventilation and perfusion in prone and supine postures with reference to anesthetized and mechanically ventilated healthy volunteers. *Anesthesiology*. 2010;112(3):682-7.
100. Guerin C, Reignier J, Richard JC, Beuret P, Gacouin A, Boulain T, et al. Prone positioning in severe acute respiratory distress syndrome. *N Engl J Med*. 2013;368(23):2159-68.
101. Samanta S, Samanta S, Wig J, Baronia AK. How safe is the prone position in acute respiratory distress syndrome at late pregnancy? *Am J Emerg Med*. 2014;32(6):687 e1-3.
102. Enforcement Policy for Ventilators and Accessories and other Respiratory Devices During the Coronavirus Disease 2019 Public Health Emergency. FDA Guidance Article. March 2020 [Available from: <https://www.fda.gov/regulatory-information/search-fda-guidance-documents/enforcement-policy-ventilators-and-accessories-and-other-respiratory-devices-during-coronavirus>].
103. Consensus Statement on Multiple Patients Per Ventilator. American Association for Respiratory Care, American Society of Anesthesiologists, Anesthesia Patient Safety Foundation, American Association of Critical-Care Nurses, and American College of Chest Physicians. 3/26/2020 2020 [Available from: <https://www.sccm.org/Disaster/Joint-Statement-on-Multiple-Patients-Per-Ventilato>].
104. CDC. COVID-19 – Ventilators 2020 [Available from: <https://www.cdc.gov/coronavirus/2019-ncov/hcp/ppe-strategy/ventilators.html>].
105. Optimizing Ventilator Use during the COVID-19 Pandemic. U.S. Public Health Service Commissioned Corps. 3/31/2020 2020 [Available from: <https://www.hhs.gov/sites/default/files/optimizing-ventilator-use-during-covid19-pandemic.pdf>].
106. Bangalore S, Sharma A, Slotwiner A, Yatskar L, Harari R, Shah B, et al. ST-Segment Elevation in Patients with Covid-19 - A Case Series. *N Engl J Med*. 2020.
107. Li B, Yang J, Zhao F, Zhi L, Wang X, Liu L, et al. Prevalence and impact of cardiovascular metabolic diseases on COVID-19 in China. *Clin Res Cardiol*. 2020;109(5):531-8.
108. ESC Guidance for the Diagnosis and Management of CV Disease during the COVID-19 Pandemic. (2020, April 21) 2020 [Available from: <https://www.escardio.org/Education/COVID-19-and-Cardiology/ESC-COVID-19-Guidance>].
109. Kirkpatrick J, Mitchell C, Taub C, Kort S, Hung J, Swaminathan M. ASE Statement on Protection of Patients and Echocardiography Service Providers During the 2019 Novel Coronavirus Outbreak. Updated April 1, 2020. 2020 [Available from: <https://www.asecho.org/ase-statement-covid-19/>].
110. Thygesen K, Alpert JS, Jaffe AS, Chaitman BR, Bax JJ, Morrow DA, et al. Fourth Universal Definition of Myocardial Infarction (2018). *J Am Coll Cardiol*. 2018;72(18):2231-64.
111. Shi S, Qin M, Shen B, Cai Y, Liu T, Yang F, et al. Association of Cardiac Injury With Mortality in Hospitalized Patients With COVID-19 in Wuhan, China. *JAMA Cardiology*. 2020.
112. Clerkin KJ, Fried JA, Raikhelkar J, Sayer G, Griffin JM, Masoumi A, et al. Coronavirus Disease 2019 (COVID-19) and Cardiovascular Disease. *Circulation*.0(0).
113. Choi AD, Abbara S, Branch KR, Feuchtner GM, Ghoshhajra B, Nieman K, et al. Society of Cardiovascular Computed Tomography guidance for use of cardiac computed tomography amidst the COVID-19 pandemic Endorsed by the American College of Cardiology. *Journal of Cardiovascular Computed Tomography*.
114. Ruan Q, Yang K, Wang W, Jiang L, Song J. Clinical predictors of mortality due to COVID-19 based on an analysis of data of 150 patients from Wuhan, China. *Intensive Care Med*. 2020.
115. Hu H, Ma F, Wei X, Fang Y. Coronavirus fulminantmyocarditis treated with glucocorticoid and human immunoglobulin. *European Heart Journal*. 2020.

Clinical Management of COVID-19, v4

116. Welt FGP, Shah PB, Aronow HD, Bortnick AE, Henry TD, Sherwood MW, et al. Catheterization Laboratory Considerations During the Coronavirus (COVID-19) Pandemic: From ACC's Interventional Council and SCAI. *Journal of the American College of Cardiology*. 2020.
117. Edelson DP, Sasson C, Chan PS, Atkins DL, Aziz K, Becker LB, et al. Interim Guidance for Basic and Advanced Life Support in Adults, Children, and Neonates With Suspected or Confirmed COVID-19: From the Emergency Cardiovascular Care Committee and Get With the Guidelines((R))-Resuscitation Adult and Pediatric Task Forces of the American Heart Association in Collaboration with the American Academy of Pediatrics, American Association for Respiratory Care, American College of Emergency Physicians, The Society of Critical Care Anesthesiologists, and American Society of Anesthesiologists: Supporting Organizations: American Association of Critical Care Nurses and National EMS Physicians. *Circulation*. 2020.
118. Zheng YY, Ma YT, Zhang JY, Xie X. COVID-19 and the cardiovascular system. *Nat Rev Cardiol*. 2020.
119. Zhang P, Zhu L, Cai J, Lei F, Qin J-J, Xie J, et al. Association of Inpatient Use of Angiotensin Converting Enzyme Inhibitors and Angiotensin II Receptor Blockers with Mortality Among Patients With Hypertension Hospitalized With COVID-19. *Circulation Research*.0(0).
120. Phelan D, Kim JH, Chung EH. A Game Plan for the Resumption of Sport and Exercise After Coronavirus Disease 2019 (COVID-19) Infection. *JAMA Cardiology*. 2020.
121. Baggish AL, Drezner JA, Kim JH, Martinez MW, Prutkin JM. The resurgence of sport in the wake of COVID-19: cardiac considerations in competitive athletes. *British Journal of Sports Medicine- Blog*. 2020.
122. Dores H, Cardim N. Return to play after COVID-19: a sport cardiologist's view. *Br J Sports Med*. 2020;bjsports-2020-102482.
123. Maron BJ, Udelson JE, Bonow RO, Nishimura RA, Ackerman MJ, Estes NAM, et al. Eligibility and Disqualification Recommendations for Competitive Athletes With Cardiovascular Abnormalities: Task Force 3: Hypertrophic Cardiomyopathy, Arrhythmogenic Right Ventricular Cardiomyopathy and Other Cardiomyopathies, and Myocarditis: A Scientific Statement From the American Heart Association and American College of Cardiology. *Circulation*. 2015;132(22).
124. Kellum JA, Lameire N, Aspelin P, Barsoum RS, Burdmann EA, Goldstein SL, et al. Improving global outcomes (KDIGO) acute kidney injury work group. KDIGO clinical practice guideline for acute kidney injury. *Kidney International Supplements*. 2012;2:1-138.
125. Chan L, Chaudhary K, Saha A, Chauhan K, Vaid A, Baweja M, et al. Acute Kidney Injury in Hospitalized Patients with COVID-19. *medRxiv*. 2020:2020.05.04.20090944.
126. Mohamed MMB, Lukitsch I, Torres-Ortiz AE, Walker JB, Varghese V, Hernandez-Arroyo CF, et al. Acute Kidney Injury Associated with Coronavirus Disease 2019 in Urban New Orleans. *Kidney360*. 2020:10.34067/KID.0002652020.
127. Cheng Y, Luo R, Wang K, Zhang M, Wang Z, Dong L, et al. Kidney disease is associated with in-hospital death of patients with COVID-19. *Kidney International*. 2020.
128. Glas GJ, Van Der Sluijs KF, Schultz MJ, Hofstra JJ, Van Der Poll T, Levi M. Bronchoalveolar hemostasis in lung injury and acute respiratory distress syndrome. *J Thromb Haemost*. 2013;11(1):17-25.
129. Liu PP, Blet A, Smyth D, Li H. The Science Underlying COVID-19: Implications for the Cardiovascular System. *Circulation*. 2020.
130. Helms J, Tacquard C, Severac F, Leonard-Lorant I, Ohana M, Delabranche X, et al. High risk of thrombosis in patients with severe SARS-CoV-2 infection: a multicenter prospective cohort study. *Intensive Care Med*. 2020.
131. Llitjos J-F, Leclerc M, Chochois C, Monsallier J-M, Ramakers M, Auvray M, et al. High incidence of venous thromboembolic events in anticoagulated severe COVID-19 patients. *Journal of Thrombosis and Haemostasis*. 2020;n/a(n/a).
132. Thachil J, Tang N, Gando S, Falanga A, Cattaneo M, Levi M, et al. ISTH interim guidance on recognition and management of coagulopathy in COVID-19. *J Thromb Haemost*. 2020;18(5):1023-6.
133. Tang N, Li D, Wang X, Sun Z. Abnormal coagulation parameters are associated with poor prognosis in patients with novel coronavirus pneumonia. *J Thromb Haemost*. 2020.
134. Thachil J, Tang N, Gando S, Falanga A, Levi M, Clark C, et al. Laboratory haemostasis monitoring in COVID-19. *J Thromb Haemost*. 2020.
135. Whyte CS, Morrow GB, Mitchell JL, Chowdary P, Mutch NJ. Fibrinolytic abnormalities in acute respiratory distress syndrome (ARDS) and versatility of thrombolytic drugs to treat COVID-19. *J Thromb Haemost*. 2020.

Clinical Management of COVID-19, v4

136. Wang J, Hajizadeh N, Moore EE, McIntyre RC, Moore PK, Veress LA, et al. Tissue plasminogen activator (tPA) treatment for COVID-19 associated acute respiratory distress syndrome (ARDS): A case series. *J Thromb Haemost.* 2020.
137. Thachil J, Tang N, Gando S, Falanga A, Levi M, Clark C, et al. Type and dose of heparin in Covid-19: Reply. *J Thromb Haemost.* 2020.
138. Barrett CD, Moore HB, Yaffe MB, Moore EE. ISTH interim guidance on recognition and management of coagulopathy in COVID-19: A Comment. *J Thromb Haemost.* 2020.
139. Kollias A, Kyriakoulis KG, Dimakakos E, Poulakou G, Stergiou GS, Syrigos K. Thromboembolic risk and anticoagulant therapy in COVID-19 patients: emerging evidence and call for action. *Br J Haematol.* 2020.
140. Tang N, Bai H, Chen X, Gong J, Li D, Sun Z. Anticoagulant treatment is associated with decreased mortality in severe coronavirus disease 2019 patients with coagulopathy. *J Thromb Haemost.* 2020;18(5):1094-9.
141. Brown BL, McCullough J. Treatment for emerging viruses: Convalescent plasma and COVID-19. *Transfus Apher Sci.* 2020:102790.
142. Rajendran K, Krishnasamy N, Rangarajan J, Rathinam J, Natarajan M, Ramachandran A. Convalescent plasma transfusion for the treatment of COVID-19: Systematic review. *J Med Virol.* 2020.
143. Casadevall A, Joyner MJ, Pirofski LA. A Randomized Trial of Convalescent Plasma for COVID-19-Potentially Hopeful Signals. *JAMA.* 2020.
144. Li L, Zhang W, Hu Y, Tong X, Zheng S, Yang J, et al. Effect of Convalescent Plasma Therapy on Time to Clinical Improvement in Patients With Severe and Life-threatening COVID-19: A Randomized Clinical Trial. *JAMA.* 2020.
145. Bergman P, Lindh AU, Bjorkhem-Bergman L, Lindh JD. Vitamin D and Respiratory Tract Infections: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *PLoS One.* 2013;8(6):e65835.
146. Cannell JJ, Vieth R, Umhau JC, Holick MF, Grant WB, Madronich S, et al. Epidemic influenza and vitamin D. *Epidemiol Infect.* 2006;134(6):1129-40.
147. Holick MF. The vitamin D deficiency pandemic: Approaches for diagnosis, treatment and prevention. *Rev Endocr Metab Disord.* 2017;18(2):153-65.
148. Martineau AR, Jolliffe DA, Hooper RL, Greenberg L, Aloia JF, Bergman P, et al. Vitamin D supplementation to prevent acute respiratory tract infections: systematic review and meta-analysis of individual participant data. *BMJ.* 2017;356:i6583.
149. McCarthy MS, Elshaw EB, Szekely BM, Raju D. A Prospective Cohort Study of Vitamin D Supplementation in AD Soldiers: Preliminary Findings. *Mil Med.* 2019;184(Suppl 1):498-505.
150. Rondanelli M, Miccono A, Lamburghini S, Avanzato I, Riva A, Allegrini P, et al. Self-Care for Common Colds: The Pivotal Role of Vitamin D, Vitamin C, Zinc, and Echinacea in Three Main Immune Interactive Clusters (Physical Barriers, Innate and Adaptive Immunity) Involved during an Episode of Common Colds-Practical Advice on Dosages and on the Time to Take These Nutrients/Botanicals in order to Prevent or Treat Common Colds. *Evid Based Complement Alternat Med.* 2018;2018:5813095.
151. Dabbagh-Bazarbachi H, Clergeaud G, Quesada IM, Ortiz M, O'Sullivan CK, Fernandez-Larrea JB. Zinc ionophore activity of quercetin and epigallocatechin-gallate: from Hepa 1-6 cells to a liposome model. *J Agric Food Chem.* 2014;62(32):8085-93.
152. Prasad AS. Zinc is an Antioxidant and Anti-Inflammatory Agent: Its Role in Human Health. *Front Nutr.* 2014;1:14.
153. te Velthuis AJW, van den Worm SHE, Sims AC, Baric RS, Snijder EJ, van Hemert MJ. Zn²⁺ Inhibits Coronavirus and Arterivirus RNA Polymerase Activity In Vitro and Zinc Ionophores Block the Replication of These Viruses in Cell Culture. *PLOS Pathogens.* 2010;6(11):e1001176.
154. Xue J, Moyer A, Peng B, Wu J, Hannafon BN, Ding W-Q. Chloroquine Is a Zinc Ionophore. *PLOS ONE.* 2014;9(10):e109180.
155. Biesalski HK, Nohr D. Importance of vitamin-A for lung function and development. *Mol Aspects Med.* 2003;24(6):431-40.
156. Higdon J, Drake V, Angelo G, Delage B, Carr A, Michels A. Vitamin C. Linus Pauling Institute. Reviewed December 2018 [Available from: <https://lpi.oregonstate.edu/mic/vitamins/vitamin-C>]
<<https://lpi.oregonstate.edu/mic/vitamins/vitamin-C>.
157. Martindale R, Patel J, Taylor B, Warren M, McClave S. Nutrition Therapy in the Patient with COVID-19 Disease Requiring ICU Care. Reviewed and approved by ASPEN and Society for Critical Care Medicine. 1 April 2020

[Available from:

[https://www.nutritioncare.org/Guidelines and Clinical Resources/Resources for Clinicians Caring for Patients with Coronavirus/](https://www.nutritioncare.org/Guidelines%20and%20Clinical%20Resources/Resources%20for%20Clinicians%20Caring%20for%20Patients%20with%20Coronavirus/).

158. Rhodes A, Evans LE, Alhazzani W, Levy MM, Antonelli M, Ferrer R, et al. Surviving Sepsis Campaign: International Guidelines for Management of Sepsis and Septic Shock: 2016. *Intensive Care Med.* 2017;43(3):304-77.
159. Albright CM, Ali TN, Lopes V, Rouse DJ, Anderson BL. The Sepsis in Obstetrics Score: a model to identify risk of morbidity from sepsis in pregnancy. *Am J Obstet Gynecol.* 2014;211(1):39 e1-8.
160. Chebbo A, Tan S, Kassis C, Tamura L, Carlson RW. Maternal Sepsis and Septic Shock. *Crit Care Clin.* 2016;32(1):119-35.
161. Weiss SL, Peters MJ, Alhazzani W, Agus MSD, Flori HR, Inwald DP, et al. Surviving Sepsis Campaign International Guidelines for the Management of Septic Shock and Sepsis-Associated Organ Dysfunction in Children. *Pediatr Crit Care Med.* 2020;21(2):e52-e106.
162. Comparison of Two Fluid-Management Strategies in Acute Lung Injury. *New England Journal of Medicine.* 2006;354(24):2564-75.
163. American College of Radiology (ACR) COVID-19 Clinical Resources for Radiologists, updated 27 March 2020 [Available from: <https://www.acr.org/Clinical-Resources/COVID-19-Radiology-Resources>.
164. Davenport MS, Bruno MA, Iyer RS, Johnson AM, Herrera R, Nicola GN, et al. ACR Statement on Safe Resumption of Routine Radiology Care During the Coronavirus Disease 2019 (COVID-19) Pandemic. *J Am Coll Radiol.* 2020.
165. ACR recommendations for the use chest radiography and computed tomography (CT) for suspected COVID-19 infection, updated 22 March 2020 [Available from: <https://www.acr.org/Advocacy-and-Economics/ACR-Position-Statements/Recommendations-for-Chest-Radiography-and-CT-for-Suspected-COVID19-Infection>.
166. Mossa-Basha M, Meltzer CC, Kim DC, Tuite MJ, Kolli KP, Tan BS. Radiology Department Preparedness for COVID-19: Radiology Scientific Expert Panel. *Radiology.* 2020:200988.
167. Rubin GD, Ryerson CJ, Haramati LB, Sverzellati N, Kanne JP, Raoof S, et al. The Role of Chest Imaging in Patient Management during the COVID-19 Pandemic: A Multinational Consensus Statement from the Fleischner Society. *Radiology.* 2020:201365.
168. Ai T, Yang Z, Hou H, Zhan C, Chen C, Lv W, et al. Correlation of Chest CT and RT-PCR Testing in Coronavirus Disease 2019 (COVID-19) in China: A Report of 1014 Cases. *Radiology.* 2020:200642.
169. Bai HX, Hsieh B, Xiong Z, Halsey K, Choi JW, Tran TML, et al. Performance of radiologists in differentiating COVID-19 from viral pneumonia on chest CT. *Radiology.* 2020:200823.
170. Xie X, Zhong Z, Zhao W, Zheng C, Wang F, Liu J. Chest CT for Typical 2019-nCoV Pneumonia: Relationship to Negative RT-PCR Testing. *Radiology.* 2020:200343.
171. Oudkerk M, Buller HR, Kuijpers D, van Es N, Oudkerk SF, McCloud TC, et al. Diagnosis, Prevention, and Treatment of Thromboembolic Complications in COVID-19: Report of the National Institute for Public Health of the Netherlands. *Radiology.* 2020:201629.
172. Society of Interventional Radiology COVID-19 Planning, updated 23 March 2020 [Available from: <https://www.sirweb.org/practice-resources/toolkits/covid-19-toolkit/covid-19-planning/>.
173. Wong HYF, Lam HYS, Fong AH-T, Leung ST, Chin TW-Y, Lo CSY, et al. Frequency and Distribution of Chest Radiographic Findings in COVID-19 Positive Patients. *Radiology.* 2020:201160.
174. Chung M, Bernheim A, Mei X, Zhang N, Huang M, Zeng X, et al. CT Imaging Features of 2019 Novel Coronavirus (2019-nCoV). *Radiology.* 2020;295(1):202-7.
175. Fang Y, Zhang H, Xu Y, Xie J, Pang P, Ji W. CT Manifestations of Two Cases of 2019 Novel Coronavirus (2019-nCoV) Pneumonia. *Radiology.* 2020;295(1):208-9.
176. Ng M-Y, Lee EY, Yang J, Yang F, Li X, Wang H, et al. Imaging Profile of the COVID-19 Infection: Radiologic Findings and Literature Review. *Radiology: Cardiothoracic Imaging.* 2020;2(1):e200034.
177. Song F, Shi N, Shan F, Zhang Z, Shen J, Lu H, et al. Emerging 2019 Novel Coronavirus (2019-nCoV) Pneumonia. *Radiology.* 2020;295(1):210-7.
178. Pan F, Ye T, Sun P, Gui S, Liang B, Li L, et al. Time Course of Lung Changes On Chest CT During Recovery From 2019 Novel Coronavirus (COVID-19) Pneumonia. *Radiology.* 2020:200370.
179. Wang Y, Dong C, Hu Y, Li C, Ren Q, Zhang X, et al. Temporal Changes of CT Findings in 90 Patients with COVID-19 Pneumonia: A Longitudinal Study. *Radiology.* 2020:200843.

Clinical Management of COVID-19, v4

180. Simpson S, Kay FU, Abbara S, Bhalla S, Chung JH, Chung M, et al. Radiological Society of North America Expert Consensus Statement on Reporting Chest CT Findings Related to COVID-19. Endorsed by the Society of Thoracic Radiology, the American College of Radiology, and RSNA. *Radiology: Cardiothoracic Imaging*. 2020;2(2):e200152.
181. Beach SR, Praschan NC, Hogan C, Dotson S, Merideth F, Kontos N, et al. Delirium in COVID-19: A case series and exploration of potential mechanisms for central nervous system involvement. *Gen Hosp Psychiatry*. 2020;65:47-53.
182. Poyiadji N, Shahin G, Noujaim D, Stone M, Patel S, Griffith B. COVID-19-associated Acute Hemorrhagic Necrotizing Encephalopathy: CT and MRI Features. *Radiology*. 2020:201187.
183. Radmanesh A, Derman A, Lui YW, Raz E, Loh JP, Hagiwara M, et al. COVID-19 -associated Diffuse Leukoencephalopathy and Microhemorrhages. *Radiology*. 2020:202040.
184. Kandemirli SG, Dogan L, Sarikaya ZT, Kara S, Akinci C, Kaya D, et al. Brain MRI Findings in Patients in the Intensive Care Unit with COVID-19 Infection. *Radiology*. 2020:201697.
185. Mahammedi A, Saba L, Vagal A, Leali M, Rossi A, Gaskill M, et al. Imaging in Neurological Disease of Hospitalized COVID-19 Patients: An Italian Multicenter Retrospective Observational Study. *Radiology*. 2020:201933.
186. Bhayana R, Som A, Li MD, Carey DE, Anderson MA, Blake MA, et al. Abdominal Imaging Findings in COVID-19: Preliminary Observations. *Radiology*. 2020:201908.
187. National Academies of Sciences, Engineering, and Medicine. Integrating clinical research into epidemic response: The Ebola experience. Washington DC: The National Academies Press; 2017.
188. Lee N, Allen Chan KC, Hui DS, Ng EK, Wu A, Chiu RW, et al. Effects of early corticosteroid treatment on plasma SARS-associated Coronavirus RNA concentrations in adult patients. *J Clin Virol*. 2004;31(4):304-9.
189. Wang Y, Jiang W, He Q, Wang C, Wang B, Zhou P, et al. A retrospective cohort study of methylprednisolone therapy in severe patients with COVID-19 pneumonia. *Signal Transduct Target Ther*. 2020;5(1):57.
190. Low-cost dexamethasone reduces death by up to one third in hospitalised patients with severerespiratory complications of COVID-19 [press release]. JUNE 16, 2020 2020.
191. Mo P, Xing Y, Xiao Y, Deng L, Zhao Q, Wang H, et al. Clinical characteristics of refractory COVID-19 pneumonia in Wuhan, China. *Clin Infect Dis*. 2020.
192. Infectious Diseases Society of America Guidelines on the Treatment and Management of Patients with COVID-19: IDSA; 2020 [Available from: <https://www.idsociety.org/practice-guideline/covid-19-guideline-treatment-and-management/>]
193. de Wit E, Feldmann F, Cronin J, Jordan R, Okumura A, Thomas T, et al. Prophylactic and therapeutic remdesivir (GS-5734) treatment in the rhesus macaque model of MERS-CoV infection. *Proceedings of the National Academy of Sciences*. 2020:201922083.
194. Sheahan TP, Sims AC, Leist SR, Schafer A, Won J, Brown AJ, et al. Comparative therapeutic efficacy of remdesivir and combination lopinavir, ritonavir, and interferon beta against MERS-CoV. *Nat Commun*. 2020;11(1):222.
195. Wang M, Cao R, Zhang L, Yang X, Liu J, Xu M, et al. Remdesivir and chloroquine effectively inhibit the recently emerged novel coronavirus (2019-nCoV) in vitro. *Cell Res*. 2020;30(3):269-71.
196. Wang Y, Zhang D, Du G, Du R, Zhao J, Jin Y, et al. Remdesivir in adults with severe COVID-19: a randomised, double-blind, placebo-controlled, multicentre trial. *The Lancet*.
197. Beigel JH, Tomashek KM, Dodd LE, Mehta AK, Zingman BS, Kalil AC, et al. Remdesivir for the Treatment of Covid-19 - Preliminary Report. *N Engl J Med*. 2020.
198. Goldman JD, Lye DCB, Hui DS, Marks KM, Bruno R, Montejano R, et al. Remdesivir for 5 or 10 Days in Patients with Severe Covid-19. *New England Journal of Medicine*. 2020.
199. Gautret P, Lagier J-C, Parola P, Hoang VT, Meddeb L, Mailhe M, et al. Hydroxychloroquine and azithromycin as a treatment of COVID-19: results of an open-label non-randomized clinical trial. *International Journal of Antimicrobial Agents*. 2020.
200. Molina JM, Delaugerre C, Le Goff J, Mela-Lima B, Ponscarne D, Goldwirt L, et al. No evidence of rapid antiviral clearance or clinical benefit with the combination of hydroxychloroquine and azithromycin in patients with severe COVID-19 infection. *Médecine et Maladies Infectieuses*. 2020.
201. Cao B, Wang Y, Wen D, Liu W, Wang J, Fan G, et al. A Trial of Lopinavir-Ritonavir in Adults Hospitalized with

Clinical Management of COVID-19, v4

Severe Covid-19. N Engl J Med. 2020.

202. Liu F, Xu A, Zhang Y, Xuan W, Yan T, Pan K, et al. Patients of COVID-19 may benefit from sustained lopinavir-combined regimen and the increase of eosinophil may predict the outcome of COVID-19 progression. Int J Infect Dis. 2020.
203. Yao TT, Qian JD, Zhu WY, Wang Y, Wang GQ. A systematic review of lopinavir therapy for SARS coronavirus and MERS coronavirus-A possible reference for coronavirus disease-19 treatment option. J Med Virol. 2020.
204. Fang L, Karakiulakis G, Roth M. Are patients with hypertension and diabetes mellitus at increased risk for COVID-19 infection? Lancet Respir Med. 2020.
205. Xu X, Han M, Li T, Sun W, Wang D, Fu B, et al. Effective Treatment of Severe COVID-19 Patients with Tocilizumab. 2020.
206. Tiberghien P, de Lambalerie X, Morel P, Gallian P, Lacombe K, Yazdanpanah Y. Collecting and evaluating convalescent plasma for COVID-19 treatment: why and how. Vox Sang. 2020.
207. Shen C, Wang Z, Zhao F, Yang Y, Li J, Yuan J, et al. Treatment of 5 Critically Ill Patients With COVID-19 With Convalescent Plasma. JAMA. 2020.
208. Liu STH, Lin H-M, Baine I, Wajnberg A, Gumprecht JP, Rahman F, et al. Convalescent plasma treatment of severe COVID-19: A matched control study. medRxiv. 2020:2020.05.20.20102236.
209. Tanne JH. Covid-19: FDA approves use of convalescent plasma to treat critically ill patients. BMJ. 2020;368:m1256.
210. Boelig RC, Manuck T, Oliver EA, Di Mascio D, Saccone G, Bellussi F, et al. Labor and Delivery Guidance for COVID-19. Am J Obstet Gynecol MFM. 2020:100110.
211. Rasmussen SA, Smulian JC, Lednicky JA, Wen TS, Jamieson DJ. Coronavirus Disease 2019 (COVID-19) and Pregnancy: What obstetricians need to know. Am J Obstet Gynecol. 2020.
212. ReproTox [Available from: www.reprotox.org.
213. Kreuziger LB, Lee A, Garcia D, Cuker A, Cushman M, DeSancho M, et al. COVID-19 and VTE/Anticoagulation: Frequently Asked Questions (Version 3.0; last updated May 18, 2020) 2020 [Available from: <https://www.hematology.org/covid-19/covid-19-and-vte-anticoagulation>.
214. Oxford CM, Ludmir J. Trauma in pregnancy. Clinical obstetrics and gynecology. 2009;52(4):611-29.
215. Jeejeebhoy FM, Zelop CM, Lipman S, Carvalho B, Joglar J, Mhyre JM, et al. Cardiac Arrest in Pregnancy: A Scientific Statement From the American Heart Association. Circulation. 2015;132(18):1747-73.
216. Fan C, Lei D, Fang C, Li C, Wang M, Liu Y, et al. Perinatal Transmission of COVID-19 Associated SARS-CoV-2: Should We Worry? Clin Infect Dis. 2020.
217. CDC. Coronavirus Disease 2019 (COVID-19) Caring for Newborns (Updated May 20, 2020) 2020 [Available from: <https://www.cdc.gov/coronavirus/2019-ncov/hcp/caring-for-newborns.html>.
218. Center for Disease Control and Prevention. Pregnancy & Breastfeeding: Information about Coronavirus Disease 2019 2020.
219. CDC. Interim Considerations for Infection Prevention and Control of Coronavirus Disease 2019 (COVID-19) in Inpatient Obstetric Healthcare Settings 2020 [Available from: <https://www.cdc.gov/coronavirus/2019-ncov/hcp/inpatient-obstetric-healthcare-guidance.html>.
220. COVIDView 2020 [updated 17 April 2020. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/covid-data/covidview/index.html>.
221. Paret M, Lighter J, Pellett Madan R, Raabe VN, Shust GF, Ratner AJ. SARS-CoV-2 infection (COVID-19) in febrile infants without respiratory distress. Clinical Infectious Diseases. 2020.
222. Patek P, Corcoran J, Adams L, Khandhar P. SARS-CoV-2 Infection in a 2-Week-Old Male With Neutropenia. Clin Pediatr (Phila). 2020:9922820920014.
223. CDC COVID-19 Response Team. Coronavirus Disease 2019 in Children - United States, February 12-April 2, 2020. MMWR Morb Mortal Wkly Report. 2020 Apr 10;69(14):422-426. 2020.
224. CDC. COVID-NET: A Weekly Summary of U.S. COVID-19 Hospitalizations 2020 [Available from: https://gis.cdc.gov/grasp/covidnet/COVID19_5.html.
225. National Vital Statistics System Provisional Death Counts for Coronavirus Disease (COVID-19). 2020 [Available from: <https://www.cdc.gov/nchs/nvss/vsrr/COVID19/>.
226. European Center for Disease Prevention and Control's Situation dashboard- COVID-19 cases in Europe and worldwide. 2020 [Available from: <https://gap.ecdc.europa.eu/public/extensions/COVID-19/COVID-19.html>.

Clinical Management of COVID-19, v4

227. Kim D, Quinn J, Pinsky B, Shah NH, Brown I. Rates of Co-infection Between SARS-CoV-2 and Other Respiratory Pathogens. *Jama*. 2020;323(20):2085-6.
228. Chiotos K, Hayes M, Kimberlin D, Jones S, James S, Pinninti S, et al. Multicenter initial guidance on use of antivirals for children with COVID-19/SARS-CoV-2. *Journal of the Pediatric Infectious Diseases Society*. 2020.
229. Lu Y, Li Y, Deng W, Liu M, He Y, Huang L, et al. Symptomatic Infection is Associated with Prolonged Duration of Viral Shedding in Mild Coronavirus Disease 2019: A Retrospective Study of 110 Children in Wuhan. *The Pediatric infectious disease journal*. 2020.
230. Verdoni L, Mazza A, Gervasoni A, Martelli L, Ruggeri M, Ciuffreda M, et al. An outbreak of severe Kawasaki-like disease at the Italian epicentre of the SARS-CoV-2 epidemic: an observational cohort study. *Lancet*. 2020;395(10239):1771-8.
231. Licciardi F, Pruccoli G, Denina M, Parodi E, Taglietto M, Rosati S, et al. SARS-CoV-2-Induced Kawasaki-Like Hyperinflammatory Syndrome: A Novel COVID Phenotype in Children. *Pediatrics*. 2020.
232. Rivera-Figueroa EI, Santos R, Simpson S, Garg P. Incomplete Kawasaki Disease in a Child with Covid-19. *Indian pediatrics*. 2020.
233. Balasubramanian S, Nagendran TM, Ramachandran B, Ramanan AV. Hyper-inflammatory Syndrome in a Child With COVID-19 Treated Successfully With Intravenous Immunoglobulin and Tocilizumab. *Indian pediatrics*. 2020.
234. CDC. Clinical Outreach and Communication Activity (COCA): Multisystem Inflammatory Syndrome in Children (MIS-C) Associated with Coronavirus Disease 2019 (COVID-19); May 19, 2020 [Available from: https://emergency.cdc.gov/coca/calls/2020/callinfo_051920.asp?deliveryName=USCDC_1052%29-DM28626].
235. Mahase E. Covid-19: concerns grow over inflammatory syndrome emerging in children. *Bmj*. 2020;369:m1710.
236. Riphagen S, Gomez X, Gonzalez-Martinez C, Wilkinson N, Theocharis P. Hyperinflammatory shock in children during COVID-19 pandemic. *Lancet*. 2020;395(10237):1607-8.
237. CDC. COVID-19 Older Adults 2020 [Available from: <https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/older-adults.html>].
238. D'Adamo H, Yoshikawa T, Ouslander JG. Coronavirus Disease 2019 in Geriatrics and Long-term Care: The ABCDs of COVID-19. *J Am Geriatr Soc*. 2020.
239. Han JH, Wilson A, Vasilevskis EE, Shintani A, Schnelle JF, Dittus RS, et al. Diagnosing delirium in older emergency department patients: validity and reliability of the delirium triage screen and the brief confusion assessment method. *Ann Emerg Med*. 2013;62(5):457-65.
240. Inouye SK, van Dyck CH, Alessi CA, Balkin S, Siegal AP, Horwitz RI. Clarifying confusion: the confusion assessment method. A new method for detection of delirium. *Ann Intern Med*. 1990;113(12):941-8.
241. Lindroth H, Bratzke L, Purvis S, Brown R, Coburn M, Mrkobrada M, et al. Systematic review of prediction models for delirium in the older adult inpatient. *BMJ Open*. 2018;8(4):e019223.
242. Statement from the British Geriatrics Society. Managing delirium in confirmed and suspected cases 2020 [Available from: <https://www.bgs.org.uk/resources/coronavirus-managing-delirium-in-confirmed-and-suspected-cases>].
243. Hwang U, Malsch AJ, Biese KJ, Inouye SK. Preventing and Managing Delirium in Older Emergency Department Patients During the COVID-19 Pandemic. *Journal of Geriatric Emergency Medicine*. 2020;1(4):Supplement 1.
244. Neufeld KJ, Yue J, Robinson TN, Inouye SK, Needham DM. Antipsychotic Medication for Prevention and Treatment of Delirium in Hospitalized Adults: A Systematic Review and Meta-Analysis. *J Am Geriatr Soc*. 2016;64(4):705-14.
245. Farah A. Atypicality of atypical antipsychotics. *Prim Care Companion J Clin Psychiatry*. 2005;7(6):268-74.
246. Clinical Issues and Guidance From the American College of Surgeons 2020 [Available from: <https://www.facs.org/covid-19/clinical-guidance>].
247. COVID-19: Guidance for Triage of Non-Emergent Surgical Procedures 2020 [Available from: <https://www.facs.org/covid-19/clinical-guidance/triage>].
248. Create a Surgical Review Committee for COVID-19-Related Surgical Triage Decision Making 2020 [Available from: <https://www.facs.org/covid-19/clinical-guidance/review-committee>].
249. Prachand VN, Milner R, Angelos P, Posner MC, Fung JJ, Agrawal N, et al. Medically Necessary, Time-Sensitive *Guideline Only/Not a Substitute for Clinical Judgment*


Clinical Management of COVID-19, v4

Procedures: Scoring System to Ethically and Efficiently Manage Resource Scarcity and Provider Risk During the COVID-19 Pandemic. J Am Coll Surg. 2020.

250. Joint Statement: Roadmap for Resuming Elective Surgery after COVID-19 Pandemic 2020 [Available from: <https://www.facs.org/covid-19/clinical-guidance/roadmap-elective-surgery>.
251. ACS. American College of Surgeons Post-COVID-19 Readiness Checklist for Resuming Surgery 2020 [Available from: <https://www.facs.org/covid-19/checklist>.
252. Guidelines: Opening Up American Again 2020 [Available from: <https://www.whitehouse.gov/openingamerica/#criteria>.
253. The ASA and APSF Joint Statement on Perioperative Testing for the COVID-19 Virus 2020 [Available from: <https://www.asahq.org/about-asa/newsroom/news-releases/2020/04/asa-and-apsf-joint-statement-on-perioperative-testing-for-the-covid-19-virus>.
254. American College of Surgeons. COVID-19: Considerations for Optimum Surgeon Protection Before, During, and After Operation 2020 [Available from: <https://www.facs.org/covid-19/clinical-guidance/surgeon-protection>.
255. APSF. Perioperative Considerations for the 2019 Novel Coronavirus (COVID-19) 2020 [Available from: <https://www.apsf.org/news-updates/perioperative-considerations-for-the-2019-novel-coronavirus-covid-19/>.
256. SAGES and EAES Recommendations Regarding Surgical Response to COVID-19 Crisis 2020 [Available from: <https://www.sages.org/recommendations-surgical-response-covid-19/>.
257. CDC. Guidelines for Environmental Infection Control in Health-Care Facilities (2003) 2020 [Available from: <https://www.cdc.gov/infectioncontrol/guidelines/environmental/appendix/air.html>.
258. Joint AGA/DHPA Guidance: Recommendations for Resumption of Elective Endoscopy During the COVID-19 Pandemic 2020 [updated April 27, 2020. Available from: <https://www.dhpassociation.org/2020/04/27/aga-dhpa-resume-endoscopy-covid19/>.
259. ASGE. ASGE COVID-19 Frequently Asked Questions. American Society for Gastrointestinal Endoscopy 2020 [updated March 31, 2020. Available from: https://www.asge.org/docs/default-source/default-document-library/faq_asge_covid19_updated3-31-2020_final.pdf?sfvrsn=dcb28c52_2&zs=oewhb&zl=nLI32.
260. JTS. Management of COVID-19 in Austere Operational Environments v2.0 Clinical Practice Guideline 2020 [Available from: https://jts.amedd.army.mil/index.cfm/PI_CPGs/cpgs.
261. Sher Y, Rabkin B, Maldonado JR, Mohabir P. A CASE REPORT OF COVID-19 ASSOCIATED HYPERACTIVE ICU DELIRIUM WITH PROPOSED PATHOPHYSIOLOGY AND TREATMENT. Psychosomatics. 2020.
262. LaHue SC, James TC, Newman JC, Esmaili AM, Ormseth CH, Ely EW. Collaborative Delirium Prevention in the Age of COVID-19. J Am Geriatr Soc. 2020;68(5):947-9.
263. Alosaimi FD, Alghamdi A, Alsuhaibani R, Alhammad G, Albatili A, Albatly L, et al. Validation of the Stanford Proxy Test for Delirium (S-PTD) among critical and noncritical patients. J Psychosom Res. 2018;114:8-14.
264. Ferrando SJ, Klepacz L, Lynch S, Tavakkoli M, Dornbush R, Baharani R, et al. COVID-19 Psychosis: A potential new neuropsychiatric condition triggered by novel coronavirus infection and the inflammatory response? Psychosomatics. 2020.
265. Bilbul M, Paparone P, Kim AM, Mutalik S, Ernst CL. Psychopharmacology of COVID-19. Psychosomatics. 2020;10.1016/j.psych.2020.05.006.

APPENDIX A : CRISIS LEVEL SURGE – CRITICAL CARE TRIAGE TOOL EXAMPLE

Crisis Level Surge – Critical Care Triage Tool



THEN

- Tie-breaking- A. Allocation by incremental ICU benefit- saving the most life-years
B. If still tie- first come, first served
- Re-assess priority every 24h for patients waiting for ICU admission.
- Re-assess patients at day 10-14 or earlier if significant deterioration.

APPENDIX B : INFECTION PREVENTION AND CONTROL RELEVANT RESOURCES AND DOCUMENTS

Mask Guidance
(For Use in Crisis Standards Only)

SURGICAL MASKS

DISCARD MASK IF:

- Contaminated with blood, respiratory or nasal secretions, or other bodily fluids from patients.
- Obviously damaged or hard to breathe through.
- At the conclusion of your shift.

EXTENDED USE:

- Wear mask for ENTIRE shift unless soiled, damaged, or hard to breathe through.
- Do not touch the mask. If you touch or adjust your mask, you must immediately perform proper hand hygiene.
- Leave the patient care area if you need to remove your mask.
- Consider use of a face shield over mask.

REUSE:

- Masks that fasten via ties that are unable to be undone and are torn need to be discarded.
- Masks should be carefully folded so the outer surface is held inward and against itself to reduce contact with the outer surface during storage.
- Keep used masks in a clean, breathable container such as a paper bag between uses. Do not store in a plastic bag. Keep in a clean space outside patient room, such as a wall locker next to patient room or top of the isolation cart. To prevent accidental use of another’s mask, label the container:
 - First initial and last name of owner
 - Strap of mask with first initial and last name of owner

Glossary

Extended Use — The practice of wearing the same mask/respirator for repeated close contact encounters with several patients, without removing the mask/respirator between patient encounters.

Reuse — The practice of using the same mask/respirator for multiple encounters with several patients but removing it after each encounter.

N95 RESPIRATORS

Extended and limited reuse of respirators were recommended for conserving respirators during previous respiratory pathogen outbreaks and pandemics.

Use face shield over N95 respirator to reduce surface contamination.

Perform hand hygiene with soap and water or an alcohol-based hand sanitizer before and after touching or adjusting respirator.

DISCARD N95 RESPIRATOR IF:

- Used for aerosol-generating procedure.
- Contaminated with blood, respiratory or nasal secretions, or other bodily fluids from patients.
- Obviously damaged or hard to breathe through.
- Reused (donned/doffed) a maximum of five times.

EXTENDED USE:

Extended use may be implemented when multiple patients are infected with the same respiratory pathogen and patients are placed together in dedicated waiting rooms or hospital wards.

REUSE:

- Keep used respirators in a clean, breathable container such as a paper bag between uses. Do not store in a plastic bag. Keep in a clean space outside patient room such as a wall locker near patient’s room or top of the isolation cart. To prevent accidental use of another person’s respirator, label the container:
 - First initial and last name of owner
 - Strap of respirator with first initial and last name of owner
- Avoid touching the inside of the respirator. If inadvertent contact with the inside of the respirator, perform hand hygiene as described above.
- Use a pair of clean (non-sterile) gloves when donning a used N95 respirator and performing a user seal check. Discard gloves after the N95 respirator is donned and any adjustments are made to ensure the respirator is sitting comfortably on your face with a good seal.

Standard Precautions **FOR THE CARE OF ALL PATIENTS**

Includes Blood, Body Fluids, Secretions, Excretions, and Contaminated Items

<p>Wash hands BEFORE and AFTER patient care regardless of whether gloves are worn. - Wash hands immediately after gloves are removed and between patient contacts.</p>
<p>Wear gloves when touching blood, body fluids, secretions, excretions, and contaminated items. - Put on clean gloves just before touching mucous membranes and non-intact skin.</p>
<p>Wear mask and eye protection or a face shield to protect mucous membranes of the eyes, nose, and mouth during procedures and patient care activities that are likely to generate splashes or sprays of blood/body fluids.</p>
<p>Wear gown to protect skin and prevent soiling of clothing during procedures and patient care activities that are likely to generate splashes or sprays of blood & body fluids. Remove soiled gown as promptly as possible and wash hands.</p>
<p>Take care to prevent injuries when using needles, scalpels, and other sharp instruments or devices; when handling sharp instruments after procedures; when cleaning used instruments; and when disposing of used needles.</p>
<p>Use mouthpieces, resuscitation bags, or other ventilation devices as an alternative to mouth-to-mouth resuscitation.</p>
<p>Cover your cough and sneeze with tissues or cough and sneeze into your sleeve.</p>
<p>Avoid touching your face (eyes, nose and mouth) with unclean hands.</p>
<p>Clean and disinfect shared patient equipment.</p>
<p>Use aseptic technique.</p>

POWERED AIR PURIFYING RESPIRATORS (PAPRs) April 2020 NOTE: This document was adapted from UNMC - Nebraska Medicine COVID-19 PPE Guidance for Extended Use and Limited Reuse of Disposable Facemasks, Respirators and Protective Eyewear

Introduction: PAPRs are reusable respirators that are loose-fitting hoods or helmets. Caution should be applied with use of PAPRs in surgical settings due to concerns that the blower exhaust and exhaled air may contaminate the sterile field. The FDA issued an update Mar 2020 to address NIOSH-approved air purifying respirators for use in health care settings during the COVID-19 emergency available for review at the following link:

<https://www.fda.gov/media/135763/download>. Facilities using elastomeric respirators and PAPRs are required to have up-to-date cleaning and disinfection procedures to facilitate protection against infectious agents.

Recommendations: This document provides an overview of current industry recommendations for consideration. Such recommendations are not all-inclusive, and decision-making must address the unique readiness challenges and concerns faced at each individual facility.

- Staff are required to receive training on correct use of PAPRs.
 - Training ensures HCPs are knowledgeable and proficient in the donning and doffing of PAPR and other PPE prior to engaging in patient care. In addition, during practice, HCPs and their trainers will assess their proficiency and comfort with performing required duties while wearing PAPR and other PPE.
- A trained observer is required.
 - The observer should be a dedicated and knowledgeable individual with the responsibility of ensuring adherence to the entire donning and doffing process, including disposal of used PPE. The sequence and actions involved in each donning and doffing step are critical, therefore a trained observer must read aloud to the HCP each step in the procedure checklist and visually confirm, document that the step has been completed correctly, and provide immediate corrective instruction if the HCP is not following the recommended steps.
- The following supplies are gathered in preparation for PAPR use:
 - One pair of extended cuff gloves (two pairs if practicing double gloving technique)
 - One long-sleeve gown
 - One PAPR*
 - One PAPR hood
 - One airflow indicator

*Note: The PAPR must be inspected and a function check completed in accordance with the manufacturer’s instructions for use. DO NOT USE and remove from service if airflow does not reach six cubic feet/minute (CFM). Change the filters and repeat the function test. If after changing filter the function test fails, take out of service.

- PPE must remain in place and worn correctly for the duration of exposure to potentially contaminated areas. Avoid adjusting PPE during patient care. If PAPR malfunctions during patient care, the HCP must move immediately to the doffing area to assess the situation.

Donning PAPR Equipment:

- Healthcare facilities that decide to add additional PPE or modify this PPE guidance, must consider the risk versus benefit of any modification, and train HCPs on modified donning and doffing procedures.
- The practice of double-gloving provides an extra layer of safety during direct patient care and during the PPE removal process, however more than two pairs of gloves can make it more difficult to perform patient care duties.
- PAPR and all other PPE must be donned correctly in proper order before entry into the patient care area. Donning activities must be directly observed by a trained observer. The following steps for donning must be followed:
 1. Perform hand hygiene
 2. Don PAPR
 - a. Don PAPR belt with assistance
 - b. Position PAPR around waist
 - c. Fold/tuck extra belt webbing into belt
 - d. Test range of motion
 - e. Power ON PAPR motor

Clinical Management of COVID-19, v4

3. Don PAPR hood assembly
 - a. Place hood on head. Ensure hood fits comfortably and is positioned properly
4. Don surgical gown & secure gown over the hood shroud and hose (if possible), secure both neck & waist ties
5. Don extended cuff gloves over gown wrist cuff (if desired, may use second pair of gloves)
6. Check range of motion
7. Donning partner will inspect member for defects in PPE. Pay close attention to gown/glove junction

Doffing PAPR Equipment:

- Appropriate PAPR doffing procedures must be followed. All PPE must be removed slowly and deliberately in the correct sequence. Anytime a PAPR is used, a process checklist with a designated trained observer is required.
- The following steps must be followed for doffing:
 1. Doffing will begin in the patient's room. Doffing partner will be prepared to assist outside patient's room by performing hand hygiene and donning the surgical mask and gloves. Doffing partner will prepare the area outside the room, and gather the following supplies:
 - a. Intravenous (IV) Pole
 - b. Disinfectant wipes
 - c. Biohazard bags
 - d. Plastic bag
 2. HCP performs hand hygiene over gloves
 3. Gown is removed by pulling away from the shoulders, taking care to avoid jerking motion; may remove gloves in conjunction with the gown (if using the double-gloving technique, remove outer pair of gloves prior to removing gown)
 4. If gloves are still on, remove gloves using the "glove in glove" technique
 5. Perform hand hygiene
 6. HCP will leave the patient's room
 7. Keeping the blower motor ON, HCP will disconnect belt, and hand it to the doffing partner
 8. Doffing partner will hang belt on the IV pole
 9. HCP completes hand hygiene
 10. Doffing partner thoroughly disinfects PAPR hose and motor using approved disinfectant wipe
 11. Doffing Partner will tell HCP that the hose will be disconnected from PAPR motor
 12. Doffing partner will hold the hose and instruct HCP to lean forward and remove the hood
 - a. HCP will reach under the sides of the hood and carefully remove the hood over and off head
 - b. Alternative method: HCP will pinch the crown of the hood and carefully pull the hood over and off head
 13. Doffing partner will place the hood and hose into plastic bag. *Note: the hood may be reused if supplies are low
 14. HCP will complete hand hygiene and exit the area
 15. Doffing partner will perform hand hygiene
- Appropriate steps for doffing area cleanup must be performed as follows by doffing partner:
 1. Dons new pair of gloves
 2. Disinfects high-touch surfaces
 3. Disinfects the IV pole
 4. Place PAPR in biohazard bag and stores in designated area
 5. Remove regulated medical waste (RMW) bags from waste receptacles
 - a. Secure bags with tape
 - b. Do not express any trapped air from the bag
 - c. Place bags in the designated area/soiled utility room
 - d. Perform hand hygiene
 - e. Replace red bag
 - f. Perform hand hygiene

Clinical Management of COVID-19, v4

- Steps for disinfection and storage of PAPR components including hood for re-use:
 1. Perform hand hygiene
 2. Don gloves and a procedure mask, and carry the PAPR to the PAPR processing area without allowing it to come in contact with clothing or skin
 3. Visually inspect the PAPR hood for contamination; discard and do not re-use if visibly contaminated
 - a. If visible contamination is not observed and PAPR will be reused during the shift, do not disconnect any of the PAPR components
 - b. Do not remove the PAPR filters from the motor unless flow test fails
 4. Disinfect the PAPR motor, belt, hose and hood using Environmental Protection Agency (EPA) approved disinfectant wipes, while observing contact time
 5. Disinfect in the following order (using a new wipe for each component):
 - a. PAPR motor and filters (avoid introducing liquid into the filter holes)
 - b. Belt
 - c. Tubing sleeve
 - d. Hood (wipe the hood inside, then the outside)
 6. Once completely dry, place the PAPR in a clean area
 7. Ensure battery is charged or place on charger in accordance with the manufacturer instructions for use (IFU)
- Steps for terminal disinfection and storage of used PAPR components:
 1. Follow the above procedure for cleaning and disinfecting PAPR with the following additional steps:
 - a. Disconnect PAPR belt to disinfect separately and reattach to PAPR motor when dry
 - b. Disconnect and dispose of PAPR hood
 - c. Return PAPR motor with filters, belt, and tubing attached to unit storage area
 - d. Plug in PAPR motor to recharge battery in accordance with manufacturer IFU

References

1. Food and Drug Administration. (2020). NIOSH-Approved Air Purifying Respirators for Use in Health Care Settings during Response to the COVID-19 Public Health Emergency. <https://www.fda.gov/media/135763/download>
2. Strategies for Optimizing the Supply of N95 Respirators 2 Apr 2020 Personal Protective Equipment: Respiratory Protection: Use of alternatives to N95 respirators <https://www.cdc.gov/coronavirus/2019-ncov/hcp/respirators-strategy/index.html>
3. Guidance on Personal Protective Equipment (PPE) To Be Used By Healthcare Workers during Management of Patients with Confirmed Ebola or Persons under Investigation (PUIs) for Ebola who are Clinically Unstable or Have Bleeding, Vomiting, or Diarrhea in U.S. Hospitals, Including Procedures for Donning and Doffing PPE 30 August 2018 <https://www.cdc.gov/vhf/ebola/healthcare-us/ppe/guidance.html>
4. Verbeek JH, Rajamaki B, Ijaz S, Tikka C, Ruotsalainen JH, Edmond MB, Sauni R, Kilinc Balci FS. Personal protective equipment for preventing highly infectious diseases due to exposure to contaminated body fluids in healthcare staff. *Cochrane Database Syst Rev.* 2019 Jul 1;7:CD011621. doi: 10.1002/14651858.CD011621.pub3. PubMed [citation] PMID: 31259389, PMCID: PMC6601138
5. Chughtai AA, Stelzer-Braid S, Rawlinson W, Pontivivo G, Wang Q, Pan Y, Zhang D, Zhang Y, Li L, MacIntyre CR. Contamination by respiratory viruses on outer surface of medical masks used by hospital healthcare workers. *BMC Infect Dis.* 2019 Jun 3;19(1):491. doi: 10.1186/s12879-019-4109-x. PubMed [citation] PMID: 31159777, PMCID: PMC6547584

DECONTAMINATION OF N95 RESPIRATORS: June 2020

Situation and Background

COVID-19 has caused significant disruption in the manufacturing of N95 filtering facemask respirators (FFRs), subsequently generating a need for strategies to decontaminate for reuse. To ensure existing resources are leveraged effectively, and Military Medical Treatment Facilities (MTFs) are equipped to optimally care for patients in a crisis situation, an evaluation of alternative strategies is warranted.

Disposable FFRs, are not approved for routine decontamination as conventional standards of care. Per the CDC guidance for crisis standards these FFRs should not be worn by healthcare providers when performing or present for an aerosol-generating procedures. Refer to table on Mask Guidance in this document.

Assessment

There are currently four strategies for decontamination of N95 FFRs. These include high-concentration hydrogen peroxide (e.g., the Battelle Decontamination System), hydrogen peroxide sterilization systems (STERRAD, STERIZONE, STERIS, and Steriluent), heat and humidity, and ultraviolet (UV) decontamination (e.g., Xenex). The implementation of each of these strategies carries with it unique benefits and challenges, as highlighted below:

- On 28 MAR 2020, the Food and Drug Administration (FDA) issued an Emergency Use Authorization (EUA) for the Battelle Decontamination System.¹ This system can decontaminate thousands of N95 FFRs at one time with up to 20 decontamination cycles per mask. Additionally, FDA does not require masks to be returned to the same user. A unique challenge associated with this product is that implementation requires transport of the respirators to and from the decontamination site, and therefore requires logistics support.
- FDA has now issued Emergency use Authorization to four VHP sterilizer companies.²⁻⁵ These authorizations are only for specific models and do not include all systems that various companies make. A benefit associated with these devices, is that many MTFs already have them, but the disadvantage is the decontamination capacity of each system is anywhere from 10-20 masks every 30-60 minutes depending on the system. Another disadvantage is that FDA requires single-user per masks and STERRAD and STERIZONE are only recommended for two decontamination cycles, while STERIS and Steriluent are authorized for up to ten decontamination cycles.
- Heat and Humidity has been proposed as an option for decontamination of N95 FFRs. This method has not been authorized by FDA. Various studies have demonstrated that under the right circumstances of about 70-85 °C with relative humidity of 50-85% for 60 minutes, inactivation of SARS-CoV-2 is likely. Users are cautioned that if humidity is not maintained, viable viruses were present on the mask. The advantage of this strategy is that systems that can achieve these parameters are inexpensive and widely available, and most N95 FFRs have maintained good fit and filter performance for up to 3 cycles. The disadvantage is that mask integrity as it relates to decontamination cycles varies based on the make and model of N95 FFR, and therefore facilities must assure that the masks that are being decontaminated were studied within those parameters.
- The use of UV-C disinfection is now gaining recognition in the literature as a potentially viable strategy for N95 FFR decontamination during this crisis⁶, and a number of reputable hospital systems have publicly supported the practice. ECRI has provided communications indicating this approach is acceptable as a last resort, and additional information regarding use of this method is available at the following link: <https://www.nebraskamed.com/sites/default/files/documents/covid-19/n-95-decon-process.pdf> Currently some MTFs have developed protocols and may already be implementing this decontamination strategy. The advantages of using this process is that many masks could be decontaminated in fairly short period of time, but numerous disadvantages should be considered when implementing this strategy:
 - The UV-C light systems are not regulated as medical devices by FDA, and therefore must be validated for appropriate output.
 - UV-C light must shine directly on all surfaces, which is difficult to accomplish with curved masks (any shadows may leave masks still contaminated).

Clinical Management of COVID-19, v4

- UV-C light must be delivered at proper dose. This should be verified by a UV-C-specific sensor.
- UV light degrades mask components, and determining the number of decontamination cycles depends on the amount of UV light delivered per cycle.
- It is likely that due to kinking, straps would not receive proper amount of UV-C light. Experts recommend that decontamination of straps is conducted manually.

The following considerations must be taken into account if using any decontamination strategy:

- Decontaminated compatible N95 FFRs are not sterile, and in most cases (with exception of Battelle's method) must go back to the original wearer.
- All hydrogen peroxide systems cannot decontaminate masks that contain cellulose-based materials
- Each of the aforementioned systems have different requirements regarding the number of times they can be used.
- If any of the N95 FFRs are visibly soiled (e.g., blood, dried sputum, makeup, body fluids) they must be disposed of.
- If a good seal cannot be maintained, the mask must be disposed of.
- Any individual handling contaminated respirators must wear full personal protective equipment (PPE), including an N95 FFR and eye protection.

Recommendation

Leadership should consider hydrogen peroxide sterilization systems as a first line decontamination strategy, and avoid the use of UV light and heat and humidity strategies for decontamination until proper validation of effectiveness is achieved. If MTFs have concerns regarding an inability to maintain adequate supply of N95 respirators, the DHA IPC Tiger Team should be contacted at the following e-mail to address such concerns: dha.ncr.clinic-support.list.ipc-group@mail.mil

References

1. Food and Drug Administration. (2020). *Battelle Decontamination System Letter of Authorization*. <https://www.fda.gov/media/136529/download>
2. Food and Drug Administration. (2020). *STERIS Sterilization Systems Letter of Authorization*. <https://www.fda.gov/media/136843/download>
3. Food and Drug Administration. (2020). *STERRAD Sterilization Systems Letter of Authorization*. <https://www.fda.gov/media/136884/download>
4. Food and Drug Administration. (2020). *Sterilucent Letter of Authorization*. <https://www.fda.gov/media/137167/download>
5. Stryker. (2020). *Emergency Use Decontamination of Compatible N95 or N95-equivalent Respirators in Stryker's STERIZONE VP4 Sterilizer Using the Stryker STERIZONE VP4 N95 Respirator Decontamination Cycle*. <https://www.fda.gov/media/136979/download>
6. Card, K., Crozier, D., Dhawan, A., Dinh, M., Dolson, E., Farrokhan, N., Gopalakrishnan, V., Ho, E., King, E., Krishnan, N., Kuzmin, G., Maltas, J., Pelesko, J., Scarborough, J., Scott, J., Sedor, G., & Weaver, D. (2020). *UV Sterilization of Personal Protective Equipment with Idle Laboratory Biosafety Cabinets During the Covid-19 Pandemic*. <https://doi.org/10.1101/2020.03.25.20043489>
7. Association for the Advancement of Medical Instrumentation. (2020, April 24). *N95 Mask Decontamination and Reuse Webinar*. www.aami.org
8. CDC Coronavirus Disease 2019: Decontamination and Reuse of Filtering Facepiece Respirators <https://www.cdc.gov/coronavirus/2019-ncov/hcp/ppe-strategy/decontamination-reuse-respirators.html>

INTUBATION BARRIER STRATEGIES for Use during the COVID–19 Pandemic: April 2020

Situation and Background

Concerns have been raised over the need to implement additional strategies to prevent COVID–19 disease transmission during patient intubation. To address these concerns, a number of products and strategies have been developed by facilities and vendors. To date, there are no clear industry standards or guidance that definitively recommend one approach as more effective over another. Given the lack of clear guidance, careful review and evaluation of each method is warranted prior to consideration for implementation.

Assessment

Although there are additional prototypes and products currently in use, three primary strategies for barrier protection during endotracheal intubation include the COVID-19 Airway Management Isolation Chamber (CAMIC), the Aerosol Box, and the Intubation Shield. Each of these products/strategies maintain unique benefits and challenges with implementation, as highlighted below:

- The CAMIC is constructed primarily out of PVC pipes and a clear polyethylene bag. One unique benefit of this approach is that its flexible structure is believed to allow for improved provider mobility during intubation. Caution must be applied in setting up this system, as it is designed to be hooked to suction on one side, and air or oxygen input on the other. If both ports are accidentally hooked up to oxygen or air, there is a potential to increase aerosolization of the virus. The CAMIC is relatively inexpensive to construct, and initial testing supports this method as an effective means of reducing respiratory droplet spread during aerosol generating procedures (AGPs).¹ Understanding that the CAMIC is not intended to be disposed of after a single use, careful consideration must be made regarding the use of effective cleaning practices. Specifically, the following steps should be followed for CAMIC cleaning:
 - At the point of generation, dispose of polyethylene bag and wipe down the PVC pipe with a hospital–approved germicidal wipe.
 - Place wiped down equipment in a transport container and transport to soiled utility room.
 - Obtain 1:10 premixed bleach disinfectant and submerge pieces for 5 minutes (refer to guidelines provided as an appendix to this document regarding premixed bleach disinfectant).
 - Remove equipment and rinse, ensuring removal of any residue.
 - Dry for 24 hours in an upright position to ensure drainage of remaining water.
 - Place in peel pouch and return to clean storage.
- The Aerosol Box is constructed from acrylic or transparent polycarbonate, and is intended to serve as an additional barrier for protection while allowing the provider to insert their hands through pre–drilled holes to perform intubation.² As with the CAMIC, the Aerosol Box must undergo thorough cleaning prior to reuse in order to avoid inadvertent disease transmission. Aligned with developer recommendations, cleaning should be performed with an Environmental Protection Agency approved disinfectant wipe. Clean water or alcohol may then be utilized to remove any visible residue.
- The Intubation Shield is a single-use/disposable, clear plastic drape placed over the patient during intubation. The Intubation Shield can be utilized as a single–method for protection, or in conjunction with additional barrier strategies. One unique benefit of the Intubation Shield is that since it is disposed of after a single use, there are no cleaning requirements. Caution must be applied with this product, understanding that it may pose a risk of suffocation if left in place without direct supervision (clipping the sheet to an IV pole may reduce this risk). Healthcare workers must receive proper training in the use and supervision of these barriers.

Recommendation

Military Medical Treatment Facility (MTF) leadership, in collaboration with frontline providers, may consider each of the aforementioned strategies as options for implementing additional barrier precautions during AGPs such as endotracheal intubation. Healthcare workers must be trained properly on the use of these barriers, and operators should be ready to abandon their use should airway management prove difficult. Decision–making should take into account each MTF’s existing resources and needs. Commanders must pay close attention that proper cleaning procedures are implemented for use these products, as well as any strategy involving reuse of materials.

References

1. Blood, T., Perkins, J., Riley, C., Ruhl, D., Wistermayer, P., Fisher, N., & Hong, S. (2020). *COVID-19 Airway Management Isolation Chamber (CAMIC)*.
2. Lai, H. Y. (2020). *Aerosol Box*. <https://sites.google.com/view/aerosolbox/home>
3. Protection Products Inc. (2020). *Intubation Shield*. <https://p-p-i.com/intubation-shield>
4. Canelli, R. (2020) Barrier Enclosure during Endotracheal Intubation. <https://www.nejm.org/doi/full/10.1056/NEJMc2007589>

BLEACH USE FOR INTERMEDIATE DISINFECTION (of CAMIC PVC)

Preparing a 0.5–0.6% sodium hypochlorite (i.e. “1:10 bleach”) solution for disinfection (refer to table for correct dilution ratios).

5.25% to 6% Sodium hypochlorite (household bleach)	Cold Tap Water
380 mL (1 cup and 5 ounces)	3.8 Liters (1 gallon)
65 mL (2 ounces)	650 mL (22 ounces)
45 mL (3 TABLEspoons)	474 mL (16 ounces or 2 cups)
23 mL (1.5 TABLEspoons)	237 mL (8 ounces or 1 cup)

- When mixing bleach, wear gloves and eye protection.
- Consider a waterproof apron or gown to avoid getting on clothing.
- Mix bleach in a well ventilated area.
- Mix bleach using cold water, as hot water decomposes it.
- Do not mix with other chemicals.
- Discard after each use.
 - If bleach is reused, mix in an opaque bottle/container and discard after each shift/day.
 - Clearly label and date the container of the bleach solution.
 - Keep diluted bleach covered and protected from sunlight, and if possible in a dark container.

Procedure for Disinfection

- Items will be cleaned with a 0.5% to 0.6% sodium hypochlorite (i.e. “1:10 bleach”) solution, then thoroughly rinsed or wiped with clean water to remove any residual and then dried.
- Sodium hypochlorite solutions (mixed/diluted) will gradually lose strength, so fresh solutions must be prepared frequently.
- Diluted solutions of bleach will be replaced after each use.
- Items soaking for 5 minutes must be in a well vented room.
- Proper PPE must be worn.
 - Wear disposable gloves when cleaning and disinfecting surfaces.
 - Gloves should be discarded after each cleaning. If reusable gloves are used, those gloves should be dedicated for cleaning and disinfection of surfaces for COVID-19 and should not be used for other purposes.
 - Clean hands immediately after gloves are removed.
 - If surfaces are dirty, they should be cleaned using a detergent or soap and water prior to disinfection.
- Allow proper ventilation during and after application.
- Never mix household bleach with ammonia or any other cleanser.
- Unexpired household bleach will be effective against coronaviruses when properly diluted.

Situation and Background

Concerns have been raised regarding an increased risk for the bacterium *Legionella pneumophila* and other waterborne pathogens as a result of facility/unit closure throughout the SARS-CoV-2 pandemic. In persons at risk for infection (e.g., individuals who are over 50 years of age, are smokers, immunocompromised, or have underlying medical conditions), this bacterium can lead to a life-threatening pneumonia, called Legionnaires' disease. It is particularly important to note that *Legionella* infection can oftentimes mimic SARS-CoV-2 presentation. Outbreaks are linked to poorly maintained building water systems, especially those that are extensive or complex. Even in the setting of a long-term disinfection program, outbreaks of *Legionella* were noted. Transmission can occur via aerosols from devices such as showerheads, cooling towers, hot tubs, and water fountains.

Throughout the SARS-CoV-2 pandemic, many facility units/areas were either closed or experienced reduced operations. In some instances, water was completely shut off (e.g., water fountains). Such closures and interruptions in normal operations have created an ecosystem that supports the growth of *Legionella* and other waterborne pathogens. It is therefore important to implement strategies to prevent healthcare-associated infections prior to reopening units, bringing employees back from telework, and/or turning on water fountains. In order to appropriately mitigate the risk for opportunistic infections, Military Medical Treatment Facilities (MTFs) must develop and adhere to policies and procedures that inhibit microbial growth and spread of *Legionella* and other waterborne pathogens in building water systems.¹ Dental Treatment Facilities (DTFs) will continue to follow American Dental Association (ADA) and CDC guidelines for dental water lines and/or other governing authority standards as applicable (e.g, The Joint Commission).

Assessment

The Joint Commission (TJC) maintains standards requiring facilities to protect the health and safety of patients through establishment of a water management program that reduces the risk of growth and spread of *Legionella* and other opportunistic pathogens in facility water systems. TJC evaluates evidence of compliance with the following key elements. (Non-accredited DTFs will follow the national CDC and ADA guidelines in regard to water management programs):

- Completion of a facility risk assessment to identify where *Legionella* and other opportunistic waterborne pathogens could grow;
- Development and implementation of a water management program with corresponding testing protocols; and
- Establishment of testing protocols and acceptable ranges for control measures.²⁻³
 - Although TJC recommends establishment of testing protocols and acceptable ranges for control measures, more recent information is showing that interpretative results from *Legionella* cultures are variable. As a result, the Centers for Disease Control and Prevention (CDC) discourages the use of thresholds using colony forming units (CFU)/mL. Until more precise tests are available, any detectable level at a single site should be considered a hazard.

In 2016, the CDC and its partners developed a toolkit to facilitate implementation of the American Society of Heating, Refrigerating, and Air Conditioning Engineers (ASHRAE) standards. This comprehensive toolkit provides environmental, clinical, and epidemiologic considerations for healthcare facilities, and can be accessed with the following link:⁴ <https://www.cdc.gov/legionella/maintenance/wmp-toolkit.html>.

Risk Factors

Several factors associated with building water systems can contribute to amplification of pathogenic bacteria, thereby leading to increased risk of exposure:

1. Water temperature fluctuations

Clinical Management of COVID-19, v4

- Temperatures between 25° - 42° C (77° - 108° F) are ideal for *Legionella* growth.⁴
 - *Legionella* can continue to proliferate at temperatures outside the above range.
2. The presence of free chlorine (chlorine residual) available in water to kill microorganisms. Click here for more information: <https://www.cdc.gov/safewater/chlorine-residual-testing.html>
 - Free chlorine levels should be maintained at a minimum of 0.5 mg/L (0.5 ppm or parts per million) after at least 30 minutes contact time, at pH less than 8.0.
 3. pH level
 - The pH of potable water impacts the efficacy of chlorine disinfection. Such disinfectants are most effective within a narrow pH range of approximately 6.5-8.5.
 4. Stagnation/flow
 - Low flow or stagnant water leads to increased water age, which depletes free chlorine.
 - Stagnation encourages biofilm growth and promotes sediment accumulation, which uses up disinfectant and provides nutrients for bacteria colonization.
 - Changes in water pressure can dislodge biofilm, subsequently releasing *Legionella* into the water.
 5. Aerosolization
 - Devices or processes that aerosolize water increase risk for inhalation and infection.
 6. Immunocompromised patients are at higher risk of contracting Legionnaires' disease from inhalation of contaminated water.

Recommendation

As considerations are made regarding reopening of facilities and units previously closed during the SARS-CoV-2 pandemic, leadership must ensure that a comprehensive water management program addressing safety concerns related to disruption in water flow are in place, and fully implemented.

As units begin to reopen, a multidisciplinary team to ensure proper implementation of the water management program is essential. At minimum, this team should include the following representatives: Preventive Medicine, Infection Prevention and Control, Facilities Management, Department of Medicine or Infectious Disease Physician (as applicable to in-patient facilities), Division of Nursing, Industrial Hygiene, Clinical Laboratory (Microbiology), and contracted subject matter experts. Primary responsibilities of this team must include, but are not limited to, the following:

- Reviewing the facility's water management plan and implementing strategies to mitigate risks prior to turning on water systems where a disruption in flow occurred.
 - This applies both to areas completely shut down, and those where a significant reduction in water use was noted.
- Performing a risk assessment prior to opening areas or turning on water systems (e.g., water fountains).
 - The risk assessment must take into account water temperatures, pH levels, and chlorine concentration following water disruptions.
- Monitoring microbiological data as the systems are returned to normal operation.
- Reporting cases of suspected and confirmed hospital-associated *Legionella* transmission (includes patients and staff).

Clinical Management of COVID-19, v4

- Implementing water testing/treatment protocols as described in the facility's water management plan.

During the SARS-CoV-2 crisis, clinicians have maintained a high degree of suspicion for SARS-CoV-2 in patients presenting with respiratory illnesses. However, clinicians should also test patients with healthcare-associated pneumonia for Legionnaires' disease as described in the CDC toolkit. This is especially important in circumstances where *Legionella* growth risk factors are/may have been present (e.g., areas where water stagnation may have occurred). The preferred diagnostic tests for Legionnaires' disease include cultures of lower respiratory secretions on selective media and the *Legionella* urinary antigen test.

Facilities should utilize the CDC's comprehensive toolkit referenced in this document to ensure their water management program appropriately incorporates all industry recommended *Legionella* and other waterborne pathogen prevention strategies. Lastly, facilities must ensure their water management program is properly aligned with current TJC standards to effectively protect the safety and health of those in the facility, as well as to avoid adverse accreditation action.

References

1. Center for Medicare & Medicaid Services. (2017). Requirement to Reduce Legionella Risk in Healthcare Facility Water Systems to Prevent Cases and Outbreaks of Legionnaires' Disease (LD).
2. The Joint Commission (2020) *Comprehensive accreditation manual for hospitals*.
3. Centers for Disease Control and Prevention. (2017). *Toolkit: Developing a Water Management Program to Reduce Legionella Growth and Spread in Buildings*. <https://www.cdc.gov/legionella/wmp/toolkit/index.html>
4. American Society of Heating, Refrigerating, and Air Conditioning Engineers. (2018). *Legionellosis: Risk Management for Building Water Systems*. www.ashrae.org
5. Lin, Y. E., Stout, J. E., & Yu, V. L. (2011). Controlling Legionella in hospital drinking water: An evidence-based review of disinfection methods. *Infection Control and Hospital Epidemiology*, 32(2), 166–173. <https://doi.org/10.1086/657934>
6. The American Dental Association website: <https://www.ada.org/en/member-center/member-benefits/practice-resources/the-joint-commission-tjc>

COVID–19 Recovery Plan: Guidance for IPC Programs - 21 May 2020

Introduction:

As with any crisis situation, Military Medical and Dental Treatment Facilities (MTFs/DTFs) will need to take a strategic approach to optimize efficiency in recovery from the COVID–19 pandemic. Infection Prevention and Control (IPC) Programs at the MTF/DTF level face unique challenges, and the guidance provided within this document is intended to facilitate a coordinated approach to integration of best practices in alignment with current Centers for Disease Control and Prevention (CDC) and other evidence–based guidelines and standards.

Recommendations

Infection Preventionist' s (IPs) must work closely with multidisciplinary team members, leadership, and logistics to optimize recovery efforts and ensure a comprehensive strategy remains in place. This document provides a high–level overview of recommendations for the following topics, as they relate to preparation for return of operations with pandemic resolution:

1. General IPC Program Preparation
2. Administrative Controls
3. Environment of Care
4. Water Plans
5. Personal Protective Equipment (PPE)
6. Additional IPC Considerations

Understanding that identified needs and existing resources are unique to each facility, the recommendations provided in this document are intended to serve as a guide, and are not an all–inclusive list of necessary actions.

General IPC Program Preparation

1. A comprehensive IPC Risk Assessment for each facility must be performed based on review of national, state, and local COVID–19 specific epidemiology. As leaders make decisions to resume patient care services, it is important that IPC guidance is provided to prevent and/or mitigate potential harm to patients and health care employees.

In collaboration with the IPC committee and senior leadership, IPs should develop an IPC plan for de-escalation that addresses cleaning of patient care areas, equipment, and other environment of care requirements prior to re-opening clinical spaces. IPs should consider using a checklist similar to one that is used when reopening areas after a major renovation project. Additionally, the Patient Safety Checklist from the Defense Health Agency Memorandum “COVID-19 Guidance for Resuming Full Healthcare Operations” signed by RADM Riggs May 2020

2. Basic IPC education should resume. At a minimum, such training should address standard precautions and disease transmission based precautions. Training on donning and doffing of PPE should become a routine requirement to prepare for a potential surge or other emerging disease. Additional information regarding PPE is available at the following link: <https://www.cdc.gov/coronavirus/2019-ncov/hcp/using-ppe.html>
3. Facilities must continue to follow the most up to date guidance from the CDC regarding management of known or suspected cases of COVID–19.
4. Health care personnel must continue to assess potential risks of exposure during patient encounters, as well as ensure safe work practices, administrative, and engineering controls are in place in alignment with current guidelines and standards of practice.
5. IPC policies should be flexible, allowing for updates to be made based on new CDC and DHA guidance regarding universal source control.

Clinical Management of COVID-19, v4

6. If patient care items are reused between patients, health care workers must follow manufacturer's instructions for use (IFUs) and adhere to guidelines for cleaning and disinfection.
7. As COVID-19 patient care units are no longer required, IPC plans must address optimal patient placement of known or suspected cases within a facility. If admitted, place a patient with known or suspected COVID-19 in a single-person room with the door closed. The patient should have a dedicated bathroom. Airborne Infection Isolation Rooms (AIIRs) should be reserved for patients who will be undergoing aerosol generating procedures (AGPs).

Administrative Controls

1. Administrative controls are defined as changes in policy or procedures to reduce and/or minimize exposure to infectious diseases.
2. Facilities must maintain heightened awareness in triaging/assessing patients and staff for potential COVID-19 related symptoms. Leadership must also consider how to implement measures to mitigate risk of disease transmission while returning to normal operations.
3. Disease-specific clearance requirements for return to work must be established by occupational health and implemented based on CDC guidelines. Also, a process for documenting clearance results for both staff and patients must be in place.
4. Sick employees must be encouraged to remain home.
5. Leadership should consider establishing alternating days or extra shifts that reduce the total number of employees in a facility at a given time, allowing them to maintain distance from one another while maintaining a full work week during the ongoing COVID-19 pandemic.
6. All facilities should consider the implementation of a visitation policy that is in alignment with processes established for screening patients and staff.

Environment of Care

Facility Considerations

IPs, in collaboration with Facilities Management and first-line leadership (e.g., C-suite) maintain responsibility for the following:

1. Establishing non-COVID-19 care zones that screen all patients for symptoms of COVID-19 (including temperature checks). As stated previously, all staff, patients, and visitors should continue to be routinely screened.
2. Considering areas for non-COVID care (i.e., separate building, designated rooms, or floor with a separate entrance and minimal crossover with COVID-19 areas). The use of segregated hallways/paths for transport of COVID-19 positive patients to minimize exposure to others should be also be considered.
3. Ensuring proper signage is in place to instruct patients on requirements for building entry (e.g., screening, source control (face coverings), and social distancing).
4. Ensuring facility, administrative, and engineering controls have been established to facilitate social distancing. Examples include, but are not limited to, minimizing time in waiting areas, spacing chairs at least 6 feet apart, installing stanchion barriers, and maintaining low patient volumes.
5. Considering installing physical barriers (e.g., plastic sneeze guards) in non-clinical areas such as pharmacy and medical records.
6. Ensuring all ventilation requirements are met. This includes evaluating all AIIR and rooms that have been, or may have been modified with air scrubbers/high-efficiency particulate air (HEPA) filtration.

Clinical Management of COVID-19, v4

7. Delaying entry into rooms where an AGP was performed until sufficient time has elapsed to allow for enough air changes to remove potentially infectious particles. Entry should also be avoided before terminal cleaning is completed.
8. Ensuring all sinks have hand washing supplies available, and are not expired. The establishment of cough etiquette stations throughout the MTF should be considered based on local resources and needs. Items for consideration to include at each cough etiquette station include signage, tissues, masks, and alcohol-based hand sanitizer. Additionally, facilities must ensure hand washing/hygiene signs are posted at hand washing stations as appropriate.

Sanitation Protocols

IPs, in collaboration with Facilities Management and first-line leadership (e.g., C-suite) maintain responsibility for the following:

1. Ensuring there is an established plan for thorough cleaning and disinfection prior to using spaces or units that may have been closed during the COVID-19 crisis.
 - a. Consideration may be required to modify housekeeping contracts to increase frequency of cleaning.
 - b. Environmental Service (EVS) personnel should refrain from entering vacated rooms until sufficient time has elapsed for enough air changes to remove potentially infectious particles.
 - c. After the correct time has elapsed, EVS personnel may enter the room and should wear a gown and gloves when performing terminal cleaning. A mask and eye protection should be added if splashes or sprays during cleaning and disinfection activities are anticipated, or otherwise required based on the selected cleaning products.
2. Ensuring any equipment that was taken out of service is cleaned/disinfected in accordance with manufacturer's IFU prior to use.
3. Ensuring that equipment such as anesthesia machines used for COVID-19 positive patients, or any patient who has a disease that can potentially spread via the environment (e.g., Vancomycin-resistant enterococci) is thoroughly cleaned in accordance with CDC guidelines.
4. Ensuring staff understand the management of standard/office waste and regulated medical waste in accordance with local/state requirements.
5. Ensuring housekeeping rotates linen and privacy curtains in areas where COVID-19 patients were treated, and in areas that were closed during pandemic if contaminated.

Supplies and Linen

IPs, in collaboration with Facilities Management and first-line leadership (e.g., C-suite) maintain responsibility for the following:

1. Assessing supply and linen rooms to ensure they meet IPC recommendations for storage and cleanliness.
2. Ensure section/unit personnel check expiration dates for all supplies.

Cleaning, Sterilization, and High-Level Disinfection

All staff engaged in cleaning, sterilization, and high-level disinfection are responsible for the following:

1. Inspecting all sterile packages and instrument trays for integrity and expiration dates.
2. Ensuring all washer/decontaminators, sterilizers, automated endoscope reprocessors, ultrasonic machines, and other sensitive equipment have been tested (QC) to verify appropriate parameters are met.
 - a. Contact biomedical maintenance, if necessary
 - b. Maintain documentation

Clinical Management of COVID-19, v4

3. Assessing sterilant and disinfection solutions to confirm stability and date of expiration per manufacturer's IFU.
4. Reviewing endoscope reprocessing protocols if endoscopic procedures are performed. If there is a scheduled reprocessing interval (hang time), reprocess in accordance with local policy.
5. Reviewing competencies and IPC training for personnel who perform disinfection and sterilization practices, including personnel who handle instruments at the point of use. Consideration for retraining should be based on individual need and length of time passed since the activity was last performed.

Water Plans

1. To prevent waterborne pathogen outbreaks, facilities will need to take water plans into consideration as units reopen and/or water is turned back on.
2. As part of the facility's water management plan, the following minimum requirements must be reviewed prior to opening:
 - a. Certify all sinks, showers, fountains, dental water lines, etc. are flushed in spaces that were closed or not used during the COVID-19 crisis.
 - b. Confirm ice machines have been maintained in accordance with manufacturer recommendations. In the absence of manufacturer recommendations, refer to CDC guidelines and recommendations.
3. Additional guidance includes policy on *Legionella* & other waterborne pathogen risk mitigation.


Personal Protective Equipment

1. As Health Protection Condition (HPCON) levels are reduced and operational status begins to normalize, MTF/DTF leadership must ensure staff are aware that all PPE extended and reuse strategies utilized during the pandemic must be discontinued when sufficient levels of critical PPE are achieved and able to be maintained.
2. MTF/DTF leadership should work in close collaboration with logistics to develop appropriate stockpile quantities for critical PPE and supplies, in preparation for a potential second pandemic surge. In particular, the following should be considered for stockpile supply:
 - a. Alcohol-based hand sanitizer (>60% alcohol content)
 - b. Liquid hand soap
 - c. Face masks (i.e. surgical masks)
 - d. Face shields
 - e. Eye protection
 - f. NIOSH approved surgical N95 respirators
 - g. Isolation gowns
 - h. Shoe and head covers
 - i. Gloves
 - j. Disinfecting surface wipes
 - k. Isolation signs
3. Facilities should develop and maintain a plan for decontaminating N95 respirators in the event of a critical shortage during a second-wave pandemic. Such a plan must demonstrate alignment with existing resources and needs.

Clinical Management of COVID-19, v4

- All staff must understand that industry guidelines continue to evolve, however, the following algorithm provided by the Infectious Diseases Society of America (IDSA) is a helpful resource in terms of caring for patients with suspected or known COVID-19 during either conventional or crisis situations.

Figure 1. IDSA Algorithm for Appropriate PPE in Conventional and Contingency or Crisis Settings



AGP: aerosol-generating procedures; PPE: personal protective equipment

Additional IPC Considerations

Discontinuation of transmission based precautions for patients with COVID-19 should be made using one of the following three strategies, based on current clinical evidence:

1. Test-based
2. Symptom-based
3. Time-based

The decision to discontinue empiric transmission-based precautions by excluding the diagnosis of COVID-19 for a suspected COVID-19 patient can be made based on obtaining negative results from at least one Food and Drug Administration Emergency Use Authorized COVID-19 molecular assay for detection of SARS-CoV-2 RNA. Still, clinical judgment and suspicion of SARS-CoV-2 infection must be applied to determine whether to continue or discontinue empiric transmission based precautions.

Additional information regarding discontinuation of isolation is available at the following link:

<https://www.cdc.gov/coronavirus/2019-ncov/hcp/disposition-hospitalized-patients.html>

References

1. Centers for Medicare & Medicaid Services (CMS) Recommendations Re-opening Facilities to Provide Non-emergent Non-COVID-19 Healthcare: Phase I. <https://www.cms.gov/files/document/covid-flexibility-reopen-essential-non-covid-services.pdf>
2. U. S. Department of Labor, O. S. (2020, 03). *Guidance on preparing workplaces for COVID-19*. Retrieved from <https://www.osha.gov/Publications/OSHA3990.pdf>
3. Lynch, J., Davitkov, P., Anderson, D., Bhimraj, A., Chi-Chung Cheng, V., Guzman-Cottrill, J., Dhindsa, J., Duggal, A., Jain, M., Lee, G., Liang, S., McGeer, A., Lavergne, V., Murad, H., Mustafa, R., Morgan, R., Falck-Ytter, Y., & Sultan, S. (2020). *Infectious Diseases Society of America Guidelines on Infection Prevention for Health Care Personnel Caring for Patients with Suspected or Known COVID-19*. www.idsociety.org/COVID19guidelines/ip
4. Defense Health Agency. (2020). *Memorandum: COVID-19 Guidance for Resuming Full Healthcare Operations*.

USE OF MASKS IN THE COMMUNITY SETTING

As COVID-19 extends its reach into local communities throughout the United States, the use of masks in public is now a topic of debate. The leading health organizations in the world offer mixed recommendations regarding the use of masks in the community setting. To address this topic, I will look at the components of droplet-producing events (e.g. sneezing), CDC and WHO mask recommendations, the performance of different types of masks, and finally how to care for masks in the community setting.

What's in a Sneeze?

High-speed recordings of droplets following a sneeze demonstrate turbulent clouds, with the largest droplets settling 1-2 meters away. Smaller droplets remain suspended, carrying them as far as 6-8 meters away from the individual. The lifetime of these suspended droplets, particularly in warm and moist air, could extend as much as 1000 times that of larger droplets (from seconds to minutes).¹

Currently, both the Center for Disease Control (CDC) and World Health Organization (WHO) consider the airborne transmission of COVID-19 to be unlikely.^{2,3} Both organizations encourage droplet precautions for healthcare workers and social distancing of 6 feet in the community; however, research proving that COVID is not transmissible through airborne droplet nuclei is lacking. One recent study, published in the April 16th edition of the *New England Journal of Medicine*, demonstrated the viability of SARS-CoV-2 in aerosols under laboratory conditions.⁴ This does not address whether aerosolized droplet nuclei containing COVID-19 would deliver a big enough inoculum to infect another person. Ultimately, more concrete research is needed to determine the true risk to community members outside of a 1-2 meter radius.

Organizational Recommendations for Community Masks

At the time of this edition's publication, the Center for Disease Control (CDC) argues that citizens can utilize cloth face coverings in various settings, such as grocery stores, where social distancing can be more difficult. The CDC emphasizes that this measure protects those around you by limiting the spread of droplets from an otherwise asymptomatic individual. Their guidance regarding cloth face coverings include a snug fit, secured with ear-loops or ties, be multi-layered, and may be laundered without damage. Finally, the CDC cautions against placing masks on children younger than 2 or on any individual with trouble breathing or who would otherwise have difficulty removing the mask.⁵

The World Health Organization (WHO), however, recommends against the use of any type of mask in the community setting. They portend that there is no evidence that masks can prevent individuals in the community from contracting a respiratory virus. Furthermore, the WHO argues the use of masks will create complacency regarding other preventive measures. Ultimately, they recommend decision-makers keep in mind the purpose of using masks, who in the local population is at risk, characteristics of the community setting, the feasibility of producing and distributing masks, and the type of masks being recommended.⁶

So How Good are Masks?

With particles of COVID-19 measuring 60-140 nanometers (0.06-0.140 microns)⁷, there is a reasonable concern regarding how effective masks, particularly cloth ones, are in blocking the virus. One recent laboratory trial assessed the effectiveness of N95, surgical masks, homemade masks, and cloth masks in filtering avian influenza virus in aerosols. Ma and colleagues found N95 masks 99.98% of the virus, while medical masks blocked 97.14%, and homemade masks (1 layer of polyester cloth with 4 layers of kitchen paper) could block 95.15%. They cautioned that cloth alone may not confer any protection from the virus.⁸

One prospective, randomized control trial from 2015 evaluated 1607 healthcare workers across seventy-four hospital wards with the use of medical and cloth masks. They found the rate of influenza-like illness was significantly higher in the cloth mask arm (RR = 13) compared to those wearing medical masks. They additionally conducted a sodium chloride filtration test, demonstrating a 97% penetration rate in cloth masks, compared to 44% in medical masks and <0.01% in N95 masks.⁹

An additional laboratory-based study assessed the filtration efficiency of surgical masks compared to masks created from various household materials. First, the investigators assessed filtration efficiency for blocking *Bacillus*

Clinical Management of COVID-19, v4

astrophageus (0.95-1.25 microns) and Bacteriophage MS2 (0.023 microns) for all materials. Cotton was 69% efficient for *Bacillus* and 50% efficient for Bacteriophage, compared to 96% and 89% respectively for a surgical mask. Next, healthy volunteers were asked to cough twice into a 0.5m³ box. Investigators took air samples for the next five minutes, and the median colony forming units (CFUs) were measured for air and settle plates within the box. The reduction of CFUs was statistically significant for both the surgical masks and for the cloth masks; however, the surgical mask was three times better at blocking the transmission of expelled microorganisms. Davies and colleagues concluded poor fit will reduce effectiveness and that homemade masks are not only a last resort, but they would have minimal effect if not combined with hand hygiene and other preventive measures.¹⁰

Caring for Masks

A recent article in *JAMA* recommends that cloth masks should be washed routinely with laundry detergent or with soapy water.¹¹ The CDC also states these facial coverings should be washed “routinely” depending on how often they are worn. For items placed in the laundry, the warmest possible setting should be utilized.⁵

For those using surgical masks in the community, the CDC discusses several potential methods for sterilizing masks while minimizing degradation in overall fit and performance. The most practical is with moist heat, based on studies performed using microwave steam bags and microwave-generated steam.¹² Based on the need for steam temperatures of 60-65° Celsius and 80-85% humidity for only a couple of minutes, it may be reasonable for individuals at home to hold their masks in the steam from a boiling pot of water for 2-3 minutes to achieve sterilization.

Conclusion

Current recommendations that members of the community wear masks in settings where social distancing measures may be more difficult to maintain. Providers and or healthcare professionals must emphasize using the masks as a way of further limiting the spread of COVID from asymptomatic patients, rather than as a means of protecting yourself from others. Masks of any type should be form-fitting and frequently laundered or sterilized. Surgical masks should generally be reserved for healthcare personnel; however, there are some grass-roots efforts by military garrisons to produce reusable masks for service members and dependents. Finally, members of the community must understand that masks are part of a multi-faceted approach to reducing viral transmission. Frequent hand-washing, not touching your face with unwashed hands, and maintaining social distancing practices remain vital in the fight against SARS-CoV-2.

CDC webpage that includes a description with photos on how to make a mask at home:

<https://www.cdc.gov/coronavirus/2019-ncov/prevent-getting-sick/diy-cloth-face-coverings.html>

Providence St Joseph Health webpage about making surgical facemasks:

https://psjh.blob.core.windows.net/covid/2020-%20PSJH_CLINICALFACEMASK_%20Instructions%20-%20Final.pdf


References

1. Baden LR. Images in Clinical Medicine: A Sneeze. *N Engl J Med*. 2016; e15.
2. CDC. Coronavirus Disease 2019: Infection Control Guidance. 2020 Apr 13. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/hcp/infection-control-recommendations.html#adhere>.
3. WHO. Modes of transmission of virus causing COVID-19: Implications for IPC precaution recommendations. 2020 Mar 29. Available from: <https://www.who.int/news-room/commentaries/detail/modes-of-transmission-of-virus-causing-covid-19-implications-for-ipc-precaution-recommendations>.
4. van Doremalen N, Bushmaker T, et al. Aerosol and Surface Stability of SARS-CoV-2 as Compared with SARS-CoV-1. *N Engl J Med*. 2020 Apr 16;382(16): 1564-7.
5. CDC. Coronavirus Disease 2019: Cloth Face Covers. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/prevent-getting-sick/diy-cloth-face-coverings.html>.
6. WHO. Advice on the use of masks in the context of COVID-19: Interim Guidance. 2020 Apr 6. Available from: [https://www.who.int/publications-detail/advice-on-the-use-of-masks-in-the-community-during-home-care-and-in-healthcare-settings-in-the-context-of-the-novel-coronavirus-\(2019-ncov\)-outbreak](https://www.who.int/publications-detail/advice-on-the-use-of-masks-in-the-community-during-home-care-and-in-healthcare-settings-in-the-context-of-the-novel-coronavirus-(2019-ncov)-outbreak).
7. Cascella M, Rajnik M, Cuomo A, et al. Features, Evaluation and Treatment Coronavirus (COVID-19) [Updated 2020 Apr 6]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2020 Jan-. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK554776>.
8. Ma QX, Shan H, et al. Potential utilities of mask-wearing and instant hand hygiene for fighting SARS-CoV-2. *J Med Virol*. 2020 Mar 31 [Epub ahead of print available from: <https://doi.org/10.1002/jmv.25805>]
9. MacIntyre CR, Seale H, et al. A cluster randomized trial of cloth masks compared with medical masks in healthcare workers. *BMJ Open*. 2015 Apr 22;5(4):e006577.
10. Davies A, Thompson KA, et al. Testing the efficacy of homemade masks: Would they protect in an influenza pandemic? *Disaster Med Public Health Prep*. 2013 Aug;7(4): 413-8.
11. Desai AN & Aronoff DM. Masks and Coronavirus Disease 2019 (COVID-19). *JAMA*. 2020 Apr 17 [Epub ahead of print available from: <https://jamanetwork.com/journals/jama/fullarticle/2764955>]
12. CDC. Coronavirus Disease 2019: Decontamination & Reuse of Filtering Facepiece Respirators. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/hcp/ppe-strategy/decontamination-reuse-respirators.html>.

APPENDIX C : EXAMPLE TRIAGE PROTOCOLS DURING COVID-19 PANDEMIC


COVID-19 Telephone Triage Protocol

As of 1600 on 19 May 2020




COVID-19 In-clinic Protocol

As of 1100 on 11 June 2020




COVID-19 Home Isolation Discontinuation Protocol (Symptomatic)

As of 1600 on 19 May 2020




COVID-19 Home Isolation Discontinuation Protocol (Asymptomatic)

As of 1600 on 19 May 2020




Clinical Management of COVID-19, v4



COVID-19 Calculator*

		Points
C CONTACT	Does the patient have a known COVID-19 contact?	
	Yes	3
	No	0
O OTHER Sxs	Does the patient endorse cough, dyspnea, myalgias, headache, sore throat, anosmia or diarrhea?	
	Yes (1 point each)	1 to 7
	No	0
V VITALS	Does the patient have either reported/recorded fever (>100.4F) or hypoxemia (SpO2 < 90%)?	
	Yes (1 point each)	1 to 2
	No	0
I IMAGING	Is imaging suggestive of bilateral infiltrates or GGOs (without alternative explanation)?	
	Yes	1
	No	0
D DIFFERENTIAL	Does the patient have lymphopenia (< 1,500/ml)?	
	Yes	2
	No	0

*The above calculator should only be utilized in the context of a negative COVID-19 PCR test and does not have ability to detect asymptomatic cases. If further risk stratification may be indicated. Last updated: 4 May 2020



Last updated: 4 May 2020

Return to Work Guidelines for Individuals Diagnosed with Coronavirus Disease (COVID-19)

To help prevent the spread of 2019 coronavirus disease (COVID-19) in the community, Department of Defense (DoD) Commanders in coordination with medical experts have instituted transmission-based precautions. Transmission-based precautions include stay home from work policy, restriction of movement (ROM), quarantine, and isolation. Personnel who have symptoms of acute respiratory illness should notify their supervisor and avoid going to work. Personnel who have a high risk exposure to COVID-19 or who have traveled to high risk locations as defined by the Centers for Disease Control and Prevention (CDC) with ongoing community transmission of COVID-19 (Travel Health Notice Levels 2 or 3) should be placed under ROM or quarantine. Personnel who diagnosed with COVID-19 should be placed under isolation at a healthcare facility, home, or another designated isolation facility.

The decision to discontinue transmission-based precautions and allow personnel to return to work (RTW) is based on weighing the overall risks of severe illness and death; balancing the potential benefits of decreasing transmission through the exclusion of ill persons with minimizing social disruption. This guidance will continue to be updated as more information becomes available.

ROM/Quarantine

After travel to a high risk location or close contact with a COVID-19 positive patient, personnel may RTW if no symptoms develop after 14 days since the date of departure from high risk location or, if they remain free of symptoms, the last date of high risk exposure to the infected individual. You **cannot** test-out of quarantine; you must remain in quarantine for the full 14-day incubation period.

Person Under Investigation (PUI)

A PUI is defined as an individual with symptoms consistent with COVID-19 and high risk exposure from travel or close contact; OR based on clinician's judgement. After being identified as a PUI, personnel may RTW when they meet the following criteria:

- Completion of the required 14-day ROM/Quarantine period (if placed under ROM for high risk exposure); and
- Negative COVID-19 test (if testing was performed); and
- Symptom free for at least 72 hours without the use of fever-reducing or symptom-altering medications; and
- At least 10 days have passed since symptoms first appeared.


Confirmed COVID-19 Case

After being diagnosed with COVID-19, personnel may RTW using one of the following clearance criteria:

- Using an FDA Emergency Use Authorization (EUA) approved test:
 - Resolution of fever, without use of fever-reducing medication (e.g., acetaminophen, ibuprofen); and
 - Improvement in respiratory symptoms (e.g., cough, shortness of breath); and
 - Negative results from at least two consecutive nasopharyngeal swab specimens collected ≥24 hours apart (total of two negative specimens)
- Unable to get FDA EUA approved test (i.e. OCONUS facility using foreign national laboratories, or mild symptoms unable to obtain clearance test):
 - At least 3 days (72 hours) have passed since recovery, defined as resolution of fever without the use of fever-reducing medications and improvement in respiratory symptoms (e.g., cough, shortness of breath); and
 - At least 10 days have passed since symptoms first appeared.

Management of a Negative COVID-19 Lab Test

- A negative COVID-19 lab test does not rule out COVID-19 infection. Persons with a negative COVID-19 test should be treated as a presumptive COVID-19 case unless a healthcare provider diagnoses another disease, and the RTW guidance for confirmed COVID-19 cases as described above should be followed.
- If the healthcare provider determines a diagnosis other than COVID-19, e.g., flu, is appropriate, the RTW guidance for that disease should be followed.



Guideline Only/Not a Substitute for Clinical Judgment

APPENDIX D : ADULT PRONE POSITIONING PROTOCOL EXAMPLE*

*Adapted from University Medical Center (Las Vegas, NV)

Procedure for patient preparation prior to proning:

1. Obtain an order from the Fellow or Attending physician to place patient in the prone position. The order should include:
 - a. Proper sedation/pain medications and paralytic agents if necessary.
 - b. Length of time for each pronation cycle (patient should be in prone position a minimum of 16 hours, with a return to the supine position at least once a day).
 - c. Prone positioning should be performed within the first 24 hours of the diagnosis of severe hypoxemia.
2. Explain proning procedure and benefits to patient and family members when present.
3. Prior to proning patient, make sure the following criteria have been met and necessary equipment is made available:
 - a. Patient is mechanically ventilated via a secured endotracheal tube (ETT) with inline suction.
 - b. RT is at bedside to evaluate securement of ETT with commercial tape and to place bite block as needed. Twill may be used in addition to the tape if additional securement is needed. Do not secure ETT with a commercial securement device (i.e. Hollister).
 - c. Confirm patient intravenous access including central and arterial lines; verify lines are secure in place.
 - d. Remove ECG leads from anterior of torso; obtain new leads to place posteriorly once patient is prone. Electrocardiogram leads can be placed in the lateral limb position (left and right deltoid midaxillary line and left and right 12th intercostal space at the midaxillary line). The virtual lead (V1 or chest lead) can be placed on the dorsal surface.
 - e. Consider adhesive foam pads (i.e. Mepilex) to apply to bony prominences such as forehead, bilateral shoulders, chest, iliac crests and knees to prevent pressure ulcers.
 - f. Obtain positioning pillows, blanket rolls or foam prone positioning kit from materials management or supply room.
 - g. Continuous SpO2 monitoring.
 - h. Foley catheter and oral gastric tube secured in place.
 - i. Use fecal management system if needed.
 - j. It is reasonable to provide enteral feedings while patient is in prone position. Elevation of head of bed in reverse Trendelenburg position helps reduce the risk of gastric aspiration. Post pyloric tubes are preferred.
 - k. Lubricate patient's eyes prior to proning, then every six hours and as needed (Provider order needed).
 - l. Assess and document pain and provide adequate sedation and pain management throughout the procedure.
 - m. Patients may also require neuromuscular blocking agent during proning.
 - n. Remove head board and ensure bed brake is on.
 - o. RT will perform and document a complete vent check including auscultation of bilateral lung sounds, ventilator settings, ETT positioning/depth, patient tidal volumes and ETT cuff pressures pre and post turn.

Procedure of manual pronation:

1. Assemble a minimum of a 5-person team consisting of at least on RT and the patient's RN. RT is to manage airway protection at the head of the bed and the other team members are positioned on either side of the bed to manually prone the patient. A fellow or attending physician should be present for the first turn.
2. Correctly position all tubes, taking into account the direction of the turn.
3. Lines inserted in the upper torso are aligned with either shoulder, exception is chest tubes or large bore tubes.
4. Tubes in the lower torso are aligned with either leg and extended off the bed.
5. Always initially turn the patient in the direction of the ventilator.

Procedure for proper patient positioning (see diagram below):

1. Head and Neck positioning:

Place patient's head on a foam head positioner, which allows for the patient's head in a neutral position. Otherwise, support the patient's head in a rotated position paying attention to avoid pressure to the eyes and ears. Provide range of motion to the patient's head at least every hour, maintaining ETT tube alignment. Reposition head every two hours, head should be turned to the up are while in swimmer's pose, to avoid traction on the brachial plexus. Coordinate with RT to be present to maintain the airway while repositioning the head every two hours. This may

Clinical Management of COVID-19, v4

require positioning the ventilator at the head of the bed rather than on one side of the bed to allow for the head reposition. Raise the head enough to provide for proper spinal alignment: avoid hyperextension or flexion of the cervical spine. Ensure the eyes have no pressure on the orbits and ears are properly aligned, flat and not folded.

2. Arm positioning:

If using foam prone positioning kit, place patient's arms in foam positioners. While the patient is in a side lying position, gently position the arms in a swimmer's pose. The swimmer's pose entails the up arm is in a supported, flexed position at the level of the shoulder and the down arm is parallel to the body in a position of comfort. When the arm is in the up position, keep the shoulder in a neutral position, abducted to 90 degrees and the elbow flexed at 90 degrees. Utilize pillows or blanket rolls to prevent hyperextension of the shoulder and to ensure the weight of the arm is supported. Note: Head position should be turned to the up arm while in swimmer's pose, to avoid traction on the brachial plexus.

- a. Alternate the arm and head position every two hours with the patient in a side lying position and provide passive range of motion exercise to all joints of the upper and lower extremities.

3. Patient positioning:

- a. Manually reposition the patient a minimum of every 2 hours with a slight right lateral-pillow support position (20-30°) to prone (flat) to a slight left lateral-pillow supported position (20-30°) and back to prone position. The use of automatic bed rotation is not a replacement for manual repositioning.

Note: When placing the patient in the lateral-pillow support position, coordinate head and arm in the up position toward the tilted side (Do not use foam wedges for lateral turns).

- b. During lateral turns inspect the skin and positioning of the tubes, lines and catheters (tubing and penis) and reposition accordingly, i.e. Foley catheters, chest tubes, IV lines, etc.

4. Leg positioning:

While in prone and/or lateral prone position float the knees with a pillow (be careful not to cause hyperextension of the hip), and place a foam roller, pillow or blanket roll under the ankle area to elevate the toes and prevent tension on the tendons in the foot and ankle region.

5. Tilt the patient into reverse Trendelenburg:

Goal is 30 degrees, as patient tolerates.

6. Alternative position of the arms for comfort or if swimmer's position is contraindicated.

For example, the patient, family or PT/OT one-time evaluation report history of rotator cuff tear, stroke, nerve damage, osteoarthritis of shoulder complex, history of clavicle fracture, hyper flexible joints.

- a. Arms can be left in the side lying position aligned with the body and repositioned every two hours to a slightly abducted position.

Patient monitoring and care:

1. Time patient is prone/supine:

- a. It is recommended in the literature that patient is placed in the prone position for a minimum of 16 hours. The timing for prone cycling requires a physician order and is always situational. Patients should be returned to supine position for up to four hours, once per day preferably early AM to allow the interdisciplinary team time to assess while in supine position. While in supine position, reassessment of oxygenation, skin assessment and other relevant exam elements should occur. If the patient does not tolerate being supine (i.e. requiring increased ventilator settings, decreasing PaO₂/FiO₂ ration, hemodynamically unstable or decreasing SpO₂/PaO₂) return patient to the prone position.
- b. Patients in prone position should receive the same standard of care as a patient that is supine (i.e. oral care,

Clinical Management of COVID-19, v4

urinary catheter care, skin care, eye care, suctioning, etc.).


- c. Discuss supine position tolerance and PaO₂/FiO₂ ratio in bedside report and during interdisciplinary rounds.
- d. Ongoing assessment of how the patient is tolerating prone therapy and repositioning; documentation of all vital signs, capnography, patient and family education, length of time prone, patient's response to turning supine, any adverse events that occur and changes in the patient's condition.
- e. Primary RN will coordinate with RT to re-secure ETT when the patient is supine and assist with turns, checking cuff pressures and tube placement before and after repositioning the patient; coordinate with radiology for chest x-ray when supine.
- f. Monitor all tubes, lines, drains and catheters throughout the repositioning process and continue airway management, suctioning oral and ETT secretions.
- g. Continue to evaluate enteral nutrition tolerance and maintain reverse Trendelenburg to help prevent ventilator associated pneumonia (VAP).
- h. RT to change ETT tape at least once a day or more frequently if necessary due to facial swelling.
- i. PaO₂/FiO₂ ratios should be calculated every day and when ventilator settings have been changed in order to identify candidates for returning to the supine position early.

Consider discontinuation of the prone position if:

- 1. The patient no longer shows a positive response to the position change or mechanical ventilation support has been optimized.
- 2. The patient's PaO₂/FiO₂ ratio is >200 on less than 50% FiO₂ and PEEP ≤10 cm of water.

Complications related to prone positioning:

- 1. Unplanned extubation
 - a. Lines pulled
 - b. Tubes kinked
 - c. Hemodynamic instability
 - d. Facial edema
 - e. Pressure ulcers
 - f. Aspiration
 - g. Corneal abrasions



APPENDIX E : COVID-19 INTUBATION CHECKLISTS, PROTOCOLS, AND COGNITIVE AIDS

COVID-19 INTUBATION PRE-ENTRY CHECKLIST*

For Providers:

To bring inside room:

Place a priority on rapid airway placement with video laryngoscopy (ie Glidescope) to create distance between operator and patient's airway, avoidance of BVM and NIV due to risk of aerosolization:

Airway Supplies:

- ETT (7, 7.5, 8 for adults, appropriate size for children) with syringe for cuff
- Glidescope or C-MAC (facilitate intubation from a distance)
- Appropriate stylet
- Bougie
- OG tube with syringe, lube and tape
- OP/NP airway
- Colorimetric end-tidal CO₂ detector
- Suction setup

Disposable stethoscope

Sani-wipes (should be located inside room)

Keep outside room (on standby):

Back up Airway Supplies:

- Appropriate size laryngoscope blades (Mac 3 & 4 for adults) and handle (disposable preferred)
- Stylet
- BVM (avoid if possible due to risk of aerosolization of pathogen)

Airway cart (never bring in room)

EZ-IO

For Nursing:

RSI meds kit

Restraints

Foley

ABG syringe

Post-intubation meds:

- propofol
- fentanyl
- phenylephrine
- norepinephrine drip

For Respiratory Therapy:

Ventilator with appropriate filters


ET securing device

Waveform capnography adapter

Viral filter for Ambubag

*Adapted from University of Washington (<https://covid-19.uwmedicine.org/>)

COVID-19 INTUBATION PROTOCOL



COVID-19 COGNITIVE AIDS FOR INTUBATION

COVID-19 Emergency Intubation Checklist

CHECK BEFORE ENTERING ROOM



SAFE AIRWAY SOCIETY

Team	Patient	Drugs	Equipment
<ul style="list-style-type: none"> <input type="checkbox"/> Anaesthesia contacted if difficulty anticipated <input type="checkbox"/> Team introduced: <ul style="list-style-type: none"> ▪ Airway Operator ▪ Airway Assistant ▪ Team Leader/Drugs ▪ In-room Runner: optional ▪ Door Runner ▪ Outside room Runner <input type="checkbox"/> Problems anticipated? 	<ul style="list-style-type: none"> <input type="checkbox"/> ECG, BP, Sats <input type="checkbox"/> Pre-oxygenation <ul style="list-style-type: none"> ▪ FIO2 100% ▪ Sitting position 45° <input type="checkbox"/> IV access x 2 <ul style="list-style-type: none"> ▪ 1L fluid on pump set <input type="checkbox"/> Haemodynamics optimised <ul style="list-style-type: none"> ▪ Fluid bolus ▪ Pressor 	<ul style="list-style-type: none"> <input type="checkbox"/> RSI drugs drawn up, doses chosen <input type="checkbox"/> Rescue drugs <ul style="list-style-type: none"> ▪ Metaraminol <input type="checkbox"/> Post intubation sedation plan <input type="checkbox"/> Drug C/I or allergies? 	<ul style="list-style-type: none"> <input type="checkbox"/> 2 Laryngoscopes (tested) <input type="checkbox"/> Tube chosen; cuff tested <input type="checkbox"/> Bougie/stylet <input type="checkbox"/> 10ml syringe <input type="checkbox"/> Tube tie <input type="checkbox"/> Lubricant <input type="checkbox"/> Supraglottic airway sized to pt <input type="checkbox"/> Scalpel + bougie CICO kit <input type="checkbox"/> Airway trolley/bronchoscope outside room <input type="checkbox"/> ETCO2 <input type="checkbox"/> Viral filter

FINAL CHECK IN ROOM

- Patient position optimal
 - Fluid runs easily
 - Suction working
 - Facemask with viral filter connected
 - ETCO2 trace
 - O2 running at 15L.min⁻¹
 - Oropharyngeal/nasal airways
- Airway plans:
 - Plan A: Videolaryngoscopy with bougie/stylet
 - Plan B: Supraglottic airway
 - Plan C: Vice grip, 2-person +/- Guedel/NPA
 - Plan D: Scalpel/bougie/tube



SAFE AIRWAY SOCIETY

COVID-19 AIRWAY MANAGEMENT

1. Intensive training
2. Early intervention
3. Meticulous planning
4. Vigilant infection control
5. Efficient airway management
6. Clear communication

USE A 'BUDDY CHECK' FOR CORRECT PPE FITTING

Planning	Intervene early - aim to avoid emergency intubation. Negative Pressure room or Normal pressure with strict door policy. Senior clinician involvement. Is Anaesthetist needed? Early airway assessment documented by senior clinician.
Prepare	Assemble 5-6 person Airway Team (see reverse). Use COVID-19 Intubation Tray (see reverse). Ensure Viral Filter and ETCO2 in ventilation circuit. Share Airway Strategy. Use a dedicated COVID intubation checklist.
PPE	Hand Hygiene (HH). Donning: HH > Gown > Mask > Eye-protection > Hat > HH > Gloves. Spotter to perform "Buddy Check" to ensure correct PPE fit. Airway operator to consider double gloves.
Pre-Ox	45 degree head up position. Pre-oxygenate with Face Mask using 2 hands for full 5 minutes. Ensure a square ETCO2 waveform, to be confident of no leaks. Avoid Apnoeic Oxygenation techniques due to aerosolization risk.
Perform	Use VL; use the screen (indirect view) to maximise operator distance from airway. Modified RSI technique (1.5mg/kg IBW Roc OR 1.5mg/kg TBW Sux). No ventilation prior to intubation unless for rescue oxygenation. Wait 60 seconds for paralysis to take effect - avoid triggering cough.
Post-ETT	Inflate cuff BEFORE initiating ventilation and monitor cuff pressures to minimise leak. Remove outer gloves (if on), dispose of airway equipment in sealed bag. Doffing: Gloves > Gown > HH > Hat > Eye Protection > Mask > HH. Use a Spotter. Debrief and share lessons.

Awake Intubation

Risk of aerosolization. Involve Senior Anaesthetist if this airway technique is indicated.

Connection / Disconnection

Apply the viral filter directly to the ETT. Only disconnect the circuit on the ventilator side of the viral filter.

CICO Rescue

Scalpel-bougie technique to avoid aerosolization.

Collaboration between Safe Airway Society + RNS ASCAR

@SafeAirway + @RnsAscAr

v1.2 March 2020

<https://www.safeairwaysociety.org/covid19/>


Guideline Only/Not a Substitute for Clinical Judgment



SAFE AIRWAY SOCIETY

COVID-19 AIRWAY MANAGEMENT


Team Members



COVID Intubation Tray

- | | | |
|---|---|--|
| <input type="checkbox"/> Macintosh VL blade | <input type="checkbox"/> Bougie / Stylet | <input type="checkbox"/> ETCO2 |
| <input type="checkbox"/> Hyperangulated VL blade (if available) | <input type="checkbox"/> 10 mL syringe | <input type="checkbox"/> NG tube (large bore) |
| <input type="checkbox"/> Macintosh direct laryngoscope | <input type="checkbox"/> Tube tie | <input type="checkbox"/> OPA + NPA |
| <input type="checkbox"/> SGA (2nd generation) | <input type="checkbox"/> Lubricant sachet | <input type="checkbox"/> Scalpel + bougie CICO kit |
| <input type="checkbox"/> ETT (appropriate size range) | <input type="checkbox"/> Viral filter | <input type="checkbox"/> Inline suction |

Circuit Setup




Collaboration between Safe Airway Society + RNS ASCAR

@SafeAirway + @RnsAscAr

v1.2 March 2020

Tracheal intubation of critically ill adults Adapted for COVID-19



This flowchart forms part of the 2020 COVID-19 Airway Guideline for tracheal intubation. Refer to the full document for further details.

Can't Intubate, Can't Oxygenate (CICO) in critically ill adults Adapted for COVID-19

CALL FOR HELP



Declare "Can't Intubate, Can't Oxygenate"

Plan D: Front Of Neck Airway: FONA

Extend neck

Ensure neuromuscular blockade

Exclude oxygen failure and blocked circuit

Personnel and PPE

New staff **must** don full checked PPE
Most appropriate airway manager to perform FONA

Scalpel cricothyroidotomy

Equipment: 1. Scalpel (wide blade e.g. number 10 or 20)
2. Bougie (\leq 14 French gauge)
3. Tube (cuffed 5.0-6.0mm ID)

Laryngeal handshake to identify cricothyroid membrane

Palpable cricothyroid membrane

Transverse stab incision through cricothyroid membrane

Turn blade through 90° (sharp edge towards the feet)

Slide Coudé tip of bougie along blade into trachea

Railroad lubricated cuffed tube into trachea

Inflate cuff, ventilate and confirm position with capnography

Secure tube

Impalpable cricothyroid membrane

Make a large midline vertical incision

Blunt dissection with fingers to separate tissues

Identify and stabilise the larynx

Proceed with technique for palpable cricothyroid membrane as above

Post-FONA care and follow up

- Closed tracheal suction
- Recruitment manoeuvre (if haemodynamically stable)
- Chest X-ray
- Monitor for complications
- Surgical review of FONA site
- Agree airway plan with senior clinicians
- Document and complete airway alert

This flowchart forms part of the 2020 COVID-19 Airway Guideline for tracheal intubation. Refer to the full document for further details.

APPENDIX F : SAMPLE PROTOCOL FOR EXTUBATION OF COVID-19 PATIENTS*

*Adapted from University Medical Center (Las Vegas, NV)

Guidelines for Extubation of COVID-19 patients:

- Extubations require 2 HCP's one to hold the mask while the second extubates the patient.
- Whenever possible patient should be placed in negative pressure rooms, and use cube extubation device with plastic shield
- This is considered an aerosolized procedure so proper N-95 masks should be worn, along with goggles, gowns and gloves.
- Place patient at 30 degrees and place nasal cannula on patient at 5-L/M
- Suction ETT and mouth prior to deflating the cuff
- Loosen ETT holder and place anesthesia face mask with HEPA filter attached over the patients nose and mouth leaving space for ETT exiting under the face mask
- IF anesthesia bag is used, use a low oxygen flow, consider attempting to extubate at end of expiration
- Deflate ETT cuff and extubate while maintaining face-mask seal
- Maintain two-handed mask seal until any immediate post-extubation coughing has subsided.
- Remove anesthesia mask and place procedure mask over the patient while wearing nasal cannula oxygen.



Place 5L nasal cannula on patient



Anesthesia mask without anesthesia bag over face-allowing ETT to exit under face mask



Anesthesia mask with anesthesia bag over face-allowing ETT to exit under face mask

APPENDIX G : TRANSPORT VENTILATOR SET UP GUIDE

Transport Vent Set Up Guide


COVID-19 Considerations – 7 April 2020

- A. A standard HME will not suffice for viral filtration. A HMEF (heat-moisture exchanger – filter) provides sufficient bacterial & viral filtration and can be used in place of an HME. If your patient does not already have an HMEF in place, place one prior to putting them on your transport ventilator. HMEFs are intended for extended use and filtration is not degraded over time. Any increase in resistance of gas flow is negligible. A HMEF that does not become visibly soiled can be used for 2-7 days.
- B. If you need to exchange the HMEF or anytime there is a circuit break without a HMEF in-line, you must clamp the ET tube.
- C. Whenever a circuit break is required all members in the area should be wearing full PPE with N95 mask or greater.

Based on availability, transport ventilators should be used with the follow order of preference:

1. Impact 731
2. Impact 754
3. Lung Transport Ventilator (LTV)
4. LP10 (not shown)
5. Hamilton T1 (only ground evac or *Rotary-wing* transport; Not flight approved for fixed or tilt-wing aircraft)
6. SAVE II


- D. Set up patient side with an HMEF for manual ventilation (below with and without accoutrements), as well as for a transport ventilator. The below three pictures are the “gold standard” for set up and NO additional filters are required.



Bag Valve device with PEEP valve. HMEF is connected between ETT and bag.




Bag Valve device with PEEP valve. In-line suction is on patient side of HMEF. EtCO₂ is between HMEF and bag.



Vent circuit. In-line suction is on patient side of HMEF. EtCO₂ is between HMEF and vent circuit.



- E. **In the event that HMEFs are not available**, the standard bacterial/viral vent filters will be needed. At a minimum, a filter must be placed on the port that entrains room air and the exhalation valve of the circuit. When disconnecting a patient from the ventilator without a HMEF, a standard bacterial/viral filter must be placed between the BVM and ET tube.

Some examples below:





For the **Impact 731**, place filters on the gas intake and exhalation valve marked by red arrows. It is important to note, that placing a filter on the gas intake (top arrow) will bypass an anti-asphyxiation safety feature. If this filter becomes occluded, a “Fresh Gas Intake Failure” alarm is likely to occur. When this alarm occurs, the patient will no longer be ventilated and will need to be manually ventilated while the vent is reset.

For the **Impact 754 ventilator**, place a filter on the gas intake (top arrow) and at the exhalation valve (bottom arrow). The set up for this ventilator will look identical to that of the Impact 731. The same caution must be taken when placing a filter on the gas intake due to the same risk of blocking gas flow to the ventilator resulting in vent failure.



For the **LTV ventilator**, there are some important considerations. Filters should be placed as marked by the red arrows. It is important to understand that a filter cannot be placed where the vent entrains room air, instead a filter is placed between the vent and the beginning of the circuit (left arrow). Also, to place a filter on the exhalation valve (right arrow), you must remove the exhalation valve and place a filter between the valve on the circuit tubing.

For the **Hamilton T1 ventilator**, filters need to be placed on the inhalation and exhalation ports, conveniently located right next to each other. (Ground or Rotary-wing only)



For the **SAVE II ventilator**, 3 filters are necessary. The red arrows mark where room air is entrained into the circuit. The yellow arrow shows the exhalation valve. Not only does using this ventilator require more filters, it is also not ideal for managing mechanically ventilated patients requiring complex ventilator settings.

APPENDIX H : POST-COVID-19 CARDIOPULMONARY RETURN TO EXERCISE RECOMMENDATIONS

ASYMPTOMATIC COVID-19 infection:

- a. Service Member (SM) has completed 10-14 day isolation with exercise limitations. May return to duty and deploy 14 days after onset of symptoms AND meets DOD Force Health Protection Guidelines Criteria for Redeployment.¹
- b. Physician or Healthcare professional has determined SM is low-risk based on:
 - 1) Focused clinical exam without clinically significant findings
 - 2) Absence of cardiac symptoms
 - 3) Confirmation of no exercise limitations or treatment needed
- c. Electrocardiogram (ECG) not required for asymptomatic infection.
- d. If ECG is obtained and demonstrates abnormalities,² recommend further evaluation with transthoracic echocardiogram and consideration for graded exercise testing before resuming training.
- e. SM should receive an exercise prescription to gradually re-acclimatize to activity over one week. Repeat medical evaluation is only necessary if symptoms develop.

NON-HOSPITALIZED, SYMPTOMATIC COVID-19 infection:

- a. SM has completed 10-14 day isolation with exercise limitations for 2 weeks after symptom resolution.
- b. Physician or Healthcare Professional has determined SM is low-risk based on:
 - 1) Clinical exam without clinically significant findings
 - 2) Absence of residual cardiopulmonary symptoms
 - 3) ECG without abnormalitiesⁱ
 - 4) No evidence of myocardial injuryⁱⁱ
 - 5) No abnormalities on Transthoracic Echocardiogram (TTE)ⁱⁱⁱ
- c. SM should receive an exercise prescription to gradually re-acclimatize to activity over 2 weeks and undergo a repeat evaluation with recurrence of any symptoms.
- d. If evaluation demonstrates evidence of myocardial injury and/or abnormal cardiac study, follow recommendations for symptomatic COVID-19 infection with myocardial injury.

HOSPITALIZED, SYMPTOMATIC COVID-19 infection WITHOUT MYOCARDIAL INJURY:

- a. Defined as a hospital course without cardiopulmonary findings suggestive of myocardial injury (absence of chest pain/pressure, dyspnea, orthopnea, arrhythmias/palpitations, syncope, or signs of heart failure in the absence of abnormal cardiac biomarkers, or abnormal POCUS/ TTE/ cardiac CT / coronary angiogram).
- b. SM has completed 14 day isolation and exercise limitations after discharge.
- c. Physician or Healthcare Professional has determined SM is low-risk based on:
 - 1) Clinical exam without clinically significant findings
 - 2) In the absence of cardiopulmonary symptoms, only a 12-lead ECG without abnormalities is recommended.²
 - 3) In the presence of cardiopulmonary symptoms (defined as angina or angina equivalent chest pain, limiting dyspnea, orthopnea, palpitations, syncope or near syncope), the following are recommended:
 - i. ECG without abnormalitiesⁱ
 - ii. No evidence of myocardial injuryⁱⁱ
 - iii. No abnormalities on TTEⁱⁱⁱ
 - iv. Confirmation of no ongoing exercise limitations or further treatment needed
- d. SM should receive an exercise prescription to gradually re-acclimatize to activity over at least 2-4 weeks and undergo a repeat evaluation with recurrence of any symptoms.
- e. If evaluation demonstrates evidence of myocardial injury and/or abnormal cardiac study follow recommendations for symptomatic COVID-19 infection with myocardial injury.

HOSPITALIZED, SYMPTOMATIC COVID 19 infection WITH MYOCARDIAL INJURY:

Clinical Management of COVID-19, v4

- a. Defined as a hospital course with cardiopulmonary findings suggestive of myocardial injury (presence of chest pain/pressure, dyspnea, orthopnea, arrhythmias, syncope, or signs of heart failure in the presence of abnormal cardiac biomarkers, or abnormal cardiac biomarkers alone, or abnormal POCUS / TTE / cardiac CT / coronary angiogram):
- b. SM has completed 10-14 day isolation after discharge.
- c. No exercise activity for 3 to 6 months **after symptom resolution**.
- d. Must complete the following evaluation before resuming exercise:
 - 1) 12 lead ECG
 - 2) Troponin I or High Sensitivity Troponin T/I
 - 3) Natriuretic peptide (BNP or NT-pro BNP)
 - 4) Other supplemental studies to show resolution of COVID sequela and demonstrate normalization of end organ function (i.e. CXR, ESR, CRP)
 - 5) Transthoracic Echocardiogram (after completion of activity restriction)
 - 6) 2 week ambulatory cardiac event monitoring
 - 7) Cardiac MRI with late gadolinium enhancement (LGE)
 - 8) Graded Exercise stress test after completion of the tests above if no abnormalities
 - 9) Cardiology consultation.
- e. SM should receive an exercise prescription to gradually re-acclimatize to activity over 2 -6 weeks after being cleared to resume exercise in the presence of low risk findings. SM should undergo a repeat evaluation with recurrence of any symptoms.

HOSPITALIZED, SYMPTOMATIC COVID-19 infection complicated by stroke, deep venous thrombosis, respiratory failure, cardiac failure, renal failure, other end-organ failure:

- a. RTD based on expert consultation and case-by-case consideration for retention vs. referral to DES/MEB.
- b. Initial Entry Training (IET, first 180-days), such as recruit training case-by-case consideration, likely unfavorable for retention waiver, unlikely to resume training for if in a recruit cycle. If retention waiver not approved, return home (DES referrals where appropriate). May reapply through formal channels with resolution of any complications (e.g., resolved myocarditis with normalized left ventricular ejection fraction).
- c. Military & Merchant Marine Academies (M&MMA) and Reserve Officer Training Corps (ROTC): case-by-case consideration.

ⁱ ECG findings that may indicate viral induced myocardial injury include: pathological Q waves, ST segment depressions, (new) diffuse ST segment elevation, and T wave inversions that are outside of the normal parameters based on the Internal Recommendations For Electrocardiographic interpretation in athletes.

ⁱⁱ Cardiac Biomarkers indicative of myocardial injury: >99th percent upper limit of normal levels for Troponin I or High Sensitivity Troponin I/ T.

ⁱⁱⁱ Transthoracic echocardiogram findings of cardiac injury- regional wall motion abnormalities, dilated ventricles, abnormal systolic function with a reduced EF <45%

References:

1. Jones, Steve. Force Health Protection Guidance (Supplement 9)- Department of Defense Guidance for Deployment and Redeployment of Individuals and Units During the Novel Corona Virus Disease 2019 Pandemic. Published online May 26, 2020. Accessed May 29, 2020. <https://www.health.mil/Military-Health-Topics/Combat-Support/Public-Health/Coronavirus>
2. Sharma S, Drezner JA, Baggish A, et al. International Recommendations for Electrocardiographic Interpretation in Athletes. *J Am Coll Cardiol*. 2017;69(8):1057-1075. doi:10.1016/j.jacc.2017.01.015


APPENDIX I : WEIGHT-BASED HEPARIN DOSING ALGORITHM FOR VENOUS THROMBOEMBOLISM

Weight-Based Heparin Dosing for Venous Thromboembolism, anti-Xa goal 0.3-0.7	
Initial Therapy	
Bolus ^a	80 units/kg
Infusion ^a	18 units/kg/hr
Adjustments^b	
Anti-Xa <0.2	Increase by 4 units/kg/hr
Anti-Xa 0.2-0.29	Increase by 2 units/kg/hr
Anti-Xa 0.3-0.7	No Change
Anti-Xa 0.71-0.8	Decrease by 1 unit/kg/hr
Anti-Xa 0.81-0.9	Hold for 0.5 hr; Decrease by 2 units/kg/hr
Anti-Xa >0.9	Hold for 1 hr; Decrease infusion by 3 units/kg/hr
(maximum 5,000 units), and typical initial infusion dose is 12 units/kg/hr (maximum 1,000 units/hr).	
^a Round all doses to nearest 100 units.	
^b Draw Anti-Xa 6 hours after STARTING therapy and 6 hours after any CHANGE in infusion rate.	
Adapted from https://journals.sagepub.com/doi/pdf/10.1345/aph.1Q161	

APPENDIX J : ENTERAL NUTRITION CARE PATHWAY FOR PATIENTS WITH COVID-19

Enteral Nutrition (tube feeding) Care Pathway for Critically-Ill Adult Patients Diagnosed with COVID-19

This pathway provides steps and resources for managing critically-ill adult patients (pts) requiring enteral nutrition (EN).




References:




1. McClave, SA, Taylor BE, Martindale RD, et al; Society of Critical Care Medicine; American Society for Parenteral and Enteral Nutrition. Guidelines for the provision and assessment of nutrition support therapy in the adult critically ill patient: Society of Critical Care Medicine (SCCM) and American Society for Parenteral and Enteral Nutrition (ASPEN). *JPEN J Parenter Enteral Nutr.* 2016;40:159-211.
2. Martindale R, Patel JJ, Taylor B, Warren M, & McClave SA. Nutrition Therapy in the Patient with COVID-19 Disease Requiring ICU Care. Reviewed and approved by ASPEN and Society for Critical-Care Medicine. 1 April 2020. https://www.nutritioncare.org/Guidelines_and_Clinical_Resources/Resources_for_Clinicians_Caring_for_Patients_with_Coronavirus/
3. Abbott Laboratories. Guideline to Adult Tube Feeding. November 2018. https://static.abbottnutrition.com/cms-prod/abbottnutrition-2016.com/img/182393-Adult-tube-feeding-home_tcm1226-57870.pdf

APPENDIX K : SAMPLE PROTOCOLS FOR VARIOUS ICU MANAGEMENT

Intubation





Mechanical Ventilation




COVID Prone

COVID Rapid Response Team






COVID Code Blue - Roles & Responsibilities



APPENDIX L : AHA ACLS & PALS CARDIAC ARREST ALGORITHM FOR COVID-19 PATIENTS

ACLS Cardiac Arrest Algorithm for Suspected or Confirmed COVID-19 Patients

Updated April 2020



© 2020 American Heart Association

CPR Quality

- Push hard (at least 2 inches [5 cm]) and fast (100-120/min) and allow complete chest recoil.
- Minimize interruptions in compressions.
- Avoid excessive ventilation.
- Change compressor every 2 minutes, or sooner if fatigued.
- If no advanced airway, 30:2 compression-ventilation ratio.
- Quantitative waveform capnography
 - If PETCO₂ <10 mm Hg, attempt to improve CPR quality.
- Intra-arterial pressure
 - If relaxation phase (diastolic) pressure <20 mm Hg, attempt to improve CPR quality.

Shock Energy for Defibrillation

- **Biphasic:** Manufacturer recommendation (eg, initial dose of 120-200 J); if unknown, use maximum available. Second and subsequent doses should be equivalent, and higher doses may be considered.
- **Monophasic:** 360 J

Advanced Airway

- **Minimize closed-circuit disconnection**
- **Use intubator with highest likelihood of first pass success**
- **Consider video laryngoscopy**
- Endotracheal intubation or supraglottic advanced airway
- Waveform capnography or capnometry to confirm and monitor ET tube placement
- Once advanced airway in place, give 1 breath every 6 seconds (10 breaths/min) with continuous chest compressions

Drug Therapy

- **Epinephrine IV/IO dose:** 1 mg every 3-5 minutes
- **Amiodarone IV/IO dose:** First dose: 300 mg bolus. Second dose: 150 mg.
or
Lidocaine IV/IO dose: First dose: 1-1.5 mg/kg. Second dose: 0.5-0.75 mg/kg.

Return of Spontaneous Circulation (ROSC)


- Pulse and blood pressure
- Abrupt sustained increase in PETCO₂ (typically ≥40 mm Hg)
- Spontaneous arterial pressure waves with intra-arterial monitoring

Reversible Causes

- Hypovolemia
- Hypoxia
- Hydrogen ion (acidosis)
- Hypo-/hyperkalemia
- Hypothermia
- Tension pneumothorax
- Tamponade, cardiac
- Toxins
- Thrombosis, pulmonary
- Thrombosis, coronary

Pediatric Cardiac Arrest Algorithm for Suspected or Confirmed COVID-19 Patients

Updated April 2020



© 2020 American Heart Association

CPR Quality

- Push hard ($\geq\frac{1}{3}$ of anteroposterior diameter of chest) and fast (100-120/min) and allow complete chest recoil.
- Minimize interruptions in compressions.
- Avoid excessive ventilation.
- Change compressor every 2 minutes, or sooner if fatigued.
- If no advanced airway, 15:2 compression-ventilation ratio.

Shock Energy for Defibrillation

First shock 2 J/kg, second shock 4 J/kg, subsequent shocks ≥ 4 J/kg, maximum 10 J/kg or adult dose

Advanced Airway

- **Minimize closed-circuit disconnection**
- **Use intubator with highest likelihood of first pass success**
- **Consider video laryngoscopy**
- **Prefer cuffed endotracheal tube if available**
- Endotracheal intubation or supraglottic advanced airway
- Waveform capnography or capnometry to confirm and monitor ET tube placement
- Once advanced airway in place, give 1 breath every 6 seconds (10 breaths/min) with continuous chest compressions

Drug Therapy

- **Epinephrine IO/IV dose:** 0.01 mg/kg (0.1 mL/kg of the 0.1 mg/mL concentration). Repeat every 3-5 minutes.
- **Amiodarone IO/IV dose:** 5 mg/kg bolus during cardiac arrest. May repeat up to 2 times for refractory VF/pulseless VT.
- or
- **Lidocaine IO/IV dose:** Initial: 1 mg/kg loading dose. Maintenance: 20-50 mcg/kg per minute infusion (repeat bolus dose if infusion initiated >15 minutes after initial bolus therapy).

Return of Spontaneous Circulation (ROSC)

- Pulse and blood pressure
- Spontaneous arterial pressure waves with intra-arterial monitoring

Reversible Causes

- Hypovolemia
- Hypoxia
- Hydrogen ion (acidosis)
- Hypoglycemia
- Hypo-/hyperkalemia
- Hypothermia
- Tension pneumothorax
- Tamponade, cardiac
- Toxins
- Thrombosis, pulmonary
- Thrombosis, coronary

APPENDIX M : PREPARATION AND CLEANING OF ULTRASOUND ROOMS IN THE CONTEXT OF COVID-19

Reference: ISUOG Safety Statement: Safe performance of obstetric and gynecological scans and equipment cleaning in the context of COVID-19

Ultrasound units:

Preparation and cleaning of ultrasound room for all patients

The survival of severe acute respiratory syndrome (SARS)-associated viruses (including COVID-19) on dry inanimate surfaces, such as ultrasound systems, is between 48 and 96 h. The ultrasound room should be cleaned thoroughly each morning and all contents should be wiped with a compatible low-level disinfectant (LLD), including the ultrasound monitor, computer keyboard and mouse, stretcher rails, transducer holder, gel container, door handles, cabinet knobs, light switches, chairs and counter tops.

- The number of transducers connected to the ultrasound machine should be reduced to a minimum, usually one transabdominal and one transvaginal, and all other transducers should be stored safely in a clean closed cabinet and brought out as needed.
- All unnecessary accessories in the room should be removed and, where possible, stored in the cabinets.
- Fabric-covered chairs should be replaced with hard-surface chairs that can be wiped.
- Where possible, replace all washable linen, such as towels, pillow covers and sheets, with disposable covers.
- Ultrasound transducers and cables should be cleaned (as recommended below) every morning and this should also be performed after each scan.
- The patient bed or couch should be wiped with a LLD prior to replacing the disposable paper cover.
- The disposable paper cover should be removed with gloved hands and folded and disposed of immediately at the end of each examination.
- Ensure that the highly touched surfaces (e.g. keyboard, cord and screen) of the ultrasound machine are thoroughly cleaned after each examination.
- At the end of the day, soiled linen should be handled using two pairs of gloves and disposed of in the appropriate container without shaking the linen. The room and equipment should undergo terminal cleaning using a LLD. Hands should be washed for 20 sec afterwards.

Preparation and cleaning of ultrasound equipment for all patients

The transducer and ultrasound equipment must be cleaned with a compatible LLD after each patient, in accordance with local guidelines.

Preparation and cleaning of ultrasound equipment after performing an examination in a suspected or confirmed COVID-19 case

SARS coronavirus, Middle East respiratory syndrome (MERS) coronavirus and endemic human coronaviruses (HCoV) can persist on inanimate surfaces, such as metal, glass or plastic, for up to 9 days, but can be efficiently inactivated by surface disinfection procedures with 62–71% ethanol, 0.5% hydrogen peroxide or 0.1% sodium hypochlorite within 1 min. Other biocidal agents often used include 0.05–0.2% benzalkonium chloride (Clinell TM) or 0.02% chlorhexidine digluconate.

- Check the required contact time for each product.

Since information about COVID-19 is incomplete, additional use of high-level disinfectants is recommended; however, this advice is manufacturer-specific. High-level disinfectants include ethanol 80-95% (exposure time 30 sec), 2-propanol 75-100% (exposure time 30 sec), 2-propanol and 1-propanol 45% and 30% (exposure time 30

Clinical Management of COVID-19, v4

sec), sodium hypochlorite 0.21% (Antisapril Blu 2%, exposure time 30 sec), glutaraldehyde 2.5% (exposure time 5 min) and 0.5% (CIDEX OPA, exposure time 2 min), hydrogen peroxide 0.5% (REVITAL-OX RESERT, exposure time 1 min)⁶. Protective eyewear and gloves should be used when cleaning, disinfecting or sterilizing any equipment.

Preparation and cleaning of ultrasound transducer after performing an examination in a suspected or confirmed COVID-19 case

If feasible, it is recommended to have one (or more) dedicated ultrasound machine(s) for patients with suspected/probable/confirmed COVID-19 infection.

- If the patient must be scanned in the clinic, this should be done at the end of the clinic list, as the room and equipment will subsequently require a deep clean.
- It is imperative to perform hand hygiene once the gloves have been removed.

Guidelines regarding cleaning of ultrasound transducers between patients are available. Coronaviruses are enveloped viruses, which are the least resistant to inactivation by disinfection. The structure of these viruses includes a lipid envelope, which is easily disrupted by most disinfectants suitable for use on ultrasound systems and transducers.

According to the Spaulding classification system, medical devices are classified according to the infection risk they present as non-critical, semi-critical and critical (also referred to as low-risk, medium-risk and high-risk). Non-critical devices present the lowest risk for infection as they come in contact with intact skin, such as transabdominal transducers. Low- or intermediate-level disinfection is recommended, which will eradicate most bacteria (but not bacterial spores) and fungi, as well as certain types of viruses, including human immunodeficiency virus (HIV). Semi-critical devices are those that present a higher risk for infection because of contact with non-intact skin or mucous membranes. Transvaginal transducers belong to this category. High-level disinfection for destruction of all microorganisms, including COVID-19, is recommended and can be performed by means of solutions containing sodium hypochlorite or other disinfectants as detailed above. Critical devices, such as transducers used in invasive procedures, must undergo sterilization as per medical facility guidelines irrespective of whether a probe cover is used.

Preparation of the ultrasound transducer consists of two steps: cleaning and disinfection. Any products used for cleaning or disinfection must be compatible with the ultrasound equipment, as determined by the ultrasound equipment manufacturer. Certain products may damage ultrasound equipment or transducers and invalidate warranties.

1. Cleaning

This is an important first step since any remaining gel can act as a barrier to the disinfectant thus diminishing its efficacy. The USA Centers for Disease Control and Prevention (CDC) defines cleaning as ‘the removal of foreign material (e.g., soil, and organic material) from objects and is normally accomplished using water with detergents or enzymatic products’. Ineffective cleaning prior to disinfection can limit the effectiveness of chemical disinfection.

Current guidelines for cleaning transvaginal transducers recommend using running water to remove any residual gel or debris from the probe before cleansing thoroughly the transducer using a damp gauze pad, or other soft cloth, and a small amount of mild nonabrasive liquid soap (approved for use on medical instruments). The use of a small brush especially for the crevices and areas of angulation should be considered, depending on the design of the particular transducer. The transducer should then be rinsed thoroughly with running water and dried with a soft cloth or paper towel.

Clinical Management of COVID-19, v4

Based on the above guidelines, the following steps are recommended for cleaning the transducer, which should be performed wearing disposable gloves:

- a. Disconnect the transducer.
- b. Remove the transducer cover (if applicable) and dispose of in clinical waste.
- c. Rinse the operative end of the transducer with running tap water (NOT the electronic contact end).
- d. Clean the transducer with a soft brush and nonabrasive detergent.
- e. Rinse the transducer with tap water.
- f. Clean the transducer cable with a LLD wipe.
- g. Dry with a cloth or towel (residual water can dilute chemical disinfectants, if this is the preferred method).

2. Disinfection

Always refer to your facility's infection control policies and protocols, as well as the transducer manufacturer's instruction for use and labels for use. Disinfection practices are evolving constantly, and this is the most current to date. As mentioned above, high-level disinfection is recommended for transvaginal but not transabdominal transducers. Specific product instructions must be consulted. Available methods (current at the time of publication) include:

Chemical 'wet' disinfection:

- 2.4–3.2% glutaraldehyde products (such as Cidex, Metricide and Procide).
- Non-glutaraldehyde agents (such as Cidex OPA (o-phthalaldehyde) and Cidex PA (hydrogen peroxide and peroxyacetic acid)).
- Approved multistep disinfectant wipes containing chlorine dioxide, which are used extensively in the UK and Australia (Tristel Duo®).
- 7.5% hydrogen peroxide solution, which works by producing destructive hydroxyl free radicals.
- Sodium hypochlorite 0.21% (Antisapril Blu 2%).


Note that common household bleach (5.25% sodium hypochlorite) diluted to yield 500 parts per million chlorine (10 cc in one L of tap water), although effective it is not recommended by manufacturers because it can potentially cause damage to metal and plastic parts of the transducer. Mention of this disinfectant here does not imply that we consider it to be appropriate, but we are aware that it is used in some settings.

Automated high-level disinfection:

- Antigermix (Germitec, France): the transducer is placed in a closed cabinet and exposed to high-intensity ultraviolet type C radiation.
- Astra VR (CIVCO Medical Solutions, USA): automated disinfection with Cidex OPA and Metricide solutions.
- Trophon (Nanosonics, Australia): sonicated hydrogen peroxide mist.

After cleaning, store transducer in a clean closet if it is not going to be reused immediately.

APPENDIX N : DHA SUMMARY ON MULTISYSTEM INFLAMMATORY SYNDROME IN CHILDREN (MIS-C)



The Centers for Disease Control and Prevention (CDC) and the MHS are tracking instances of a life-threatening pediatric condition that appears to be occurring in patients who were diagnosed with or exposed to COVID-19. Early detection and treatment of this condition, called multisystem inflammatory syndrome in children (MIS-C), is crucial.

Though MIS-C appears to be a rare complication, the MHS Clinical Communities want to raise awareness so that any MHS staff member can recognize a patient with symptoms of MIS-C and knows how to react. This could be staff conducting intake at urgent care clinics or emergency departments, and nursing and reception staff at family medicine clinics.



What It Is and Who is Affected

The CDC has defined the syndrome and it is reported among patients with a variety of symptoms.

The CDC defines this syndrome as:

- An individual aged <21 years presenting with fever, laboratory evidence of inflammation, and evidence of clinically severe illness requiring hospitalization, with multisystem (>2) organ involvement (cardiac, renal, respiratory, hematologic, gastrointestinal, dermatologic, or neurological); **AND**
- NO alternative plausible diagnoses; **AND**
- Positive for current or recent severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection by reverse transcription polymerase chain reaction (RT-PCR), serology, or antigen test; **OR** COVID-19 exposure within the 4 weeks prior to the onset of symptoms

Note that emerging reports indicate that MIS-C may be affecting young adults over the age of 21. The MHS Clinical Communities recommend not limiting your assessment of these conditions to only patients under the age of 21.

MIS-C shares symptoms with other pediatric inflammatory conditions including Kawasaki disease, toxic shock syndrome, and macrophage activation syndrome.

It is critical for healthcare professionals to recognize syndrome symptoms early. This includes primary care physicians, pediatricians, emergency room staff, urgent care staff, and all support staff.



What to Look For

Child or young adult with:

- Fever of 100.4 °F for 4 or more days
- Abdominal pain, nausea, vomiting, diarrhea, or enteritis only on imaging
- Neck pain
- Rash
- Pink eye, bloodshot eyes
- Oral/mucosal changes
- Headache/irritability
- Cough, sore throat, pain swallowing
- Swelling in hands and/or feet
- Trouble breathing
- Low energy, tired



What to Do

It is important for staff to be prepared to handle suspected cases of MIS-C.

Staff should prepare in advance by:

- Familiarizing themselves with the signs of MIS-C
- Collecting the contact information for local military or civilian pediatric specialists
- Utilizing existing DoD specialty telemedicine resources to reach out early to specialists for advice/consultation
- Understanding transportation options for transfer of MIS-C patients to specialty centers

Health care management should include:

- Familiarizing staff, especially providers and hospital and clinic intake staff, with the signs of MIS-C
- Ensuring all health care workers are wearing appropriate personal protective equipment (PPE) before examining patients
- Contacting the nearest hospital with pediatric inpatient capabilities, to including a pediatric intensive care unit (PICU) and pediatric cardiology at a minimum, plus pediatric infectious diseases, pediatric rheumatology, or immunology capabilities
- If the patient is clinically unstable, activating the emergency response system and following Pediatric Advanced Life Support algorithms
- If the patient is clinically stable and is in an emergency or tertiary care center, obtaining the 12-lead EKG and labs recommended in clinical guidance. MHS clinical guidance for MIS-C will be available in Version 4 of the *DoD COVID-19 Practice Management Guide*
- For facilities in a remote location without subspecialty care, contacting pediatric specialists via phone for guidance on management/stabilization until able to transfer patient to higher level of care

Providers should report suspected cases to their local, state, or territorial health department.



Who to Contact

In addition to learning how to recognize MIS-C and preparing to manage cases, staff should know who to reach out to for additional guidance.

For questions about MIS-C and its management contact:

MHS Pediatric Tele-Critical Care: 833-238-7756, DSN 312-429-9089
 CDC's 24-hour Emergency Operations Center: 770-488-7100

For more information on MHS guidance, visit health.mil/coronavirus; for CDC information on MIS-C, visit <https://www.cdc.gov/mis-c/hcp>
 To read the May 14, 2020 CDC Health Advisory on MIS-C, visit <https://emergency.cdc.gov/han/2020/han00432.asp>.

June 1, 2020



APPENDIX O : DHA QUICK REFERENCE GUIDE TO VIRTUAL HEALTH AND TELEPHONE ENCOUNTERS

QUICK REFERENCE GUIDE

Virtual Health and Telephone Encounters



Disclaimers:

1. The **MHS Specific Coding Guidelines** are the source for all DoD specific coding guidance. To ensure coding compliance and standardization across the DHA, all coding training, tools and/or coding manuals must be vetted through **DHA Medical Coding Program Branch** and **DHA Coding Work Group (CWG)** prior to implementation.
2. The codes listed in this document are for Type 1 or 2 Privileged Providers. Refer to the **MHS Specific Coding Guidelines** for codes related to all other provider types.
3. Effective 1 Oct 2018, CMS eliminated the use of modifier GT for reporting Telehealth professional services, however, the MHS will continue to use GT until further notice.
4. Online Secure Messaging Encounters - 2019 CPT code 99444 was deleted and replaced with 3 new codes for 2020: 99421, 99422 and 99423.
Note: Use 99444 until the 2020 CPT tables are updated at your MTF.
5. VH Functionals have ownership of this VH Coding Cheat Sheet and will update annual code changes in conjunction with the **MHS Coding Guidelines**.
6. VH and Telephone Encounters must meet coding and documentation requirements.

Method	Virtual Health (Privileged Provider to Patient)				Virtual Health (Privileged Provider to Provider)		Audio-Only Encounters			Online Messaging Encounters
Interaction	Virtual Video Visit (V3) DISTANT SITE (where provider is located e.g. MTF)		Virtual Video Visit (V3) ORIGINATING SITE (where patient is located e.g. MTF)		Teleconsultation at distant site (where provider is located at e.g. MTF or on a ship) for professional interpretation	Teleconsultation at distant site (where provider is located at e.g. different MTF or on ship) for diagnosis and treatment	Clinical Telephone (audio only)		T-CON	Secure Messaging
PHA	✓		✓				✓			
Synchronous		✓		✓			✓	✓	✓	
Asynchronous					✓	✓				✓
Type: Scheduled		✓		✓			✓			
Type: Unscheduled					✓	✓			✓	✓
Appointment	FTR	24HR, FTR, SPEC (provider discretion)	FTR	24HR, FTR, SPEC	SPEC, FTR		24HR, FTR, SPEC	24HR, FTR, SPEC	T-CON	N/A
Detail Code	PHA and TELMED	TELMED	PHA and TELMED	TELMED	N/A	N/A	HC		N/A	N/A
Code	Office Visit Code		99499	99492	99451 ≥5 min consultant time including written report 99452 30 min requesting provider time in referral work	Verbal and Written Report: 99446 5-10 min 99447 11-20 min 99448 21-30 min 99449 +31 min	99499	99441 5-10 min 99442 11-20 min 99443 21-30 min		For 2019: 99444 For 2020: 99421 5-10 min 99422 11-20 min 99423 21+ min
Modifier	GT (patient is present at originating site e.g. clinic to clinic) or 95 (clinic to non-clinic patient location)		N/A	N/A	N/A	GC	N/A	N/A	N/A	GC
Code	96160	N/A	O3014 (and any procedures performed at the originating site within presenter's scope of practice)		Procedures per documentation	N/A	96160 + G2012	G2012	N/A	N/A
Modifier	N/A	N/A	N/A	TC (if technical component of procedure/test are completed and no interpretation and report are rendered)	26 (interpretation only; professional component)	N/A	N/A	N/A	N/A	N/A

Effective Date: 03/30/2020

APPENDIX P : EXAMPLE TRIAGE PROTOCOL FOR RESOURCE ALLOCATIONS IN TIMES OF CRISIS

TRIAGE PLANNING COMMITTEE	
<p><u>Members (Minimum)</u></p> <p>2 Senior Clinicians Senior Nursing Representative Ethics Representative Community Member Pastoral Care Palliative Care (as available)</p>	<p><u>Roles</u></p> <p>-Planning for the greatest medical benefit to greatest number of people -Establish SOPs for conventional, contingency and crisis capacity -Provide oversight support of scarce resource allocation decisions -Maintain available representative 24/7 to triage teams and command -Seek opportunities for regionalization of resources as permissible</p>
TRIAGE TEAM	
<p><u>Members</u></p> <p>Triage Officer (Senior Clinician) Acute Care Nurse Administrative Staff Member Ethics Representative (as available) Community Member (as available)</p>	<p><u>Roles</u></p> <p>-Liaison with command and planning committee on resources (ICU beds, staffing, equipment) -Initial contact with clinical teams for assessment of priority scoring -Collect only information relevant to priority scoring and maintain database -Make urgent allocation decisions within 90 minutes of clinical team request -Meet twice daily to match resources to patient needs and -Reassess patients every 72 hours (Minimum) -Report conflicts or requests for appeal/oversight to Planning Committee representative</p>

Figure 1: Example of Triage Team Composition and Roles

Example of Triage Oversight/Appeal Board (Individual to Each MTF)

Committee (≥3 for quorum)

- Chief medical officer
- Chief nursing officer
- Planning Committee Rep
- Legal counsel
- Risk management
- Chair of ethics committee
- Off-duty triage officer

Figure 2. Sample Priority Scoring Method

Priority Score (37) = SOFA score (24) + Clinical Frailty Score (9) + Comorbidity Score (4)
 (Numbers in parenthesis are maximum scores)

(Calculators Below)

*For Pediatric patients use PSOFA

Patient's Score = SOFA + Frailty Score + Comorbidity Score

SOFA score	1	2	3	4
Respiration				
PaO ₂ /FiO ₂ mmHg	< 400	< 300	< 200 with respiratory support	< 100 with respiratory support
Coagulation				
Platelets x 10 ³ /mm ³	< 150	< 100	< 50	< 20
Liver				
Bilirubin, mg/dL (µmol/L)	1.2–1.9 (20–32)	2.0–5.9 (33–101)	6.0–11.9 (102–204)	> 12.0 (> 204)
Cardiovascular				
Hypotension ^a	MAP < 70 mm Hg	Dopamine ≤ 5 or Dobutamine (any dose)	Dopamine < 5 or epinephrine ≤ 0.1 or norepinephrine ≤ 0.1	Dopamine > 1.5 or epinephrine > 0.1 or norepinephrine > 0.1
Central Nervous System				
Glasgow coma score	13–14	10–12	6–9	< 6
Renal				
Creatinine, mg/dL (µmol/L) or urine output	1.2–1.9 (110–170)	2.0–3.4 (171–299)	3.5–4.9 (300–440) or < 500 mL/day	> 5.0 (> 440) or < 200 mL/day

^a adrenergic agents administered for at least one hour (doses given are in µg/kg · min)

SOFA Score Table for Scoring below¹

Variable	Physiologic Value	Score
PaO ₂ /FiO ₂ Ratio		
SaO ₂ /FiO ₂ Ratio		
Platelet Count		
Bilirubin		
Blood Pressure		
Vasopressor dose	None or Agent/Dose:	
GCS		
Creatinine		
TOTAL SOFA SCORE		

Glasgow Coma Score (used for SOFA scoring above)

Criteria	Action	Score	Criteria Score
Best Eye Response	No eye opening	1	
	Opens to painful stimuli	2	
	Opens to verbal command	3	
	Opens spontaneously	4	
Best Verbal Response	None	1	
	Incomprehensible sounds	2	
	Inappropriate words	3	
	Confused	4	
	Oriented	5	
Best Motor Response	None	1	
	Extension	2	
	Flexion	3	
	Withdraw from pain	4	
	Localizes to pain	5	
	Obeys command	6	
Total Score			

Clinical Management of COVID-19, v4

Table 1 Detailed CFS criteria

CFS	Severity	Definition
1	Very fit	People who are robust, active, energetic and motivated. These people commonly exercise regularly. They are among the fittest for their age
2	Well	People who have no active disease symptoms but are less fit than category 1. Often, they exercise or are very active occasionally, e.g., seasonally
3	Managing well	People whose medical problems are well controlled, but are not regularly active beyond routine walking
4	Vulnerable	While not dependent on others for daily help, often symptoms limit activities. A common complaint is being “sloped up”, and/or being tired during the day
5	Mildly frail	These people often have more evident slowing, and need help in high older IADLs (finances, transportation, heavy housework, medications). Typically, mild frailty progressively impairs shopping and walking outside alone, meal preparation and housework
6	Moderately frail	People need help with all outside activities and with keeping house. Inside, they often have problems with stairs and need help with bathing and might need minimal assistance (cuing, standby) with dressing
7	Severely frail	Completely dependent for personal care, from whatever cause (physical or cognitive). Even so, they seem stable and not at high risk of dying (within ~6 months)
8	Very severely frail	Completely dependent, approaching the end of life. Typically, they could not recover from a minor illness
9	Terminally ill	Approaching the end of life. This category applies to people with a life expectancy <6 months, who are not otherwise evidently frail

CFS, clinical frailty scale.

Clinical Frailty Score Criteria⁶

Clinical Frailty Score (1-9)

Table to use for Comorbidity Score

Examples of Major Comorbidities	Examples of Severely Life Limiting Comorbidities
<ul style="list-style-type: none"> • Moderate dementia • Malignancy with < 10 year expected survival • New York Heart Associate Class III heart failure • Moderate to severe chronic lung disease (COPD, ILD) • End Stage Renal Disease (age < 75) • Severe Multivessel CAD • Cirrhosis with history of decompensation 	<ul style="list-style-type: none"> • Severe dementia • Cancer with only palliative treatment options • New York Heart Associate Class IV heart failure • Severe COPD with frailty • Cirrhosis with MELD > 20 • End Stage Renal Disease (age > 75)

2 Points for Major Comorbidities: Yes / No

4 Points for Severely Life Limiting Comorbidities: Yes / No

Pediatric SOFA³

Variables	Score ^a				
	0	1	2	3	4
Respiratory					
Pao ₂ :Fio ₂ ^b or SpO ₂ :Fio ₂ ^c	≥400	300-399	200-299	100-199 With respiratory support	<100 With respiratory support
	≥292	264-291	221-264	148-220 With respiratory support	<148 With respiratory support
Coagulation					
Platelet count, ×10 ³ /μL	≥150	100-149	50-99	20-49	<20
Hepatic					
Bilirubin, mg/dL	<1.2	1.2-1.9	2.0-5.9	6.0-11.9	>12.0
Cardiovascular					
MAP by age group or vasoactive infusion, mm Hg or μg/kg/min ^d					
<1 mo	≥46	<46	Dopamine hydrochloride ≤5 or dobutamine hydrochloride (any)	Dopamine hydrochloride >5 or epinephrine ≤0.1 or norepinephrine bitartrate ≤0.1	Dopamine hydrochloride >15 or epinephrine >0.1 or norepinephrine bitartrate >0.1
1-11 mo	≥55	<55			
12-23 mo	≥60	<60			
24-59 mo	≥62	<62			
60-143 mo	≥65	<65			
144-216 mo	≥67	<67			
>216 mo ^e	≥70	<70			
Neurologic					
Glasgow Coma Score ^f	15	13-14	10-12	6-9	<6
Renal					
Creatinine by age group, mg/dL					
<1 mo	<0.8	0.8-0.9	1.0-1.1	1.2-1.5	≥1.6
1-11 mo	<0.3	0.3-0.4	0.5-0.7	0.8-1.1	≥1.2
12-23 mo	<0.4	0.4-0.5	0.6-1.0	1.1-1.4	≥1.5
24-59 mo	<0.6	0.6-0.8	0.9-1.5	1.6-2.2	≥2.3
60-143 mo	<0.7	0.7-1.0	1.1-1.7	1.8-2.5	≥2.6
144-216 mo	<1.0	1.0-1.6	1.7-2.8	2.9-4.1	≥4.2
>216 mo ^e	<1.2	1.2-1.9	2.0-3.4	3.5-4.9	≥5

1. Moreno R, Vincent J-L, Matos R, et al. The use of maximum SOFA score to quantify organ dysfunction/failure in intensive care. Results of a prospective, multicentre study. *Intensive care medicine*. 1999;25(7):686-696.
2. Wallis S, Wall J, Biram R, Romero-Ortuno R. Association of the clinical frailty scale with hospital outcomes. *QJM: An International Journal of Medicine*. 2015;108(12):943-949.
3. Matics TJ, Sanchez-Pinto LN. Adaptation and validation of a pediatric sequential organ failure assessment score and evaluation of the sepsis-3 definitions in critically ill children. *JAMA pediatrics*. 2017;171(10):e172352-e172352.
4. Faden RR, Kass NE, Goodman SN, Pronovost P, Tunis S, Beauchamp TL. An ethics framework for a learning health care system: a departure from traditional research ethics and clinical ethics. *Hastings Center Report*. 2013;43(s1):S16-S27.
5. Emanuel EJ, Persad G, Upshur R, et al. Fair allocation of scarce medical resources in the time of Covid-19. In: *Mass Medical Soc*; 2020.
6. Shimura T, Yamamoto M, Kano S, Kagase A, Kodama A, Koyama Y, Otsuka T, Kohsaka S, Tada N, Yamanaka F, Naganuma T. Impact of frailty markers on outcomes after transcatheter aortic valve replacement: insights from a Japanese multicenter registry. *Annals of cardiothoracic surgery*. 2017 Sep;6(5):532.

APPENDIX Q : LIST OF CONTRIBUTORS

Contributors (*Denotes Section Editor)		
<p>*Lt Col Wesley M. Abadie, MC, USAF SMSgt Britton Adams, USAF LDCR Melissa Amescua, USN Maj David M. Anderson, MC, USAF Col Michelle K. Anton, MC, USAF Maj Bracken Armstrong, MC, USAF 1st Lt Chelsea Baker, BSC, USAF COL David M. Benedek, MC, USA *CDR John Benjamin, MC, USN COL Janetta Blackmore, USA *LTC Matthew A. Borgman, MC, USA Maj Sarah Bowe, MC, USAF Lt Col Saunya N. Bright, BSC, USAF Laura R. Brosch, RN, PhD Lt Col Daniel Brown, MC, USAF Robert F. Browning, MD *COL Jessica L. Bunin, MC, USA *CAPT Timothy H. Burgess, MC, USN *Maj Denise Campbell, BSC, USAF Ms. Elizabeth Campbell, RN, CIC, FAPIC *COL Andre Cap, MC USA LTC Vincent Capaldi, MC, USA Col Francis R. Carandang, MC, USAF Maj Mark Cheney, MC, USAF Col Jennifer C. Chow, MC, USAF Mr. David Christiansen Julia Christopher, DO LTC Jacob Collen, MC, USA *LTC Christopher Colombo, MC, USA Ms. Polyxeni Combs, MS CAPT Meghan L. Corso, USPHS *Ms. Helen Crouch, MPH, CIC, FAPIC COL Cord Cunningham, MC, USAR *CAPT Konrad L. Davis, MC, USN *LTC Sally DelVecchio, MC, USA Ms. Linderpal Dhillon, USAF COL Nicole Dobson, MC, USA COL Sean N. Dooley, MC, USA *J. Stephen Dumler, MD *COL Jeremy Edwards, MC, USA CAPT Eric Elster, MC, USN CAPT Bridgette Faber, USN *MAJ Emilio Fentanes, MC, USA CDR Jamie Fitch, MC, USN Ms. Sarah Flash, DHA Mr. Christopher Florez, MPH, CIC LTC Shannon C. Ford, MC, USA MAJ Brian Foster, MC, USA Mr. Al Garza, PE *Lt Col Brian J. Gavitt, MC, USAF Dr. James Giordano, PhD Mr. Joshua Girton, JD, LL.M, MBA LTC Pascale Guirand, NC, USA CDR Reina Gomez, USN Mark Haigney, MD *LTC Mitchell Hamele, MC, USA MAJ William Harner, MC, USA</p>	<p>Ms. Erica Harper, RN, MSN COL Bonnie H. Hartstein, MC, USA Maj Alison Helfrich, MC, USAF COL Brian A. Hemann, MC, USA Maj Minette Herrick, USAF Ms. Jennifer Hesch, JD *COL Patrick Hickey, MC, USA LTC Sean Hipp Mr. Roy Hirschak Vincet B. Ho, MD, MBA Charles W. Hoge, MD Col Melissa Howard, USAF Col David Hsieh, MC, USAF *MAJ Jessica Hull, MC, USA CDR Ashley E. Humphries, MC, USN Maj John Hunninghake, MC, USAF MAJ Nikhil A. Huprikar, MC, USA Col (Ret) Mylene T. Huynh, MD, MPH MSgt Ashley Jackson, USAF LTC(P) Chester Jean, MC, USA CAPT Elliot Jessie, MC, USN CDR Shane Jensen, MC, USN COL Matthew R. Jezior, MC, USA CDR Michael J. Kavanaugh, MC, USN Maj Rohin Kasuda, MC, USAF *Lt Col Jeremy Kilburn, MC, USAF Thomas D. Kirsch, MD, MPH *LTC Lisa Konitzer, USA CDR Joseph Kotora, MC, USN MAJ Benjamin Krainin, MC, USA LTC Jeffrey Laczek, MC, USA Lt Col John P. Lichtenberger III, MC, USAFR CAPT Mark G. Lieb CAPT Robert Liotta, MC, USN Maj David A. Lindholm, MC, USAF MAJ Adrianna Long, MC, USA *COL Frederick Lough, MC, USA *CDR Monica A. Lutgendorf, MC, USN LCDR Donovan L Mabe, MC, USN COL James Mancuso, MC, USA CDR Frank Margaron, MC, USN Col Leslie Matesick, MC, USAF *CAPT Ryan C. Maves, MC, USN Mr. Greg McFarland, PE CDR Sean Mckay, MC, USN MAJ Michael McMahon, MC, USA Maj Krista Mehlhaff, MC, USAF Maj Eric Meyer, MC, USAF COL Stephanie Meyer, USA Dr. Sonia Miller, RN PhD, CIC, FAPIC CDR Kelly Mokay, USN CAPT Joshua C. Morganstein, USPHS Mr. Art Murnan, PE COL Clinton K. Murray, MC, USA Ms. Lori Naumcheff Col Thomas C. Newton, MC, USAF CDR Phung Thien Nguyen, USN</p>	<p>Lt Col Jason F. Okulicz, MC, USAF COL David Olson, DC, USA COL Neil Page, MC, USA COL Jeremy C. Pamplin, MC, USA LDCR Melissa Parkes, USN *LTC P. Gabriel Peterson, MC, USA LTC Wylan C. Peterson, MC, USA Ms. Rhonda Podojil, DHA Mr. Joseph Procaccino, JD, MFS Lt Col Jamie Rand, MC, USAF MAJ Tyler Reese, MC, USA *LTC Sean Reilly, MC USA CAPT Robert Ricca, MC, USN COL Michael A. Robertson, USA Daniel J. Ross, MD, DDS Col James B. Sampson, MC, USAF Lt Col Valerie Sams, MC, USAF *COL Dennis Sarmiento, MC, USA LDCR David C. Shih, MC, USN LTC Robert Shih, MC, USA LTC Alexander Shilman, MSC, USA *Bich-Thuy T. Sim, MD Maj Erica Simon, MC, USAF *COL William Smith, MC, USA *LCDR Scott Snyder, MC, USN Lt Col Derek Sorensen, MC, USAF *Col Barton C. Staat, MC, USAF CPT Bryan Stepanenko, MC, USA *Lt Col Ian Stewart, MC, USAF Mr. Randy W. Stone, JD *Maj Matthew Streitz, MC, USAF Col Richard Strilka, MC, USAF Maj Mary Stuever, MC, USAF *Col Deena E. Sutter, MC, USAF *COL Timothy L. Switaj, MC, USA *Lt Col Bryan D. Szalwinski, MC, USAF MAJ Matthew Timlin, MC, USA Mary C. Vance, MD, MSc TSgt Alysha Waite, USAF COL Wendi M. Waits, MC, USA Maj Robert J. Walter, MC, USAF CDR Lauren A. Weber, MC, USN James C. West, MD Mr. Bryan T. Wheeler, JD CAPT David Whittaker, MC, USN Col J. Frederick Wiedenhoefer, MC, USAF Alison C. Wiesenthal, MD Maj Thomas J. Wilson, MC, USAF *CAPT Gordon Wisbach, MC USN MAJ Amanda L. Wolfe, USA *Col Leslie Wood, MC, USAF COL Gary H. Wynn, MC, USA MAJ Daniel Yourk, MC, USA Col Heather Yun, MC, USAF CAPT Luke Zabrocki, MC, USN Mr. Michael J. Zleit, JD</p>